



US00PP13507P2

(12) **United States Plant Patent**  
**Gerdts et al.**(10) **Patent No.:** US PP13,507 P2  
(45) **Date of Patent:** Jan. 21, 2003

(54) PEACH TREE NAMED 'BURPEACHTEN'

(75) Inventors: **Timothy J. Gerdts**, Kingburg, CA (US); **John K. Slaughter**, Clovis, CA (US)(73) Assignee: **The Burchell Nursery, Inc.**, Oakdale, CA (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: 10/035,411

(22) Filed: Dec. 28, 2001

(51) Int. Cl.<sup>7</sup> ..... A01H 5/00

(52) U.S. Cl. .... Plt./197

(58) Field of Search ..... Plt./197

Primary Examiner—Bruce R. Campell

Assistant Examiner—Susan B. McCormick

(74) Attorney, Agent, or Firm—Wells St. John P.S.

(57) **ABSTRACT**

A new and distinct variety of peach tree (*Prunus persica*), which is denominated varietally as 'Burpeachten', and produces an attractively colored yellow-fleshed clingstone peach which is mature for harvesting and shipment approximately June 13 to June 22 under ecological conditions prevailing in the San Joaquin Valley of Central California.

**1 Drawing Sheet****1****BACKGROUND OF THE NEW VARIETY**Botanical classification: *Prunus persica*.

Variety denomination 'Burpeachten'.

The present invention relates to a new, novel and distinct variety of peach tree, *Prunus persica*, which has been denominated varietally as 'Burpeachten'. The present variety of peach tree resulted from an on-going program of fruit and nut tree breeding. The purpose of this program is to improve the commercial quality of deciduous fruit and nut varieties and rootstocks by creating and releasing promising selections of prunus, malus and regia species. To this end we make both controlled and hybrid cross pollinations each year in order to produce seedling populations from which improved progenies are evaluated and selected.

The seedling 'Burpeachten' was originated by us from a population of seedlings grown in our experimental orchards located near Fowler, Calif. The seedlings, grown on their own roots, were the result of a controlled cross of the peach tree 'Rich Lady' (U.S. Plant Pat. No. 7,290) which was used as the seed parent, and the peach tree 'Mayfire' (non-patented) which was used as the pollen parent. One seedling, which is the present variety, exhibited especially desirable characteristics and was marked for subsequent observation. After the 1997 growing season, the new variety was selected for advanced evaluation and repropagation.

**ASEXUAL REPRODUCTION**

Asexual reproduction of the new and distinct variety of peach tree was accomplished by budding the present variety to 'Nemaguard' Rootstock (non-patented). This was performed by us in our experimental orchard located near Fowler, Calif. Subsequent evaluations have shown those asexual reproductions run true to the original tree. All characteristics of the original tree and its fruit were established, and appeared to be transmitted through succeeding asexual propagations.

**SUMMARY OF THE VARIETY**

'Burpeachten' is a new and distinct variety of peach tree, which is of medium large size, and which has vigorous

**2**

growth, and is a regular and productive bearer of large, firm, yellow flesh, and semi-freestone fruit with good flavor and eating quality. The tree has a medium-chilling requirement of approximately 600 hours. The tree also produces relatively uniformly sized fruit throughout the tree with a high degree of red skin coloration, and firm flesh. The fruit appears to have good handling and shipping quality. Still further, the 'Burpeachten' peach tree bears fruit that is ripe for commercial harvesting and shipment on approximately June 13 to June 22. In comparison to the seed parent the 'Rich Lady' Peach tree (U.S. Plant Pat. No. 7290), the new variety ripens about 5 or more days earlier.

**15 BRIEF DESCRIPTION OF THE DRAWING**

The accompanying drawing which is provided is a color photograph of the present variety. It depicts a characteristic twig bearing typical leaves; several leaves showing both the dorsal and ventral coloration; and two mature fruit and a third fruit that has been bisected along the equatorial plane and displaying an apical view of the fruit, flesh and pit intact. The external coloration of the fruit is shown sufficiently matured for harvesting and shipment. The colors are as nearly true as is reasonably possible in a color photographic representation of this type. Due to chemical development, processing and printing, the leaves, pit and fruit depicted in this photograph may or may not be accurate when compared to the actual specimen. For this reason, future color references should be made to the color plates (Royal Horticultural Society) and descriptions provided hereinafter.

**DETAILED DESCRIPTION**

Referring more specifically to the pomological details of this new and distinct variety of peach tree, the following has been observed in the fifth fruiting season under the ecological conditions prevailing at orchards located near the town of Fowler, county of Fresno, state of California. All major color code designations are by reference to The R.H.S. Colour Chart (Fourth Edition) provided by The Royal Horticultural Society of Great Britain.

## Tree:

**Size.**—Generally. — Considered medium as compared to other common commercial peach cultivars ripening in the mid season of maturity. The tree was pruned to a height of approximately 304.8 cm to 365.7 cm at maturity.

**Vigor.**—Moderately vigorous. The present variety grew from about 121.92 cm to 182.8 cm in height during the first growing season. The variety was pruned to a height of approximately 106.7 cm in the first dormant season and primary scaffolds are selected for the desired tree structure.

**Productivity.**—Productive. Fruit set varies from about 1.5 to several times more than the desired crop load. Fruit set is spaced by thinning to develop into the desired market sized fruit. Number of fruit set varies with climatic conditions and prevailing cultural practices during the bloom period and is therefore not distinctive of the variety.

**Bear.**—Regular. Fruit set has been heavy and thinning was necessary during the past 4 years.

**Form.**—Upright, and pruned to a vase shape.

**Density.**—Medium dense. It have been discovered that pruning the branches from the center of the tree to obtain a resulting vase shape allows for proper air movement and appropriate amounts of sunlight to enhance fruit color and renewal of fruiting wood throughout the tree.

**Hardiness.**—The present tree was grown and evaluated in USDA Hardiness Zone 9. Winter chilling requirements are approximately 600 hours below 7.0 degrees C. The variety appears to be hardy under typical Central San Joaquin Valley climatic conditions.

## Trunk:

**Diameter.**—Approximately 16.0 cm in diameter when measured at a distance of approximately 15.24 cm above the soil level, at the end of the fourth growing season.

**Bark texture.**—Considered moderately rough, with numerous folds of papery scarf skin being present.

**Lenticels.**—Numerous flat, oval lenticels are present. The lenticels range in size from approximately 2.0 to 6.0 millimeters in width, and from 1 to 2 millimeters in height.

**Lenticel color.**—Considered an Orange Brown. (RHS Greyed-Orange N172 A). Bark Coloration. — Variable, but it is generally considered to be a grey-brown (RHS Greyed-Orange Group 177 A).

## Branches:

**Size.**—Considered medium for the variety.

**Diameter.**—Average as compared to other varieties. The branches have a diameter of about 75.0 millimeters when measured during the third year after grafting.

**Surface texture.**—Average, and appearing furrowed on wood which is several years old.

**Crotch angles.**—Primary branches considered variable between about 45 to 49 degrees from the horizontal axis. This characteristic is not considered distinctive of the variety however.

**Current season shoots.**—Surface texture — Substantially glabrous.

**Internode length.**—Approximately 2.0 to 2.2 cm.

**Color of mature branches.**—Medium brown, (RHS Grey Brown Group N199C).

**Current seasons shoots.**—Color. Light green, (RHS Yellow Green Group 144 D). The color of new shoot tips is considered a bright and shiny green (RHS Green Group 134 C).

## Leaves:

**Size.**—Considered medium for the species. Leaf measurements have been taken from vigorous upright current season growth at approximately mid-shoot.

**Leaf length.**—Approximately 140 to 170 millimeters.

**Leaf width.**—Approximately 32 to 35 millimeters.

**Leaf base shape.**—Slightly oblique relative to the leaf longitudinal axis.

**Leaf form.**—Lanceolate.

**Leaf tip form.**—Acuminate.

**Leaf color.**—Dark green, approximately (RHS Green Group 137 A).

**Leaf texture.**—Glabrous.

**Lower surface.**—Medium green, (RHS Yellow Green Group 147)

**Leaf venation.**—Pinnately veined.

**Mid-Vein.**—Color. — Light yellow green, (RHS Yellow Green Group 151 A).

**Leaf margins.**—Slightly undulating. Form. — Considered crenate, occasionally doubly crenate. Uniformity. — Considered generally uniform. **Leaf petioles.**—Size. — Considered medium long to long. Length. — 9.0 to about 12.0 mm. Diameter. — 1.5 to about 2.5 mm. Color. — Pale green, (RHS Yellow Green Group 151 A).

**Leaf glands.**—Size. — About 1.0 mm in height and about 1.0 to 2.0 mm in width. Numbers. — Generally one per side, occasionally two per side. Type. — Reniform, and considered reasonably unappressed to the petiole margin. Color. — Greenish brown (RHS Yellow Green Group 152 B).

**Leaf stipules.**—Size. — Medium large for the variety. Number. — Typically 2 per leaf bud, and up to 6 per shoot tip. Form. — Lanceolate in form and having a serrated margin. Color. — Green, (RHS Green Group 132 A) when young, but graduating to a yellow-brown color (RHS Grey Orange group 177 A) with advancing senescence. The stipules are considered to be early deciduous.

## Flowers:

**Flower buds.**—Generally — The floral buds are considered to be large, about 15 millimeters wide; and about 21 millimeters long; conic in form; and slightly appressed relative to the bearing shoot.

**Flower Buds.**—Color — The bud scales are gray-brown, (approximately RHS Greyed Purple Group 183 A). The buds are considered hardy under typical central San Joaquin Valley climatic conditions.

**Hardiness.**—No winter injury has been noted during the last several years of evaluation in the Central San Joaquin Valley. The current variety has not been intentionally subjected to drought or heat stress and therefore this information is not available.

**Date of first bloom.**—Feb. 22, 1999.

**Blooming time.**—Considered relatively early in comparison to other commercial peach cultivars commonly grown in the Central San Joaquin valley. Date of full bloom was observed on Mar. 1, 1999. The date of bloom varies slightly with climatic conditions and cultural practices.

**Duration of bloom.**—Approximately 7 days.

**Flower type.**—The variety is considered to have a showy type flower.

*Flower size.*—Flower diameter at full bloom is approximately 35.0 to 41.0 millimeters.

*Bloom quantity.*—Considered abundant.

*Flower bud frequency.*—Normally 1 to 2 appear per node.

*Petal size.*—Generally. — Considered large for the species. Length. — Approximately 18.0 to 21.0 millimeters. Width. — Approximately 18.0 to 20.0 millimeters.

*Petal form.*—Broadly ovate.

*Petal count.*—Nearly always 5.

*Petal texture.*—Glabrous.

*Petal color.*—Light pink when young, (RHS Red Purple Group 63 C), and darkening with advancing senescence and exposure to sunlight to a medium to dark pink (RHS Red Purple Group 64 D).

*Fragrance.*—Slight.

*Petal claw.*—Form. — The claw is considered truncate and has a medium-large size when compared to other varieties. Length. — Approximately 8.0 to 10.0 millimeters. Width. — Approximately 6.0 to 8.0 millimeters.

*Petal margins.*—Generally considered variable, from nearly smooth, to moderately undulate and ruffled, especially apically.

*Petal apex.*—Generally — The petal apices appear slightly domed.

*Flower pedicel.*—Length. — Considered medium-long, and having an average length of approximately 3.0 to 5.0 millimeters. Diameter. — Considered average, approximately 2.0 millimeters. Color. — A medium Green (RHS — Grey Brown Group N199 D).

*Floral nectaries.*—Color. — A dull orange-gold (RHS Greyed Orange Group 172 B).

*Calyx.*—Surface Texture. — Generally glabrous. Color. — A dull red, (approximately RHS Greyed Purple Group 183 A).

*Sepals.*—Surface Texture. — The surface has a short, fine, wooly texture. Size. — Average, and ovate in form. Color. — A dull red, (approximately RHS Greyed-Red Group 178 A).

*Anthers.*—Generally. — Average, to above average in length. Color. — Red to reddish-orange dorsally, (approximately RHS Greyed Purple Group 179 A).

*Pollen production.*—Pollen is abundant, and has a yellow color, (approximately RHS Yellow Orange Group 17 B).

*Filaments.*—Size. — Variable in length, approximately 12.0 to 14.0 millimeters in length. Color. — White, (RHS Red Purple Group 62 D).

*Pistil.*—Number. — Usually 1, rarely 2. Generally. — Average in size. Length. — Approximately 14.0 to 16.0 millimeters including the ovary. Color. — Considered a very pale green, (approximately RHS Yellow Green Group 151 D). Surface Texture. — The variety has a long, pubescent pistil.

*Fruit:*

*Maturity when described.*—Firm ripe condition (shipping ripe). Date of first picking. — Jun. 13, 1999. Date of last picking. — Jun. 22, 1999. The date of harvest varies slightly with climatic conditions.

*Size.*—Generally — Medium large, and considered uniform.

*Average cheek diameter.*—Approximately 74.0 to 76.0 millimeters.

*Average axial diameter.*—Approximately 71.0 to 73.0 millimeters.

*Typical weight.*—Approximately 234.0 grams. This is highly dependent upon cultural practices and therefore is not distinctive of the variety.

*Fruit form.*—Generally — Moderately oblate. The fruit is generally uniform in symmetry.

*Fruit suture.*—Shallow, slightly lipped, and extending from the base to apex. No apparent callousing or stitching exists along the suture line.

*Suture.*—Color — This appears to be a yellow to golden yellow background color, (approximately RHS Yellow Orange Group 22 B) and additionally having some red-orange coloration (approximately RHS Red Group 45 A).

*Ventral surface.*—Form — Slightly indented.

*Apex.*—Varies from rounded to having a slight apical point.

*Base.*—Retuse.

*Stem cavity.*—Rounded too slightly elongated in the suture plane. Average depth of the stem cavity is about 1.25 cm. Average width is about 2.51 cm.

*Fruit skin.*—Thickness. — Considered medium in thickness, and tenacious to the flesh. Texture. — Medium, and having a slight pubescence. Taste. — Non-astringent. Tendency to crack. — None observed.

*Color.*—Blush Color. — This red blush color is variable from a dark to a lighter red, (approximately RHS Greyed-Purple Group 187 A to 187 C). Ground Color. — Pale yellow, (approximately RHS Yellow 11 B).

*Fruit stem.*—Medium in length, approximately 8.0 to about 9.0 millimeters. Diameter. — Approximately 2.0 to 3.0 millimeters. Color. — Pale yellow-green (approximately RHS Yellow een Group 145 B).

*Flesh.*—Ripens. — Evenly. Texture. — Considered firm, and dense. Fibers. — Few, small, and tender ones are found. Aroma. — Very slight. Eating Quality. — Good. Flavor. — Considered sweet and mildly acidic. The flavor is considered both pleasant and balanced. Juice. — Moderate. Brix. — About 11.0 degrees. This value varies slightly with the number of fruit per tree; prevailing cultural practices; and surrounding climatic conditions. Flesh Color. — Pale yellow, (approximately RHS Yellow Group 10 A). With increasing maturity red flecking (approximately RHS Red Group 43 B) can be found.

*Stone:*

*Type.*—Clingstone.

*Size.*—Considered medium for the variety.

*Length.*—Average, about 22.0 to about 24.0 millimeters.

*Width.*—Average, about 21.0 to about 22.0 millimeters.

*Diameter.*—Average, about 18.0 to about 19.0 millimeters.

*Form.*—Obovoid.

*Base.*—The stone is usually rounded, but may vary from rounded to straight.

*Apex.*—Shape. — The stone apex is raised and has an acute, short tip.

*Stone surface.*—Surface Texture — Irregularly furrowed toward the apex, and pitted toward the base. The stone exhibits substantial pitting laterally. Substantial grooving over the apical shoulders is evident. Surface pitting is prominent generally, and more

frequently, it is present basally. One long furrow on each side of suture will normally be found. Ridges.—Surface texture varies from sharp to rounded. Ventral Edge.—Width—Considered medium, and having a dimension of approximately 5 to about 6.5 millimeters at the mid-suture. The wings are most prominent over the basal area. Dorsal Edge.—Shape.—Full, heavily grooved, and having jagged edges. The dorsal edge is moderately eroded over the apical shoulder.

*Stone color.*—The color of the dry stone is a light to medium brown, (approximately RHS Orange Red Group 34 C).

*Tendency to split.*—Few but very infrequent splitting was noted.

*Kernel.*—Size.—Length—18.0 mm. Width—12.0 mm. Thickness—3.0 millimeters. Form.—Obovoid. Pellicle.—Pubescent. Color.—(RHS Greyed Brown Group N199 C).

*Use.*—The subject variety 'Burpeachten' is considered to be a Peach tree of the mid-season maturity, and which produces fruit which are considered very firm, attractively colored, and which are useful for both local and long distance shipping.

*Keeping quality.*—Excellent. Fruit has stored well up to 21 days after harvest at 1.0 degree Celsius.

*Shipping quality.*—Good. Fruit showed minimal bruising of the flesh or skin damage after being subjected to normal harvesting and packing procedures.

*Resistance to insects and disease.*—No particular susceptibilities were noted. The present variety has not been tested to detect for any susceptibilities or resistances to any known plant and/or fruit diseases.

Although the new variety of peach tree possesses the described characteristics when grown under the ecological conditions prevailing near Fowler, Calif. in Central part of the San Joaquin Valley of California, it will be understood that variations of the usual magnitude and characteristics incident to changes in growing conditions, fertilization, pruning, pest control and horticultural management are to be expected.

Having thus described and illustrated our new variety peach tree, what we claim is new and desire to secure by Plant Letters Patent is:


1. A new and distinct variety of peach tree, substantially as illustrated and described, and which is characterized principally as novelty by producing an attractively colored yellow-fleshed clingstone peach which is mature for harvesting and shipment approximately June 13 to June 22 under the ecological conditions prevailing in the San Joaquin Valley of Central California.

\* \* \* \* \*

**U.S. Patent**

**Jan. 21, 2003**

**US PP13,507 P2**



UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : PP 13,507 P2  
DATED : January 21, 2003  
INVENTOR(S) : Timothy J. Gerdts et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:


Column 4,  
Line 19, replace “Group 147” with -- Group 147B). --.

Column 6,  
Line 34, replace “RHS Yellow een Group” with -- RHS Yellow Green Group --.

Column 8,  
Line 11, replace “in Central part” with -- in the Central part --.

Signed and Sealed this

Fifth Day of August, 2003



JAMES E. ROGAN  
*Director of the United States Patent and Trademark Office*