



US00PP13574P2

(12) **United States Plant Patent**
Dümmen

(10) **Patent No.: US PP13,574 P2**
(45) **Date of Patent: Feb. 18, 2003**

(54) **GERANIUM PLANT NAMED ‘DUEBART’**
(75) Inventor: **Marga Dümmen**, Rheinberg (DE)
(73) Assignee: **Dümmen Jungpflanzen GbR**,
Rheinberg (DE)
(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.
(21) Appl. No.: **10/080,605**
(22) Filed: **Feb. 22, 2002**

(51) **Int. Cl.⁷** **A01H 5/00**
(52) **U.S. Cl.** **Plt./332**
(58) **Field of Search** **Plt./332**
Primary Examiner—Kent L. Bell
(74) *Attorney, Agent, or Firm*—C. A. Whealy
(57) **ABSTRACT**
A new and distinct cultivar of Ivy Geranium plant named
‘Duebart’, characterized by its compact, upright and out-
wardly spreading plant habit; freely basal branching habit;
freely flowering habit; and dark pink-colored flowers.

1 Drawing Sheet

1

**BOTANICAL CLASSIFICATION/CULTIVAR
DENOMINATION**

Pelargonium peltatum cultivar ‘Duebart’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of Ivy Geranium plant, botanically known as *Pelargonium*
peltatum, and hereinafter referred to by the name ‘Duebart’.
The new Ivy Geranium is a product of a planned breeding
program conducted by the Inventor in Rheinberg, Germany.
The objective of the breeding program was to develop new
Ivy Geraniums with attractive flower and foliage colors.
The new Ivy Geranium originated from a cross made by
the Inventor of a proprietary selection of *Pelargonium*
peltatum identified as code number N-12-001, not patented,
as the female, or seed, parent with a proprietary selection of
Pelargonium peltatum identified as code number F-23-09,
not patented, as the male, or pollen, parent. The cultivar
‘Duebart’ was discovered and selected by the Inventor as a
flowering plant within the progeny from this cross in a
controlled environment in Rheinberg, Germany.
Asexual reproduction of the new cultivar by terminal
vegetative cuttings taken at Rheinberg, Germany has shown
that the unique features of this new Ivy Geranium are stable
and reproduced true to type in successive generations of
asexual reproduction.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Duebart’.
These characteristics in combination distinguish ‘Duebart’
as a new cultivar and distinguish it from other known Ivy
Geranium cultivars:
1. Compact, upright and outwardly spreading plant habit.
2. Freely basal branching habit.
3. Freely flowering habit.
4. Dark pink-colored flowers.

Compared to plants of the female parent, the selection
N-12-001, plants of the new Ivy Geranium are more
compact, have smaller leaves and have brighter pink-colored
petals.

2

Compared to plants of the male parent, the selection
F-23-09, plants of the new Ivy Geranium are more compact,
have smaller leaves and larger flower umbels.
The new Ivy Geranium can be compared to plants of the
cultivar ‘Guimongol’, disclosed in U.S. Plant Pat. No. 9,352.
In side-by-side comparisons conducted in Rheinberg,
Germany, plants of the new Ivy Geranium differed from
plants of the cultivar ‘Guimongol’ in the following charac-
teristics:
1. Plants of the new Ivy Geranium were more compact
with shorter internodes and shorter peduncles than
plants of the cultivar ‘Guimongol’.
2. Plants of the new Ivy Geranium had longer leaves than
plants of the cultivar ‘Guimongol’.
3. Leaves of the new Ivy Geranium did not have a distinct
zonation pattern than leaves of plants of the cultivar
‘Guimongol’ did have a distinct zonation pattern.
4. Plants of the new Ivy Geranium had larger flower
umbels than plants of the cultivar ‘Guimongol’.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the
overall appearance of the new cultivar, showing the colors as
true as it is reasonably possible to obtain in colored repro-
ductions of this type. Flower and foliage colors in the
photograph may differ slightly from the color values cited in
the detailed botanical description which accurately describe
the colors of the new Ivy Geranium.
The photograph comprises a side perspective view of
typical flowering plants of ‘Duebart’ grown in a container.

DETAILED BOTANICAL DESCRIPTION

The cultivar ‘Duebart’ has not been observed under all
possible environmental conditions. The phenotype may vary
somewhat with variations in environment, such as tempera-
ture and light intensity, without, however, any variance in
genotype.
The aforementioned photographs, following observations
and measurements describe plants grown in Rheinberg,
Germany during the spring under commercial practice in a
glass-covered greenhouse with day and night temperatures
about 18° C. and light levels about 4,500 foot-candles.
Plants were grown with three plants per container. Plants
were about eight weeks from unrooted cuttings when the

photographs and the detailed botanical description were taken.

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Pelargonium peltatum* cultivar 'Duebart'.

Parentage:

Female parent.—Proprietary selection of *Pelargonium peltatum* identified as code number N-12-001, not patented.

Male parent.—Proprietary selection of *Pelargonium peltatum* identified as code number F-23-09, not patented.

Propagation:

Type cutting.—Terminal vegetative cuttings.

Time to initiate roots.—Summer: About 5 days at 20° C.

Winter: About 7 days at 20° C.

Time to develop roots.—Summer: About 21 days at 20° C. Winter: About 28 days at 20° C.

Root description.—Fine; freely branching.

Plant description:

General appearance.—Compact, upright and outwardly spreading plant habit, rounded; densely foliated.

Growth and branching habit.—Moderately vigorous. Freely basal branching, about four lateral branches per plant.

Plant height.—About 16 cm.

Plant width.—About 11 cm.

Lateral branches.—Length: About 13 cm. Internode length: About 2.25 cm. Texture: Smooth. Color: 144A.

Foliage description.—Arrangement: Alternate, single. Quantity of leaves per lateral branch: About five. Length: About 7 cm. Width: About 7.6 cm. Shape: Reniform. Apex: Acute. Base: Peltate. Margin: Crenate. Venation pattern: Palmate. Texture, upper and lower surfaces: Smooth. Color: Young and fully expanded foliage, upper surface: 137A; no distinct zonation pattern. Young and fully expanded foliage, lower surface: 144A. Venation, upper and lower surfaces: 144A. Petiole: Length: About 7.4 cm. Diameter: About 2 mm. Color, upper and lower surfaces: 144A.

Flower description:

Flower arrangement.—Dark pink-colored flowers arranged in rounded hemispherical umbels arising from apical leaf axils. Umbels displayed above the foliage on upright peduncles. Flowers rounded in form. Umbels persistent, flowers not persistent. Flowers not fragrant.

Quantity of flowers.—Freely flowering; at full flower, plants have about four open and developing umbels with about 15 flowers per umbel.

Flowering season.—Flowering continuous spring through summer.

Flower longevity.—Flowers last about five to seven days on the plant.

Umbel size.—Height: About 7 cm. Diameter: About 9.5 cm.

Flower size.—Diameter: About 4.25 cm. Depth (height): About 2 cm.

Flower buds.—Length: About 1.3 cm. Diameter: About 8 mm. Shape: Ovoid. Color: 144A.

Petals.—Quantity per flower: About five. Length: About 2.8 cm. Width: About 1.6 cm. Shape: Oblongate. Apex: Rounded. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Smooth. Color: When opening and fully opened, upper surface: 68B; towards base, 53A; color fading to 66D with subsequent development. When opening and fully opened, lower surface: 68D; towards base, 53A. Venation, upper surface: 68B. Venation, lower surface: 68D.

Petaloids.—Quantity per flower: About five. Length: About 2.8 cm. Width: About 1.6 cm. Shape: Oblongate. Apex: Rounded. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Smooth. Color: When opening and fully opened, upper surface: 68B; towards base, 53A; color fading to 66D with subsequent development. When opening and fully opened, lower surface: 68D; towards base, 53A. Venation, upper surface: 68B. Venation, lower surface: 68D.

Sepals.—Quantity per flower: Five, arranged in a single whorl. Length: About 1.5 cm. Width: About 4 mm. Shape: Elongated, tapering. Apex: Apiculate. Margin: Entire. Texture, upper and lower surfaces: Smooth. Color, upper and lower surfaces: 144A.

Peduncle (umbel stem).—Length: About 13 cm. Diameter: Less than 1 cm. Angle: Erect. Strength: Moderately strong. Texture: Smooth. Color: 144A.

Pedicel (individual flower stem).—Length: About 2.8 cm. Diameter: About 1 mm. Angle: Erect. Strength: Moderately strong. Texture: Pubescent. Color: 178A.

Reproductive organs.—Androecium: Anther quantity per flower: About five. Anther length: About 3 mm. Anther shape: Ovate. Anther color: 60A. Pollen amount: Moderate. Pollen color: 28A. Gynoecium: Pistil quantity per flower: One. Pistil length: About 1.3 cm. Stigma shape: Five-parted, star-shaped. Stigma color: 60A. Style length: About 6 mm. Style color: 2D. Ovary color: 144A.

Seed/fruit.—Development of seeds and fruit have not been observed.

Disease/pest resistance: Plants of the new Ivy Geranium have not been observed to be resistant to pathogens and pests common to Ivy Geraniums.

It is claimed:

1. A new and distinct cultivar of Ivy Geranium plant named 'Duebart', as herein illustrated and described.

* * * * *

