



US00PP13636P29

(12) **United States Plant Patent**
Dümmen

(10) **Patent No.: US PP13,636 P2**
(45) **Date of Patent: Mar. 11, 2003**

(54) **PETUNIA PLANT NAMED
‘DUESURIMROVEIN’**

(75) Inventor: **Marga Dümmen**, Rheinberg (DE)

(73) Assignee: **Dümmen Jungpflanzen GbR**,
Rheinberg (DE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/104,368**

(22) Filed: **Mar. 22, 2002**

(51) **Int. Cl.⁷ A01H 5/00**

(52) **U.S. Cl. Plt./356**

(58) **Field of Search Plt./356**

Primary Examiner—Bruce R. Campell
Assistant Examiner—Anne Marie Grünberg
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of Petunia plant named
‘Duesurimrovein’, characterized by its upright and out-
wardly spreading plant habit; freely basal branching; and
single flowers that are dark pink in color with red purple-
colored venation.

1 Drawing Sheet

1

**BOTANICAL CLASSIFICATION/CULTIVAR
DESIGNATION**

Petunia×*hybrida* cultivar Duesurimrovein.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of Petunia plant, botanically known as *Petunia*×*hybrida*,
and hereinafter referred to by the cultivar name Duesurim-
rovein.

The new Petunia is a product of a planned breeding
program conducted by the Inventor in Rheinberg, Germany.
The objective of the breeding program is to create new freely
branching Petunias with large flowers and attractive flower
coloration.

The new Petunia originated from a cross pollination made
by the Inventor in 1997 of a proprietary Petunia selection
identified as code number 94-211, not patented, as the
female, or seed parent, with a proprietary Petunia selection
identified as code number F-18-22, not patented, as the male,
or pollen parent. The new Petunia was selected as a single
plant from the resulting progeny by the Inventor in
Rheinberg, Germany in 1998, on the basis of its attractive
flower color.

Asexual reproduction of the new cultivar by terminal
vegetative cuttings taken in Rheinberg, Germany since
1999, has shown that the unique features of this new Petunia
are stable and reproduced true to type in successive genera-
tions.

SUMMARY OF THE INVENTION

Plants of the cultivar Duesurimrovein have not been
observed under all possible environmental conditions. The
phenotype may vary somewhat with variations in environ-
ment such as temperature and light intensity without,
however, any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of
‘Duesurimrovein’. These characteristics in combination dis-
tinguish ‘Duesurimrovein’ as a new and distinct cultivar:

1. Upright and outwardly spreading plant habit.
2. Freely basal branching.

2

3. Single flowers that are dark pink in color with red
purple-colored venation.

Compared to plants of the female parent, the selection
94-211, plants of the new Petunia have larger flowers and
differ in flower color. Compared to plants of the male parent,
the selection F- 18-22, plants of the new Petunia have larger
flowers and differ in flower color.

Plants of the new Petunia can be compared to plants of the
cultivar Surfinia Pink Vein, not patented. In side-by-side
comparisons conducted in Rheinberg, Germany, plants of
the new Petunia differed from plants of the cultivar Surfinia
Pink Vein in the following characteristics:

1. Plants of the new Petunia had longer internodes than
plants of the cultivar Surfinia Pink Vein.
2. Plants of the new Petunia had larger leaves with longer
petioles than plants of the cultivar Surfinia Pink Vein.
3. Plants of the new Petunia had larger flowers than plants
of the cultivar Surfinia Pink Vein.
4. Flowers of plants of the new Petunia were darker in
color than flowers of plants of the cultivar Surfinia Pink
Vein.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the
overall appearance of the new cultivar, showing the colors as
true as it is reasonably possible to obtain in colored repro-
ductions of this type. Colors in the photograph differ slightly
from the color values cited in the detailed botanical descrip-
tion which accurately describe the colors of the new Petunia.

The photograph comprises a top perspective view of
typical flowers and leaves of ‘Duesurimrovein’.

DETAILED BOTANICAL DESCRIPTION

The cultivar Duesurimrovein has not been observed under
all possible environmental conditions. The phenotype may
vary somewhat with variations in environment such as
temperature and light intensity without, however, any vari-
ance in genotype.

The aforementioned photographs, following observations
and measurements describe plants grown during the spring
in Rheinberg, Germany, under commercial practice in a

glass-covered greenhouse. Plants were about 16 weeks from cuttings and were grown in containers. During the production of the plants, day and night temperatures averaged 18° C. and light levels were about 4,500 lux. Plants were pinched once about three weeks after planting.

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Petunia*×*hybrida* cultivar Duesurimrovein.

Parentage:

Female parent.—Proprietary *Petunia*×*hybrida* selection identified as code number 94-211, not patented.

Male parent.—Proprietary *Petunia*×*hybrida* selection identified as code number F-18-22, not patented.

Propagation:

Type cutting.—Terminal vegetative cuttings.

Time to initiate roots.—Summer: About 7 days at 20° C. Winter: About 10 days at 20° C.

Time to develop roots.—Summer: About 21 days at 20° C. Winter: About 28 days at 20° C.

Root description.—Fine, fibrous, white in color.

Rooting habit.—Freely branching.

Plant description:

Form.—Annual flowering plant; indeterminate; upright and outwardly spreading; eventually cascading. Moderately vigorous.

Usage.—Appropriate for hanging baskets, window boxes, patio containers and landscape applications.

Plant height.—About 30 cm.

Plant diameter.—About 26 cm.

Branching habit.—Freely basal branching with about six lateral branches per plant.

Lateral branches.—Length: About 27 cm. Diameter: About 3 mm. Internode length: About 2.4 cm. Texture: Pubescent. Color: 144B.

Foliage description.—Arrangement: Before flowering, alternate; after flowering, opposite; simple. Quantity per lateral branch: About 15. Length: About 7.9 cm. Width: About 4 cm. Shape: Roughly spatulate. Apex: Obtuse. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Pubescent; leathery. Venation pattern: Pinnate. Color: Young and fully expanded foliage, upper surface: 137B. Young and fully expanded foliage, lower surface: 137D. Venation, upper and lower surfaces: 144B. Petiole length: About 8 mm. Petiole diameter: About 2 mm. Petiole color: 144D.

Flower description:

Flower type and habit.—Single salverform flowers; flowers face mostly upward or outward; axillary; about five flowers and flower buds per lateral branch.

Natural flowering season.—Long day responsive; spring until frost in the autumn; flowering continuous. Plants start flowering about nine weeks after planting. Flowers persistent.

Flower longevity on the plant.—About 4 days; flowers persistent.

Fragrance.—None detected.

Flower size.—Diameter: About 7 cm. Tube length: About 3 cm. Throat diameter, distal end: About 1.8 cm. Tube diameter, proximal end: About 4 mm.

Flower buds.—Length: About 4.5 cm. Diameter: About 1 cm. Shape: Ovoid. Color: 63B.

Corolla.—Quantity/arrangement: Five fused petals; funnellform. Petal length from throat: About 3.5 cm. Petal width: About 3.5 cm. Petal shape: Roughly spatulate. Petal apex: Obtuse. Petal margin: Entire; slightly ruffled. Petal texture, upper and lower surfaces: Smooth, velvety. Petal color: When opening, upper and lower surfaces: 66C. Fully opened, upper and lower surfaces: 66C; flower color does not fade with subsequent development. Flower throat (inside): 78A. Flower tube (outside): 82D. Venation, upper petal surface: 67A. Venation, lower petal surface: 144B. Venation, throat: 79D. Venation, tube: 144B.

Sepals.—Arrangement/appearance: Single whorl of five sepals fused at base, star-shaped. Length: About 1 cm. Width: About 2.5 mm. Shape: Strap-like; elongate. Apex: Rounded. Margin: Entire. Texture, upper and lower surfaces: Pubescent. Color: Upper surface: 137B. Lower surface: 137D.

Peduncles.—Length: About 2.8 cm. Width: About 1.25 mm. Angle: Erect to slightly bent. Strength: Moderately strong. Texture: Pubescent. Color: 144A.

Reproductive organs.—Stamens: Quantity per flower: About five. Anther shape: Ovoid. Anther length: About 3 mm. Anther color: 8D. Pollen amount: Abundant. Pollen color: 8D. Pistils: Quantity per flower: One. Pistil length: About 1.9 cm. Style length: About 1 cm. Style color: 144C. Stigma shape: Rounded. Stigma color: 144B. Ovary color: 144B.

Seed/fruit.—Seed nor fruit production has not been observed.

Disease/pest resistance: Plants of the new *Petunia* have not been noted to be resistant to pathogens or pests common to *Petunia*.

Temperature tolerance: Plants of the new *Petunia* have been observed to tolerate temperatures from 2 to 38° C.

It is claimed:

1. A new and distinct cultivar of *Petunia* plant named 'Duesurimrovein', as illustrated and described.

* * * * *

