



US00PP13648P29

(12) **United States Plant Patent**  
**Uchneat**

(10) **Patent No.:** **US PP13,648 P2**

(45) **Date of Patent:** **Mar. 11, 2003**

(54) **DOUBLE IMPATIENS PLANT NAMED**  
**'BALOLESAL'**

PP11,549 P \* 10/2000 Cosner et al. .... Plt./317  
PP11,859 P2 \* 5/2001 Knosher ..... Plt./317

(75) Inventor: **Michael S. Uchneat**, St. Charles, IL  
(US)

**OTHER PUBLICATIONS**

(73) Assignee: **Ball FloraPlant, a division of Ball**  
**Horticultural Company**, West  
Chicago, IL (US)

UPOV ROM GTITM computer database, GTI JOUVE  
retrieval software 2002/02, citations for 'Balolesal'.\*

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

\* cited by examiner

(21) Appl. No.: **10/115,052**

*Primary Examiner*—Bruce R. Campell  
*Assistant Examiner*—W C Haas  
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(22) Filed: **Apr. 1, 2002**

(57) **ABSTRACT**

(51) **Int. Cl.**<sup>7</sup> ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./317**

(58) **Field of Search** ..... **Plt./317**

A new and distinct cultivar of Double Impatiens plant named  
'Balolesal', characterized by its compact, upright, outwardly  
spreading and mounded plant habit; freely branching growth  
habit; dark green-colored leaves; freely flowering habit; and  
light red-colored flowers that are fully double and positioned  
above the foliage.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

**1 Drawing Sheet**

PP9,691 P \* 11/1996 Leue ..... Plt./317

**1**

**2**

**BOTANICAL CLASSIFICATION/CULTIVAR**  
**DESIGNATION**

**SUMMARY OF THE INVENTION**

*Impatiens walleriana* cultivar Balolesal.

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of 'Balole-  
sal'. These characteristics in combination distinguish  
'Balolesal' as a new and distinct cultivar:

**BACKGROUND OF THE INVENTION**

1. Compact, upright, outwardly spreading and mounded  
plant habit.
2. Freely branching growth habit.
3. Dark green-colored leaves.
4. Freely flowering habit.
5. Light red-colored flowers that are fully double and  
positioned above the foliage.

The present Invention relates to a new and distinct culti-  
var of Double Impatiens plant, botanically known as *Impa-*  
*tiens walleriana*, and hereinafter referred to by the name  
'Balolesal'.

Plants of the new Impatiens differ primarily from plants of  
the parent selections in flower color.

The new Impatiens is a product of a planned breeding  
program conducted by the Inventor in Elburn, Ill. The  
objective of the breeding program was to develop new  
compact Double Impatiens cultivars with fully double  
flowers, freely branching growth habit, and attractive flower  
and foliage coloration.

The new Impatiens can be compared to the *Impatiens*  
*walleriana* cultivar Salmon Sunrise, disclosed in U.S. Plant  
Pat. No. 9,691. However, in side-by-side comparisons con-  
ducted in West Chicago, Ill. plants of the new Impatiens  
differed from plants of the cultivar Salmon Sunrise in the  
following characteristics:

The new Impatiens originated from a cross-pollination  
made by the Inventor in June, 1999, of a proprietary *Impa-*  
*tiens walleriana* selection identified as 3039C-1, not  
patented, as the female, or seed parent, with a proprietary  
*Impatiens walleriana* selection identified as 3111-1-9, not  
patented, as the male, or pollen parent. The cultivar Balole-  
sal was discovered and selected by the Inventor as a flow-  
ering plant within the progeny of the stated cross-pollination  
in a controlled environment in Elburn, Ill.

1. Plants of the new Impatiens had longer lateral branches  
with longer internodes than plants of the cultivar Salmon  
Sunrise.

2. Plants of the new Impatiens had larger and darker  
green-colored leaves than plants of the cultivar Salmon  
Sunrise.

Asexual reproduction of the new cultivar by terminal  
cuttings taken at Elburn, Ill. since October, 1999, has shown  
that the unique features of this new Impatiens are stable and  
reproduced true to type in successive generations of asexual  
reproduction.

3. Plants of the new Impatiens were more freely flowering  
than plants of the cultivar Salmon Sunrise.

4. Plants of the new Impatiens had larger flowers with  
more petals per flower than plants of the cultivar Salmon  
Sunrise.

5. Flower color of plants of the new *Impatiens* was more orange than flower color of plants of the cultivar Salmon Sunrise.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Impatiens*.

The photograph comprises a side perspective view of multiple flowering plants of 'Balolesal' in a container.

#### DETAILED BOTANICAL DESCRIPTION

The cultivar Balolesal has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype. The aforementioned photograph and following observations and measurements describe plants grown in West Chicago, Ill. under commercial practice in a polycarbonate-covered greenhouse with day temperatures ranging from 20 to 24°C., night temperatures ranging from 10 to 17° C. and light levels ranging from 4,000 to 6,000 footcandles. Rooted young plants were planted in containers and had been growing for about five to six weeks when the photographs and the description were taken.

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens walleriana* cultivar Balolesal.

Parentage:

*Female parent*.—Proprietary *Impatiens walleriana* selection identified as 3039C-1, not patented.

*Male parent*.—Proprietary *Impatiens walleriana* selection identified as 3111-1-9, not patented.

Propagation:

*Type cutting*.—Terminal cuttings.

*Time to initiate roots*.—About 7 days at 18° C.

*Time to produce a rooted young plant*.—About 21 days at 18° C.

*Root description*.—Fibrous and fine.

*Rooting habit*.—Freely branching.

Plant description:

*General appearance*.—Compact, upright, outwardly spreading and mounded plant habit. Appropriate for 10-cm and larger containers.

*Growth and branching habit*.—Vigorous and freely-branching growth habit with about three basal branches and lateral branches potentially forming at every node; dense and bushy growth. Pinching, that is, removal of the terminal apices, is not required, but will enhance branching.

*Plant height*.—About 11.5 cm.

*Plant diameter or spread*.—About 25 cm.

*Lateral branches*.—Length: About 13.4 cm. Diameter: About 8 mm. Internode length: About 3.2 cm. Texture: Smooth, glabrous. Color: 146C; random dark red purple, 59A, spots and streaks at nodes.

*Foliage description*.—Arrangement: Opposite, simple. Length: About 4.6 cm. Width: About 3 cm. Shape: Ovate. Apex: Acuminate. Base: Attenuate. Margin: Serrate to crenate with ciliation. Texture, upper and lower surfaces: Smooth, glabrous. Venation pattern: Pinnate, arcuate. Color: Young and fully expanded foliage, upper surface: 139A. Young and fully expanded foliage, lower surface: 138B. Venation, upper and lower surfaces: 144B. Petiole: Length: About 8 mm. Diameter: About 2 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color: 146D.

Flower description:

*Flower type and habit*.—Numerous and consistently double flowers. Flower buds open similar to a rose in fullness; flowers rounded in shape. Flowers arise from leaf axils. Freely flowering, usually about 16 flowers and flower buds per lateral branch. Flowers positioned above the foliage and typically face upright or outward. Flowers last about five to seven days under greenhouse conditions. Flowers not persistent. Flowers not fragrant.

*Flowering season*.—Year-round under greenhouse conditions. In the garden, flowering is continuous from spring until fall.

*Flower size*.—Diameter: About 4.3 cm. Depth: About 2.3 cm.

*Flower buds (at stage of showing color)*.—Length: About 1.7 cm. Diameter: About 1.3 cm. Shape: Ovoid. Color: 48C.

*Petals*.—Quantity/arrangement: About 37 per flower; imbricate. Length, largest petals: About 2 cm. Width, largest petals: About 1.8 cm. Shape: Obovate. Apex: Emarginate. Base: Truncate to attenuate. Margin: Entire. Texture, upper and lower surfaces: Velvety, smooth. Color: When opening and fully opened, upper surface: 39A; becoming closer to 52A with subsequent development. When opening and fully opened, lower surface: 48C.

*Sepals*.—Quantity/arrangement: Five in a single whorl; one modified into an elongated spur. Calyx length: About 1 cm. Calyx diameter: About 1 cm. Shape: Elliptic. Apex: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: 144C. Spur length: About 2.9 cm. Spur color: 145A overlain with 59A.

*Peduncles*.—Length: About 2.1 cm. Diameter: About 1.25 mm. Angle: Acute. Strength: Strong. Color: 144B.

*Reproductive organs*.—None observed.

*Seed/fruit*.—None observed.

Disease-pest resistance: Plants of the new *Impatiens* have not been observed to be resistant to pathogens and pests common to *Impatiens*.

It is claimed:

1. A new and distinct cultivar of Double *Impatiens* plant named 'Balolesal', as illustrated and described.

\* \* \* \* \*

