



US00PP13801P29

(12) **United States Plant Patent**  
**Vandenberg**

(10) **Patent No.:** **US PP13,801 P2**  
(45) **Date of Patent:** **May 13, 2003**

(54) **CHRYSANTHEMUM PLANT NAMED**  
**‘SWEET YOSTACY’**

(75) Inventor: **Cornelis P. Vandenberg**, Salinas, CA  
(US)

(73) Assignee: **Yoder Brothers, Inc.**, Barberton, OH  
(US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/094,269**

(22) Filed: **Mar. 8, 2002**

(51) Int. Cl.<sup>7</sup> ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./286**  
(58) **Field of Search** ..... **Plt./286**

*Primary Examiner*—Bruce R. Campell  
*Assistant Examiner*—Anne Marie Grünberg  
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A distinct cultivar of Chrysanthemum plant named ‘Sweet  
Yostacy’, characterized by its upright and mounded plant  
habit; freely branching habit; uniform and freely flowering  
habit; daisy-type inflorescences; pale yellow-colored ray  
florets with pink-colored apices; and natural season flower-  
ing in early October in the Northern Hemisphere.

**1 Drawing Sheet**

**1**

**BOTANICAL CLASSIFICATION/CULTIVAR  
DESIGNATION**

*Chrysanthemum*×*morifolium* cultivar Sweet Yostacy.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar  
of Chrysanthemum plant, botanically known as  
*Chrysanthemum*×*morifolium*, commercially known as a  
garden-type Chrysanthemum and hereinafter referred to by  
the name ‘Sweet Yostacy’.

The new cultivar is a product of a mutation induction  
breeding program conducted by the Inventor in Salinas,  
Calif. and Fort Myers, Fla. The objective of the program is  
to create new garden-type Chrysanthemum cultivars having  
inflorescences with desirable inflorescence forms, attractive  
florete colors and good garden performance.

The new Chrysanthemum originated by exposing  
unrooted cuttings of the Chrysanthemum cultivar Stacy,  
disclosed in U.S. Plant Pat. No. 11,852, to X-ray radiation in  
March, 1997 in Fort Myers, Fla. Following the radiation  
treatment, the cuttings were rooted and terminal apices were  
removed (pinched) three times to promote lateral branch  
development. After lateral branches from the third pinch  
reached sufficient size, terminal cuttings were harvested,  
planted and flowered in a controlled environment in Salinas,  
Calif. The new Chrysanthemum was discovered and  
selected by the Inventor as a single flowering plant within  
this population in October, 1997. The selection of this plant  
was based on its desirable inflorescence form, attractive ray  
florete color and good garden performance.

Asexual reproduction of the new cultivar by terminal  
cuttings taken in a controlled environment in Salinas, Calif.  
since December, 1997, has shown that the unique features of  
this new Chrysanthemum are stable and reproduced true to  
type in successive generations.

**SUMMARY OF THE INVENTION**

The cultivar Sweet Yostacy has not been observed under  
all possible environmental conditions. The phenotype may  
vary somewhat with variations in environment such as

**2**

temperature, daylength and light intensity, without,  
however, any variance in genotype.

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of ‘Sweet  
Yostacy’. These characteristics in combination distinguish  
‘Sweet Yostacy’ as a new and distinct cultivar:

1. Upright and mounded plant habit.
2. Freely branching habit; dense and full plants.
3. Uniform and freely flowering habit.
4. Daisy-type inflorescences.
5. Pale yellow-colored ray florets with pink-colored api-  
ces.
6. Natural season flowering in early October in the  
Northern Hemisphere.

Plants of the new Chrysanthemum are most similar to  
plants of the cultivar Stacy. In side-by-side comparisons  
conducted in Fort Myers, Fla., plants of the new Chrysan-  
themum differed from plants of the cultivar Stacy in the  
following characteristics:

1. Plants of the new Chrysanthemum flowered about three  
to four days earlier than plants of the cultivar Stacy when  
flowered under artificial daylength conditions.
2. Ray florets of the new Chrysanthemum and the cultivar  
Stacy differed in color as ray florets of the cultivar Stacy  
were white with pink apices.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying photographs illustrate the overall  
appearance of the new Chrysanthemum. These photographs  
show the colors as true as it is reasonably possible to obtain  
in colored reproductions of this type. Colors in the photo-  
graphs may differ slightly from the color values cited in the  
detailed botanical description which accurately describe the  
colors of the new Chrysanthemum.

The photograph at the top of the sheet comprises a side  
perspective view of a typical flowering plant of ‘Sweet  
Yostacy’.

The photograph at the bottom of the sheet comprises a  
close-up view of typical inflorescences of the cultivar  
‘Sweet Yostacy’.



## DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. The following observations and measurements describe plants grown in an outdoor nursery in Pendleton, S.C., under natural season conditions and practices which approximate those generally used in commercial garden-type Chrysanthemum production. One rooted cutting was planted in a 16.5-cm container in late July, 2002. Plants were not pinched, that is, the terminal apex was not removed to enhance branching. During the production of the plants, day temperatures ranged from 29 to 32° C. and night temperatures ranged from 16 to 21° C. Measurements and numerical values represent averages for typical flowering plants.

Botanical classification: *Chrysanthemum*×*morifolium* cultivar Sweet Yostacy.

Commercial classification: Daisy-type garden Chrysanthemum.

Parentage: Induced mutation of the *Chrysanthemum*×*morifolium* cultivar Stacy, disclosed in U.S. Plant Pat. No. 11,852.

Propagation:

*Type*.—Terminal tip cuttings.

*Time to initiate roots*.—About four days at 21° C.

*Time to produce a rooted cutting*.—About ten days at 21° C.

*Root description*.—White, fine and fibrous.

*Rooting habit*.—Freely branching.

Plant description:

*Appearance*.—Perennial herbaceous daisy-type garden Chrysanthemum. Inverted triangle. Stems initially upright, then somewhat outwardly spreading giving a uniformly mounded appearance to the plant. Freely branching with lateral branches forming at every node.

*Plant height*.—About 23 cm.

*Plant diameter*.—About 39 cm.

*Lateral branches*.—Length: About 20 cm. Diameter: About 3.5 mm. Internode length: About 3.2 cm. Aspect: Upright and outwardly spreading. Texture: Pubescent. Color: 146A.

*Foliage description*.—Leaf arrangement: Alternate. Length: About 4.6 cm. Width: About 3.7 cm. Apex: Cuspidate to mucronate. Base: Attenuate with truncate tendencies. Margin: Palmately lobed, sinuses parallel to divergent. Texture: Both surfaces, pubescent; veins prominent on lower surface. Color: Young and mature foliage upper surface: 147A. Young and mature foliage lower surface: 147B. Venation, upper surface: 147A to 147B. Venation, lower surface: 147B. Petiole length: About 7.5 mm. Petiole diameter: About 3.5 mm. Petiole color, both surfaces: Close to 147B.

Inflorescence description:

*Appearance*.—Daisy-type inflorescence form with elongated oblong-shaped ray florets. Inflorescences borne on terminals above foliage, arising from leaf axils. Disk and ray florets arranged acropetally on a capitulum. About 6 inflorescences per lateral.

*Flowering response*.—Under natural season conditions, plants flower in early October in the Northern Hemisphere and continue to flower for at least three weeks depending on weather conditions.

*Inflorescence bud (before showing color)*.—Height: About 4 mm. Diameter: About 6 mm. Phyllary color: 143A.

*Inflorescence size*.—Diameter: About 4.3 cm. Depth (height): About 1.3 cm. Disc diameter: About 1.3 cm. Receptacle diameter: About 3.5 mm.

*Ray florets*.—Shape: Elongated oblong. Length: About 2.1 cm. Corolla tube length: About 6.5 mm. Width: About 5 mm. Apex: Mostly emarginate. Margin: Entire. Texture: Smooth, glabrous, satiny. Surface: Mostly flat. Orientation: Initially upright, then about 65° from vertical. Number of ray florets per inflorescence: About 57 in about 2 to 3 rows. Color: When opening, upper surface: 4A to 4B; towards apices, faintly overlain with 53A. When opening, lower surface: 4B to 4C; towards apices, faintly underlain with 53A. Opened inflorescence, upper surface: 8B to 8C; towards apices, overlain with 53A to 59A. Opened inflorescence, lower surface: 8C; towards apices, faintly underlain with 53A to 59A.

*Disc florets*.—Shape: Tubular, apex dentate. Length: About 4.5 mm. Width: Apex: About 1.5 mm. Base: About 1 mm. Number of disc florets per inflorescence: About 110. Color: Immature: 154A to 12A. Mature: Apex: 12A. Mid-section: 154C to 154D. Base: 155D.

*Peduncle*.—Aspect: Flexible, angled about 45° from the stem. Length: First peduncle: About 1.4 cm. Fourth peduncle: About 3.4 cm. Diameter: About 1 mm. Texture: Pubescent. Color: 146A.

*Reproductive organs*.—Androecium: Present on disc florets only. Anther color: 12A. Pollen: Moderate. Pollen color: 15A to 17A. Gynoecium: Present on both ray and disc florets.

*Seed*.—Seed production has not been observed.

Disease/pest resistance: Plants of the new Chrysanthemum have not been shown to be resistant to pathogens and pests common to Chrysanthemums.

Garden performance: Plants of the new Chrysanthemum have been observed to be tolerant to rain, wind and temperatures ranging from 0 to more than 40° C.

It is claimed:

1. A new and distinct cultivar of Chrysanthemum plant named 'Sweet Yostacy', as illustrated and described.

\* \* \* \* \*



