



US00PP13880P29

(12) **United States Plant Patent**
Smith

(10) **Patent No.:** **US PP13,880 P2**
(45) **Date of Patent:** **Jun. 17, 2003**

(54) **CHRYSANTHEMUM PLANT NAMED**
'YOMINDY'

(58) **Field of Search** Plt./297

(75) **Inventor:** **Mark A. Smith**, Fort Myers, FL (US)

Primary Examiner—Bruce R. Campell

Assistant Examiner—Annette Para

(73) **Assignee:** **Yoder Brothers, Inc.**, Barberton, OH
(US)

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(*) **Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(57) **ABSTRACT**

A distinct cultivar of Chrysanthemum plant named
'Yomindy', characterized by its upright and mounded plant
habit; freely branching habit; uniform and freely flowering
habit; daisy-type inflorescences; purple-colored ray florets
that do not fade with bright yellow-colored disc florets; and
natural season flowering in late September in the Northern
Hemisphere.

(21) **Appl. No.:** **10/094,268**

(22) **Filed:** **Mar. 8, 2002**

(51) **Int. Cl.⁷** **A01H 5/00**

(52) **U.S. Cl.** **Plt./297**

1 Drawing Sheet

1

2

**BOTANICAL CLASSIFICATION/CULTIVAR
DESIGNATION**

Chrysanthemum×*morifolium* cultivar Yomindy.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of Chrysanthemum plant, botanically known as
Chrysanthemum×*morifolium*, commercially known as a
garden-type Chrysanthemum and hereinafter referred to by
the name 'Yomindy'.

The new cultivar is a product of a planned breeding
program conducted by the Inventor in Salinas, Calif. and
Fort Myers, Fla. The objective of the breeding program is to
create new garden-type Chrysanthemum cultivars having
inflorescences with desirable inflorescence forms, attractive
florete colors and good garden performance.

The new Chrysanthemum originated from a cross made in
December, 1996, in Salinas, Calif., of an unnamed Chry-
santhemum proprietary seedling selection, not patented, as
the female, or seed, parent with the Chrysanthemum cultivar
Empire Crown Jewel, disclosed in U.S. Plant Pat. No. 9,738,
as the male, or pollen, parent. The new Chrysanthemum was
discovered and selected by the Inventor as a single flowering
plant within the progeny of the stated cross grown in a
controlled environment in Fort Myers, Fla. in October, 1998.
The selection of this plant was based on its desirable
inflorescence form, attractive ray floret color and good
garden performance.

Asexual reproduction of the new cultivar by terminal
cuttings taken in a controlled environment in Fort Myers,
Fla. since December, 1998, has shown that the unique
features of this new Chrysanthemum are stable and repro-
duced true to type in successive generations.

SUMMARY OF THE INVENTION

The cultivar Yomindy has not been observed under all
possible environmental conditions. The phenotype may vary
somewhat with variations in environment such as
temperature, daylength and light intensity, without,
however, any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of
'Yomindy'. These characteristics in combination distinguish
'Yomindy' as a new and distinct cultivar:

1. Upright and mounded plant habit.
2. Freely branching habit; dense and full plants.
3. Uniform and freely flowering habit.
4. Daisy-type inflorescences.
5. Purple-colored ray florets that do not fade with bright
yellow-colored disc florets.
6. Natural season flowering in late September in the
Northern Hemisphere.

In side-by-side comparisons conducted in Fort Myers,
Fla., plants of the new Chrysanthemum differed from plants
of the female parent, the unnamed selection, in the following
characteristics:

1. Plants of the new Chrysanthemum flowered earlier than
plants of the female parent under natural season conditions.
2. Ray florets of the new Chrysanthemum and the female
parent differed in ray floret coloration.

Plants of the new Chrysanthemum are most similar to
plants of the male parent, the cultivar Empire Crown Jewel.
In side-by-side comparisons conducted in Fort Myers, Fla.,
plants of the new Chrysanthemum differed from plants of the
cultivar Empire Crown Jewel in the following characteris-
tics:

1. Plants of the new Chrysanthemum were fuller and more
dense than plants of the cultivar Empire Crown Jewel.
2. Plants of the new Chrysanthemum flowered more
uniformly than plants of the cultivar Empire Crown Jewel.
3. Ray florets of the new Chrysanthemum were darker in
color than ray florets of the cultivar Empire Crown Jewel.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall
appearance of the new Chrysanthemum. These photographs
show the colors as true as it is reasonably possible to obtain
in colored reproductions of this type. Colors in the photo-
graphs may differ slightly from the color values cited in the

detailed botanical description which accurately describe the colors of the new Chrysanthemum.

The photograph at the top of the sheet comprises a side perspective view of a typical flowering plant of 'Yomindy'.

The photograph at the bottom of the sheet comprises a close-up view of typical inflorescences of the cultivar 'Yomindy'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. The following observations and measurements describe plants grown in an outdoor nursery in Pendleton, S.C., under natural season conditions and practices which approximate those generally used in commercial garden-type Chrysanthemum production. One rooted cutting was planted in a 16.5-cm container in late July, 2002. Plants were not pinched, that is, the terminal apex was not removed to enhance branching. During the production of the plants, day temperatures ranged from 29 to 32° C. and night temperatures ranged from 16 to 21° C. Measurements and numerical values represent averages for typical flowering plants.

Botanical classification: *Chrysanthemum*×*morifolium* cultivar Yomindy.

Commercial classification: Daisy-type garden Chrysanthemum.

Parentage:

Female, or seed, parent.—Unnamed *Chrysanthemum*×*morifolium* proprietary seedling selection, not patented.

Male, or pollen, parent.—*Chrysanthemum*×*morifolium* cultivar Empire Crown Jewel, disclosed in U.S. Plant Pat. No. 9,738.

Propagation:

Type.—Terminal tip cuttings.

Time to initiate roots.—About four days at 21° C.

Time to produce a rooted cutting.—About ten days at 21° C.

Root description.—White, fine and fibrous.

Rooting habit.—Freely branching.

Plant description:

Appearance.—Perennial herbaceous daisy-type garden Chrysanthemum. Inverted triangle. Stems initially upright, then somewhat outwardly spreading giving a uniformly mounded appearance to the plant. Freely branching with lateral branches forming at every node.

Plant height.—About 31 cm.

Plant diameter.—About 46 cm.

Lateral branches.—Length: About 27 cm. Diameter: About 5 mm. Internode length: About 2.4 cm. Aspect: Upright and outwardly spreading. Texture: Pubescent. Color: Close to 146A.

Foliage description.—Leaf arrangement: Alternate. Length: About 4.3 cm. Width: About 3.4 cm. Apex: Cuspidate to mucronate. Base: Attenuate with truncate tendencies. Margin: Palmately lobed, sinuses mostly parallel. Texture: Both surfaces, pubescent; veins prominent on lower surface. Color: Young foliage upper surface: Darker than 147A. Young

foliage lower surface: Slightly darker than 147B. Mature foliage upper surface: Slightly darker than 147A. Mature foliage lower surface: Closest to 147B. Venation, upper surface: 147A to 147B. Venation, lower surface: 147B to 147C. Petiole length: About 2.1 cm. Petiole diameter: About 2.5 mm. Petiole color: Upper surface: 147A to 147B. Lower surface: Closest to 147C.

Inflorescence description:

Appearance.—Daisy-type inflorescence form with elongated oblong-shaped ray florets. Inflorescences borne on terminals above foliage, arising from leaf axils. Disk and ray florets arranged acropetally on a capitulum. About 5 inflorescences per lateral.

Flowering response.—Under natural season conditions, plants flower in early October in the Northern Hemisphere and continue to flower for at least three weeks depending on weather conditions.

Inflorescence bud (before showing color).—Height: About 6 mm. Diameter: About 7.5 mm. Phyllary color: Darker than 143A.

Inflorescence size.—Diameter: About 5.4 cm. Depth (height): About 2.1 cm. Disc diameter: About 1.3 cm. Receptacle diameter: About 5.5 mm.

Ray florets.—Shape: Elongated oblong. Length: About 2.75 cm. Corolla tube length: About 4.5 mm. Width: About 6 mm. Apex: Acute, emarginate or dentate. Margin: Entire. Texture: Smooth, glabrous, satiny. Surface: Initially concave, then flat. Orientation: Initially upright, then about 60° from vertical. Number of ray florets per inflorescence: About 51 in about two rows. Color: When opening, upper surface: Close to 155D, heavily overlain with 70A to 71A; towards base, close to 155D. When opening, lower surface: Close to 155D, underlain with 77A. Opened inflorescence, upper surface: Close to 155D, overlain with 70A to 71A; towards base, close to 155D; with subsequent development, less purple coloration. Opened inflorescence, lower surface: Close to 155D, underlain with 77A.

Disc florets.—Shape: Tubular, apex dentate. Length: About 6 mm. Width: Apex: About 2 mm. Base: About 1 mm. Number of disc florets per inflorescence: About 110. Color: Immature: 154A to 9A. Mature: Apex: 9A. Mid-section: 154D. Base: 155D.

Peduncle.—Aspect: Flexible, angled about 45 to 50° from the stem. Length: First peduncle: About 5.7 cm. Fourth peduncle: About 7.3 cm. Diameter: About 3 mm. Texture: Pubescent. Color: Close to 146A.

Reproductive organs.—Androecium: Present on disc florets only. Anther color: 9A. Pollen: Moderate. Pollen color: 15A. Gynoecium: Present on both ray and disc florets.

Seed.—Seed production has not been observed.

Disease/pest resistance: Plants of the new Chrysanthemum have not been shown to be resistant to pathogens and pests common to Chrysanthemums.

Garden performance: Plants of the new Chrysanthemum have been observed to be tolerant to rain, wind and temperatures ranging from 0 to more than 40° C.

It is claimed:

1. A new and distinct cultivar of Chrysanthemum plant named 'Yomindy', as illustrated and described.

* * * * *

