



US00PP13906P29

(12) **United States Plant Patent**  
**Dümmen**

(10) **Patent No.: US PP13,906 P2**  
(45) **Date of Patent: Jun. 24, 2003**

(54) **GERANIUM PLANT NAMED ‘DUEVIDARED’**

(75) Inventor: **Marga Dümmen**, Rheinberg (DE)

(73) Assignee: **Dümmen Jungpflanzen GbR**,  
Rheinberg (DE)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/228,461**

(22) Filed: **Aug. 27, 2002**

(51) **Int. Cl.<sup>7</sup>** ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./330**

(58) **Field of Search** ..... **Plt./330**

*Primary Examiner*—Kent Bell

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of Zonal Geranium plant named ‘Duevidared’, characterized by its upright, outwardly spreading and rounded plant habit; freely branching habit; freely flowering habit; and bright red-colored flowers.

**1 Drawing Sheet**

**1**

**BOTANICAL CLASSIFICATION/CULTIVAR  
DENOMINATION**

*Pelargonium×hortorum* cultivar ‘Duevidared’.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of Zonal Geranium plant, botanically known as *Pelargonium×hortorum*, and hereinafter referred to by the name ‘Duevidared’.

The new Zonal Geranium is a product of a planned breeding program conducted by the Inventor in Rheinberg, Germany. The objective of the breeding program was to develop new Zonal Geraniums with attractive flower and foliage colors.

The new Zonal Geranium originated from a cross-pollination made by the Inventor in September, 1998, of a proprietary selection of *Pelargonium×hortorum* identified as code number F-21-019, not patented, as the female, or seed, parent with a proprietary selection of *Pelargonium×hortorum* identified as code number F-05-018, not patented, as the male, or pollen, parent. The cultivar Duevidared was discovered and selected by the Inventor as a flowering plant within the progeny from this cross in a controlled environment in Rheinberg, Germany in September, 2000.

Asexual reproduction of the new cultivar by terminal vegetative cuttings taken at Rheinberg, Germany since September, 2000, has shown that the unique features of this new Zonal Geranium are stable and reproduced true to type in successive generations of asexual reproduction.

**SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Duevidared’. These characteristics in combination distinguish ‘Duevidared’ as a new cultivar and distinguish it from other known Zonal Geranium cultivars:

1. Upright, outwardly spreading and rounded plant habit.
2. Freely branching habit.
3. Freely flowering habit.
4. Bright red-colored flowers.

Compared to plants of the female parent, the selection F-21-019, plants of the new Zonal Geranium are more vigorous and have a brighter red flower color. Compared to

**2**

plants of the male parent, the selection F-05-018, plants of the new Zonal Geranium differ in flower color.

The new Zonal Geranium can be compared to plants of the cultivar ‘HWD Ventura’, disclosed in U.S. Plant Pat. No. 9,747. In side-by-side comparisons conducted in Rheinberg, Germany, plants of the new Zonal Geranium differed from plants of the cultivar HWD Ventura in the following characteristics:

1. Plants of the new Zonal Geranium were larger than plants of the cultivar HWD Ventura.
2. Plants of the new Zonal Geranium had larger flower umbels and more flowers per umbel than plants of the cultivar HWD Ventura.
3. Plants of the new Zonal Geranium had semi-double flowers whereas plants of the cultivar HWD Ventura had double flowers.
4. Plants of the new Zonal Geranium had more rounded flowers than plants of the cultivar HWD Ventura.
5. Flower color of plants of the new Zonal Geranium was brighter red than flower color of plants of the cultivar HWD Ventura.
6. Plants of the new Zonal Geranium had shorter peduncles than plants of the cultivar HWD Ventura.

**BRIEF DESCRIPTION OF THE PHOTOGRAPH**

The accompanying colored photograph illustrates the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Flower and foliage colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Zonal Geranium. The photograph comprises a side perspective view of a typical flowering plant of ‘Duevidared’ grown in a container.

**DETAILED BOTANICAL DESCRIPTION**

The cultivar Duevidared has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment, such as temperature and light intensity, without, however, any variance in genotype.

The aforementioned photograph, following observations and measurements describe plants grown in Rheinberg, Germany during the summer under commercial practice in



a glass-covered greenhouse with day and night temperatures about 18° C. and light levels about 4,500 foot-candles. Plants were grown in 10.5-cm containers. Plants were about eight weeks from unrooted cuttings when the photograph and the detailed botanical description were taken.

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Pelargonium×hortorum* cultivar Duevidared, (Known as Z98-2476-04 during testing).

Parentage:

*Female parent*.—Proprietary selection of *Pelargonium×hortorum* identified as F-21-019, not patented.

*Male parent*.—Proprietary selection of *Pelargonium×hortorum* identified as F-05-018, not patented.

Propagation:

*Type cutting*.—Terminal vegetative cuttings.

*Time to initiate roots*.—Summer: About 5 days at 20° C. Winter: About 7 days at 20° C.

*Time to develop roots*.—Summer: About 21 days at 20° C. Winter: About 28 days at 20° C.

*Root description*.—Fine, fibrous; white in color.

*Rooting habit*.—Freely branching.

Plant description:

*General appearance*.—Upright, outwardly spreading and rounded plant habit; densely foliated.

*Growth and branching habit*.—Moderately vigorous. Basal branching, about three lateral branches per plant.

*Plant height, from soil level to top of umbels*.—About 24 cm.

*Plant width*.—About 23 cm.

*Lateral branches*.—Length: About 6 cm. Internode length: About 1 cm. Texture: Smooth. Color: 144A.

*Foliage description*.—Arrangement: Alternate, single. Quantity of leaves per lateral branch: About six. Length: About 7.1 cm. Width: About 8.7 cm. Shape: Reniform. Apex: Rounded. Base: Cordate. Margin: Crenate. Venation pattern: Palmate. Texture, upper and lower surfaces: Pubescent. Color: Young foliage, upper surface: 137B. Young foliage, lower surface: 138B. Fully expanded foliage, upper surface: 137A. Fully expanded foliage, lower surface: 138B. Venation, upper and lower surfaces: 144A. Zonation pattern: Very faint. Width: About 3 cm. Color: Slightly darker than 137A. Petiole: Length: About 7.4 cm. Diameter: About 2.6 mm. Color, upper and lower surfaces: 144A.

Flower description:

*Flower arrangement*.—Bright red-colored semi-double flowers arranged in rounded hemispherical umbels arising from apical leaf axils. Umbels displayed above and beyond the foliage on peduncles. Flowers rounded in form. Umbels persistent, flowers not persistent. Flowers not fragrant.

*Quantity of flowers*.—Freely flowering; at full flower, plants have about five open and developing umbels with about 35 flowers per umbel.

*Flowering season*.—Flowering continuous spring until the autumn.

*Flower longevity*.—Flowers last about five to seven days on the plant.

*Umbel size*.—Height: About 7.5 cm. Diameter: About 14 cm.

*Flower size*.—Diameter: About 4.4 cm. Depth (height): About 1.7 cm.

*Flower buds*.—Length: About 1.4 cm. Diameter: About 9 mm. Shape: Ovoid. Color: 144A.

*Petals*.—Quantity per flower: About five. Length: About 2.7 cm. Width: About 2 cm. Shape: Obovate. Apex: Rounded. Base: Attenuate. Margin: Entire; sinuate. Texture, upper and lower surfaces: Smooth. Color: When opening and fully opened, upper surface: 42A to 45B. When opening and fully opened, lower surface: 43C.

*Petaloids*.—Quantity per flower: About three or four. Length: About 2.7 cm. Width: About 2 cm. Shape: Obovate. Apex: Rounded. Base: Attenuate. Margin: Entire; sinuate. Texture, upper and lower surfaces: Smooth. Color: When opening and fully opened, upper surface: 42A to 45B. When opening and fully opened, lower surface: 43C.

*Sepals*.—Quantity per flower: Five, arranged in a single whorl. Length: About 1.1 cm. Width: About 2.3 mm. Shape: Elongated, tapering. Apex: Apiculate. Margin: Entire. Texture, upper and lower surfaces: Smooth; glabrous. Color, upper and lower surfaces: 144A.

*Peduncle (umbel stem)*.—Length: About 10.9 cm. Diameter: Less than 1 cm. Angle: Mostly erect. Strength: Moderately strong. Texture: Smooth; glabrous. Color: 144A.

*Pedicel (individual flower stem)*.—Length: About 3.8 cm. Diameter: About 1 mm. Angle: Mostly erect. Strength: Moderately strong. Texture: Pubescent. Color: 59B.

*Reproductive organs*.—Androecium: Anther quantity per flower: About ten. Anther length: About 2 mm. Anther shape: Ovate. Anther color: 47B. Pollen amount: Moderate. Pollen color: 25A. Gynoecium: Pistil quantity per flower: One. Pistil length: About 1 cm. Stigma shape: Five-parted, star-shaped. Stigma color: 46A. Style length: About 5 mm. Style color: 59B. Ovary color: 144A.

*Seed/fruit*.—Development of seeds and fruit have not been observed.

Disease/pest resistance: Plants of the new Zonal Geranium have not been observed to be resistant to pathogens and pests common to Zonal Geraniums.

Temperature tolerance: Plants of the new Zonal Geranium have been observed to be tolerant to temperatures ranging from 5 to 40° C.

It is claimed:

1. A new and distinct cultivar of Zonal Geranium plant named 'Duevidared', as herein illustrated and described.

\* \* \* \* \*



