



US00PP14028P29

(12) **United States Plant Patent**
Takeshita et al.

(10) **Patent No.: US PP14,028 P2**
(45) **Date of Patent: Jul. 29, 2003**

(54) **CALIBRACHOA PLANT NAMED
‘ILLUMINATION ROSE 2’**

(75) Inventors: **Daigaku Takeshita**, Utsunomiya (JP);
Tsutomu Tokita, Omiya (JP); **Saori**
Watanabe, Ujiie-machi (JP); **Noboru**
Onishi, Thousand Oaks, CA (US);
Kunihiro Hayashida, Utsunomiya (JP)

(73) Assignees: **Kirin Brewery Co. Ltd.**, Tokyo (JP);
Tokita Seed Co. Ltd., Saitama-Ken
(JP)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(22) Filed: **Mar. 8, 2002**

(51) **Int. Cl.⁷** **A01H 5/00**
(52) **U.S. Cl.** **Plt./263**
(58) **Field of Search** **Plt./263**

Primary Examiner—Bruce R. Campell
Assistant Examiner—June Hwu
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A distinct cultivar of Calibrachoa plant named ‘Illumination
Rose 2’, characterized by its cascading and prostrate plant
habit; freely branching growth habit; early flowering habit;
purple-colored flowers with yellow-colored throat; and good
weather tolerance.

(21) Appl. No.: **10/094,319** **2 Drawing Sheets**

1

**BOTANICAL CLASSIFICATION/CULTIVAR
DESIGNATION**

Calibrachoa sp. cultivar Illumination Rose 2.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of Calibrachoa plant, botanically known as Calibrachoa
sp., and hereinafter referred to by the name ‘Illumination
Rose 2’.

The new Calibrachoa is a product of a planned breeding
program conducted by the Inventors in Tochigi, Japan. The
objective of the breeding program is to create new early
flowering Calibrachoa cultivars that have stronger growth
and attractive flower coloration.

The new Calibrachoa originated from a cross-pollination
made by the Inventors in August, 1997, in Tochigi, Japan, of
an unnamed proprietary Calibrachoa selection, not patented,
as the female, or seed, parent with an unnamed proprietary
Calibrachoa selection, not patented, as the male, or pollen,
parent. The new Calibrachoa was discovered and selected by
the Inventors as a single flowering plant within the progeny
of the stated cross-pollination grown in a controlled envi-
ronment in Tochigi, Japan, in October, 1998. The selection
of this plant was based on its strong plant growth habit and
attractive flower coloration.

Asexual reproduction of the new cultivar by terminal
cuttings taken in a controlled environment in Saitama-ken,
Japan, since January, 1999, has shown that the unique
features of this new Calibrachoa are stable and reproduced
true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the cultivar Illumination Rose 2 have not been
observed under all possible environmental conditions. The
phenotype may vary somewhat with variations in environ-
ment such as temperature and light intensity without,
however, any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Illumi-
nation Rose 2’. These characteristics in combination distin-

2

guish ‘Illumination Rose 2’ as a new and distinct cultivar of
Calibrachoa:

1. Cascading and prostrate plant habit.
2. Freely branching growth habit.
3. Early flowering habit.
4. Purple-colored flowers with yellow-colored throat.
5. Good weather tolerance; tolerant to wind, rain, and low
and high temperatures.

Plants of the new Calibrachoa can be compared to plants
of the female parent, an unnamed proprietary Calibrachoa
selection. In side-by-side comparisons conducted by the
Inventors in Tochigi, Japan, plants of the new Calibrachoa
differed from plants of the female parent primarily in growth
habit as plants of the new Calibrachoa were more prostrate
and not as upright as plants of the female parent. In addition
plants of the new Calibrachoa and the female parent differed
in petal color as plants of the female parent had dark
pink-colored petals.

Plants of the new Calibrachoa can be compared to plants
of the male parent, an unnamed proprietary Calibrachoa
selection. In side-by-side comparisons conducted by the
Inventors in Tochigi, Japan, plants of the new Calibrachoa
differed from plants of the male parent primarily in flower-
ing time as plants of the new Calibrachoa flowered earlier
than plants of the male parent.

Plants of the new cultivar can also be compared to plants
of the cultivar Sunbelkupi, disclosed in U.S. Plant Pat. No.
10,287. In side-by-side comparisons conducted by the
Inventors in Tochigi, Japan, plants of the new Calibrachoa
differed from plants of the cultivar Sunbelkupi in the fol-
lowing characteristics:

1. Plants of the new Calibrachoa had lighter green-colored
leaves than plants of the cultivar Sunbelkupi.
2. Plants of the new Calibrachoa had larger flowers than
plants of the cultivar Sunbelkupi.
3. Plants of the new Calibrachoa and the cultivar Sun-
belkupi differed in corolla coloration.
4. Plants of the new Calibrachoa had shorter peduncles
than plants of the cultivar Sunbelkupi.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Calibrachoa.

The photograph on the first sheet comprises a side perspective view of a typical flowering plant of 'Illumination Rose 2'.

The photograph on the second sheet comprises a close-up view of a typical flower of 'Illumination Rose 2'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used. Twenty-centimeter containers with three plants per container were used for the aforementioned photographs and the following description. Plants were grown under conditions which closely approximate commercial production conditions during the winter in Santa Paula, Calif. in a polyethylene-covered greenhouse. Plants used for the photographs and the description were about 14 weeks from planting rooted young plants. During the production period, day temperatures averaged 28° C., night temperatures averaged 21° C., and light levels were about 5,000 footcandles.

Botanical classification: Calibrachoa sp. cultivar Illumination Rose 2.

Parentage:

Female, or seed, parent.—Unnamed proprietary selection of Calibrachoa sp., not patented.

Male, or pollen, parent.—Unnamed proprietary selection of Calibrachoa sp., not patented.

Propagation:

Type cutting.—Terminal vegetative cuttings.

Time to initiate roots.—Summer: About 3 days at 25° C.

Winter: About 7 days at 20° C.

Time to produce a rooted young plant.—Summer: About 21 days at 25° C. Winter: About 23 days at 22° C.

Root description.—Fine, fibrous, and white in color.

Rooting habit.—Freely branching; dense.

Plant description:

Form.—Annual flowering plant; cascading and prostrate plant habit; low mounded; short internodes, dense and bushy appearance.

Plant height (from soil level to top of plant plane).—About 20 cm.

Plant diameter (area of spread).—About 84 cm.

Growth rate.—Relatively rapid.

Branching habit.—Freely basal branching, about 6 to 8 basal branches per plant; lateral branches develop at potentially every node; pinching is typically not required.

Lateral branch description.—Length: About 44 cm. Diameter: About 2.75 mm. Internode length: About 1 cm. Orientation: Initially upright, then horizontal. Texture: Pubescent; short, fine hairs. Strength: Strong, but flexible. Color: 146C.

Foliage description.—Leaves simple, generally symmetrical. Arrangement: Alternate before flowering, then opposite. Length: About 2.8 cm. Width: About 8 mm. Shape: Narrowly elliptic. Apex: Rounded. Base: Acute. Margin: Entire. Texture, upper surface: Coarse; slightly glandular; slightly pubescent. Texture, lower surface: Coarse; slightly glandular;

glabrous. Venation pattern: Pinnate; arcuate. Color: Young foliage, upper and lower surfaces: 147A. Fully expanded foliage, upper surface: 147A. Fully expanded foliage, lower surface: 147B. Venation, upper surface: 147A. Venation, lower surface: 147B. Petiole length: About 5 mm. Petiole diameter: About 1 mm. Petiole color: 144A.

Flower description:

Flower type and habit.—Flowers face upright or outward; solitary and axillary; salverform. Freely flowering habit, about 22 to 26 flowers and flower buds per lateral stem. Flowers persistent. Flowers not fragrant.

Natural flowering season.—Spring until frost in the autumn; flowering continuous during this period.

Time to flower.—Early flowering; plants begin flowering about 2 weeks after planting.

Flower longevity on the plant.—About 2 weeks.

Flower size.—Diameter: About 3.4 cm. Tube length: About 2.5 cm.

Flower buds (showing color).—Length: About 2 cm. Diameter: At apex, about 5 mm; at base, about 3 mm.

Shape: Roughly oblong. Color: 79D.

Corolla.—Arrangement/appearance: Single whorl of five petals, fused into flared trumpet. Petal length from throat: About 1.5 cm. Petal width: About 1.6 cm. Petal shape: Spatulate to flabellate. Petal apex: Rounded with shallow emargination. Petal margin: Entire. Petal texture: Smooth, glabrous; velvety. Color: Petal, when opening, upper surface: Brighter than 71B; venation, close to 71B. Petal, when opening, lower surface: 77B to 77C; venation, close to 77B to 77C. Petal, opened flower, upper surface: 74A; towards throat, 77B; venation, close to 74A; color does not fade with subsequent development. Petal, opened flower, lower surface: 77B to 77C; venation, close to 77B to 77C. Flower throat (inside): 3B to 3D; venation, close to 3B to 3D. Flower tube (outside): 145D; venation, close to 145D.

Sepals.—Arrangement/appearance: Single whorl of five sepals fused at base; star-shaped. Length: About 1.2 cm. Width: About 2.5 mm. Shape: Narrowly elliptic. Apex: Broadly acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: Young sepals, upper surface: 147A. Young sepals, lower surface: 147B. Fully expanded sepals, upper and lower surfaces: 147A.

Peduncles.—Length: About 2.2 cm. Width: About 1 mm. Strength: Moderately strong. Angle: About 45° from stem. Texture: Pubescent. Color: 144B.

Reproductive organs.—Stamens: Quantity: Five per flower. Anther shape: Oval. Anther length: Less than 1 mm. Anther color: 12C. Pollen amount: Scarce. Pollen color: 12C. Pistils: Quantity: One per flower. Pistil length: About 1.2 cm. Stigma shape: Anvil-shaped; curved. Stigma color: 144B. Style length: About 8 mm. Style color: 144C. Ovary color: 150C.

Seed/fruit.—Seed and fruit production has not been observed.

Disease/pest resistance: Plants of the new Calibrachoa have not been noted to be resistant to pathogens and pests common to Calibrachoa.

Weather/temperature tolerance: Plants of the new Calibrachoa are tolerant to rain and wind and have been observed to tolerate temperatures from 3 to 35° C.

It is claimed:

1. A new and distinct cultivar of Calibrachoa plant named 'Illumination Rose 2', as illustrated and described.

* * * * *



