



US00PP14032P29

(12) **United States Plant Patent**
Kientzler(10) **Patent No.:** **US PP14,032 P2**
(45) **Date of Patent:** **Jul. 29, 2003**(54) **NEW GUINEA IMPATIENS PLANT NAMED
'KIOTOA'**(75) Inventor: **Ludwig Kientzler**, Gensingen (DE)(73) Assignee: **Innovaplant GmbH + Co. KG**,
Gensingen (DE)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/159,064**(22) Filed: **May 31, 2002**(51) Int. Cl.⁷ **A01H 5/00**(52) U.S. Cl. **Plt./318**
(58) Field of Search **Plt./318**

Primary Examiner—Bruce R. Campell

Assistant Examiner—Susan B. McCormick

(74) Attorney, Agent, or Firm—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of New Guinea Impatiens plant named 'Kiotoa', characterized by its rounded, upright and compact plant habit; freely branching growth habit; bushy appearance; variegated foliage; freely flowering habit with flowers held above and beyond the foliage; and red-colored flowers.

1 Drawing Sheet**1****BOTANICAL CLASSIFICATION/CULTIVAR
DESIGNATION***Impatiens hawkeri* cultivar Kiotoa.**BACKGROUND OF THE INVENTION**

The present Invention relates to a new and distinct cultivar of New Guinea Impatiens plant, botanically known as *Impatiens hawkeri*, and hereinafter referred to by the cultivar name Kiotoa.

The new Impatiens is a product of a planned breeding program Inventor in Gensingen, Germany. The objective of the breeding program is to develop new Impatiens cultivars with interesting and unique flower and foliage colors.

The new Impatiens originated from a cross-pollination made by the Inventor in September, 1998 of the *Impatiens hawkeri* cultivar Eurema, disclosed in U.S. Plant Pat. No 6,681, as the female, or seed parent, with a proprietary *Impatiens hawkeri* selection identified as 97-286, not patented, as the male, or pollen parent. The cultivar Kiotoa was discovered and selected by the Inventor as a flowering plant within the progeny of the stated cross-pollination in a controlled environment in Gensingen, Germany in March, 1999.

Asexual reproduction of the new cultivar by terminal cuttings taken at Gensingen, Germany, since April, 1999, has shown that the unique features of this new Impatiens are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Kiotoa'. These characteristics in combination distinguish 'Kiotoa' as a new and distinct Impatiens cultivar:

1. Rounded, upright and compact plant habit.
2. Freely branching growth habit; bushy appearance.
3. Variegated foliage.
4. Freely flowering habit with flowers held above and beyond the foliage.
5. Red-colored flowers.

2

Plants of the new Impatiens differ from plants of the female parent, the cultivar Eurema, in the following characteristics:

1. Leaves of plants of the new Impatiens are more strongly variegated than leaves of plants of the cultivar Eurema.

2. Flower color of plants of the new Impatiens is red whereas flower color of plants of the cultivar Eurema is orange.

10 3. Plants of the new Impatiens differ from plants of the male parent, the selection 97-286, primarily in growth habit as plants of the new Impatiens are more freely branching than plants of the selection 97-286.

15 4. Plants of the new Impatiens can be compared to plants of the cultivar Moala, disclosed in U.S. Plant Pat. No. 10,301. In side-by-side comparisons conducted in Gensingen, Germany, plants of the new Impatiens differed from plants of the cultivar Moala primarily in leaf coloration as plants of the cultivar Moala did not have variegated leaves.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which more accurately describe the actual colors of the new Impatiens. The photograph comprises a side perspective view of three typical flowering plants of 'Kiotoa' grown in a 22-cm container.

DETAILED BOTANICAL DESCRIPTION

35 The cultivar Kiotoa has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype. The aforementioned photograph and the following observations and measurements describe plants grown in Encinitas, Calif., during the late spring, under commercial practice in a polyethylene-covered greenhouse with day temperatures about 24° C., night temperatures about 18° C., and light levels typically about 4,000 foot-candles. Plants

40

used in the photograph and following description were about 14 weeks old and grown in 22-cm containers with three plants per container.

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens hawkeri* cultivar Kiotoa.
Parentage:

Female, or seed, parent.—*Impatiens hawkeri* cultivar Eurema, disclosed in U.S. Plant Pat. No. 6,681.

Male, or pollen, parent.—Proprietary *Impatiens hawkeri* selection identified as 97-286, not patented.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—Summer: About 14 days at 23° C. Winter: About 17 days at 20° C.

Time to produce a rooted cutting or liner.—Summer: About 21 days at 23° C. Winter: About 24 days at 20° C.

Root description.—Fine, fibrous, and freely branching.

Plant description:

General appearance.—Rounded and somewhat upright; compact. Appropriate for 10 to 25-cm containers; multiple plants are typically planted in larger containers.

Growth and branching habit.—Freely branching with about ten lateral branches developing at the base, dense and bushy growth. Pinching, that is, removal of the terminal apices, is typically not required. Moderately vigorous.

Plant height.—About 17 cm.

Plant diameter, single plant.—About 18 cm.

Lateral branches.—Length: About 10.5 cm. Diameter: About 6 mm. Internode length: About 3.5 cm. Texture: Sparsely pubescent. Color: 182A to 182B.

Foliage description.—Arrangement: Opposite or in whorls; simple. Length: About 9.25 cm. Width: About 4.3 cm. Shape: Elliptic. Apex: Acute. Base: Attenuate. Margin: Serrulate with ciliation. Texture, upper surface: Slightly pubescent. Texture, lower surface: Smooth, glabrous. Color: Young foliage, upper surface: Towards margin, 147A; towards center, 154A. Young foliage, lower surface: Towards margin, 147C; towards center, 145C. Fully expanded foliage, upper surface: Towards margin, 147A; towards center, 10B to 10C. Fully expanded foliage, lower surface: Towards margin, 147B to 147C; towards center, 10D. Venation, upper surface: 63C. Venation, lower surface: 58D. Petiole: Length: About 4.5 cm. Diameter: About 4 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: 50C.

Flower description:

Flower type and flowering habit.—Single axillary flowers. Freely flowering, usually about seven flowers and flower buds per lateral branch. Flowers typically face upward or outward and held above and beyond the foliage. Flowers somewhat cupped and rounded triangular in shape. Flowers last about eight days on the plant depending on temperature and weather conditions. Petals self-cleaning; gynoecium persistent. Flowers not fragrant.

Flowering season.—Year-round under greenhouse conditions. In the garden, flowering from spring until fall.

Flower height.—About 5.5 cm.

Flower width.—About 5 cm.

Flower depth.—About 2 cm.

Flower buds (at stage of showing color).—Rate of opening: From showing color to fully open flower, typically about 2 to 3 days depending on temperature. Length: About 1.5 cm. Diameter: About 8 mm. Shape: Ovoid. Color: 46C.

Petals.—Quantity: Single, five per flower. Length: Banner petal: About 2.2 cm. Lateral petals: About 2.5 cm. Base petals: About 3 cm. Width: Banner petal: About 3.5 cm. Lateral petals: About 2.5 cm. Base petals: About 3 cm. Shape: Cordate. Apex: Emarginate. Base: Acute. Margin: Entire. Texture: Smooth; velvety. Color: When opening, upper surface: 45A. When opening, lower surface: 44A. Fully opened, upper surface: 45A. Fully opened, lower surface: 44C.

Spur.—Length: About 4.6 cm. Texture: Smooth, glabrous. Color: 47A to 47B.

Peduncles.—Length: About 3.5 cm. Diameter: About 1 mm. Strength: Strong. Angle: About 45° from vertical. Texture: Smooth, glabrous. Color: 53C.

Reproductive organs.—Androecium: Stamen number: Five fused at anthers, filaments free. Anther shape: Obovate. Anther size: About 4 mm by 5 mm. Anther color: 10C. Amount of pollen: Moderate. Pollen color: 158B. Gynoecium: Pistil length: About 5 mm. Stigma color: 155A. Style color: 146A. Ovary color: 146A.

Seed/fruit.—Seed and fruit development has not been observed.

Disease/pest resistance: Plants of the new *Impatiens* have not been observed to be resistant to pathogens and pests common to *Impatiens*.

Temperature tolerance: Plants of the new *Impatiens* have been observed to tolerate temperatures from 16 to 30° C. It is claimed:

1. A new and distinct cultivar of New Guinea *Impatiens* plant named 'Kiotoa', as illustrated and described.

* * * * *

