



US00PP14404P39

(12) **United States Plant Patent**
Pieters

(10) **Patent No.: US PP14,404 P3**
(45) **Date of Patent: Dec. 23, 2003**

(54) **CHRYSANTHEMUM PLANT NAMED ‘GEDI TWO COP’**

(50) Latin Name: *Chrysanthemum morifolium*
Varietal Denomination: **Gedi Two Cop**

(75) Inventor: **Dirk Pieters**, Staden (BE)

(73) Assignee: **Pieters Plant Production, BVBA**,
Oostnieuwkerke (BE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/137,801**

(22) Filed: **May 1, 2002**

(65) **Prior Publication Data**
US 2003/0208817 P1 Nov. 6, 2003

(51) **Int. Cl.⁷ A01H 5/00**

(52) **U.S. Cl. Plt./287**

(58) **Field of Search Plt./292, 287, 291**

(56) **References Cited**
U.S. PATENT DOCUMENTS
PP11,300 P 3/2000 Boeder Plt./287

Primary Examiner—Bruce R. Campell
Assistant Examiner—June Hwu

(74) *Attorney, Agent, or Firm*—Knobbe, Martens, Olson &
Bear, LLP

(57) **ABSTRACT**
A new and distinct Chrysanthemum plant cultivar is
disclosed, characterized by a decorative-type inflorescence,
consistent flowering response to short days, blooming con-
sistently after 42 days of short day length, free branching
habit, purple ray florets, a very uniform round growth habit,
and a large quantity of blooms per flowering branch.

1 Drawing Sheet

1

Latin name of the genus and species: The present inven-
tion relates to a new and distinct cultivar of Chrysanthemum
plant, botanically known as *Chrysanthemum morifolium*.
Variety denomination: The new and distinct Chrysante-
mum plant is hereinafter referred to by the cultivar name
‘Gedi Two Cop.’

BACKGROUND OF THE INVENTION

The new cultivar is a product of hybridization of a female
parent Chrysanthemum plant variety ‘Gedi One Car’ (U.S.
Plant Pat. No. 13,794) and a male parent Chrysanthemum
plant variety ‘Gedi One Gal’ (U.S. Plant Pat. No. 13,807).
The new cultivar was discovered and selected by the inven-
tor in September 1996.

Asexual reproduction of the new cultivar by apical tip
cutting and meristem tissue culture was performed in
Oxnard, Calif. and has shown that the unique features of this
cultivar are stable and reproduced true to type on successive
generations.

SUMMARY OF THE INVENTION

The cultivar ‘Gedi Two Cop’ has not been observed under
all possible environmental conditions. The phenotype may
vary somewhat with variations in environment such as
temperature, daylength, and light intensity, without,
however, any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Gedi Two
Cop.’ These characteristics in combination distinguish ‘Gedi
Two Cop’ as a new and distinct Chrysanthemum cultivar:

1. Decorative-type inflorescence,
2. Consistent flowering response to short days, blooming
consistently after 42 days of short day length,
3. Free branching habit,
4. Purple ray florets,

2

5. Very uniform round growth habit, and

6. Large quantity of blooms per flowering branch.

Plants of the new cultivar are similar to plants of the
female parent variety, ‘Gedi One Car’ in most horticultural
characteristics, however plants of the new cultivar bloom
earlier, have fewer ray florets, and have a smaller diameter
inflorescence than plants of the female parent variety.

Plants of the new cultivar are similar to plants of the male
parent variety, ‘Gedi One Gal’ in most horticultural
characteristics, however plants of the new cultivar have
inflorescences of a different color, fewer ray florets, and a
smaller diameter inflorescence than plants of the male parent
variety.

In comparison to the commercially available variety ‘Cle-
argar VZP1’ (U.S. Plant Pat. No. 11,300), ‘Gedi Two Cop’
blooms earlier and has fewer ray florets per inflorescence,
ray florets without irregular lobes, and fewer disc florets.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The photograph shows a plant of ‘Gedi Two Cop’ grown
in a six-inch container. One cutting was used in the pot.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to
The Royal Horticultural Society Colour Chart except where
general terms of ordinary dictionary significance are used.
The following observations and measurements describe
plants grown in Oxnard, Calif. during the month of Decem-
ber and January. The age of the observed plants is 86 days
from a rooted cutting. The growing temperature ranged from
55° to 68° F. at night to 60° to 75° F. during the day.
Measurements and numerical values represent averages of
typical flowering types.-

Botanical classification: *Chrysanthemum morifolium* cultivar 'Gedi Two Cop'.

Commercial classification: Garden-type Chrysanthemum

Parentage: Hybridization of a female parent Chrysanthemum plant variety, 'Gedi One Car' (U.S. Plant Pat. No. 13,794), and a male parent Chrysanthemum plant variety 'Gedi One Gal' (U.S. Plant Pat. No. 13,807)

PROPAGATION

Time to rooting: 7 to 14 days at approximately 21° C.

Root description: Fine, fibrous.

PLANT

Growth habit: Mounding herbaceous perennial.

Height: Approximately 19.5 cm.

Spread: Approximately 23 cm.

Growth rate: Moderate.

Branching characteristics: Free Branching.

Length of lateral branches: Approximately 20 cm.

Number of leaves per lateral branch: Approximately 36.

FOLIAGE

Leaf:

Arrangement.—Alternate.

Average length.—Approximately 5.2 cm.

Average width.—Approximately 4.6 cm.

Shape of blade.—Ovate.

Apex.—Cuspidate.

Base.—Attenuate.

Attachment.—Stalked.

Margin.—Palmately lobed.

Texture of top surface.—Lightly pubescent.

Texture of bottom surface.—Lightly pubescent.

Color.—young foliage upper side: Near 139A. young foliage under side: Near 137B. mature foliage upper side: Near 139A. mature foliage under side: Near 137B. venation upper side: Near 138B. venation under side: Near 137D.

Venation type.—Palmately net.

Petiole:

Average length.—Approximately 2.6 cm.

Color.—Near 139B.

Diameter.—Approximately 0.2 cm.

BLOOM

Inflorescence:

Flowering habit.—Induced by darkness period greater than 13.5 hours, approximately 42 days of appropriate day length required to induce and develop blooms.

Inflorescence form.—Decorative.

Natural flowering season.—Approximately the middle two weeks of September.

Number of inflorescences per lateral branch.—Approximately 15.

Inflorescence diameter.—Approximately 4.6 cm.

Inflorescence depth.—Approximately 1.4 cm.

Inflorescence longevity on plant.—Approximately 27 days.

Persistence.—Persistent.

Ray florets:

Appearance.—Matte.

Texture.—Smooth.

Average number per flower.—140.

Shape.—Oblong.

Aspect.—Flat.

Margin.—Entire.

Apex.—Retuse.

Length.—Approximately 2.6 cm.

Width.—Approximately 0.5 cm.

Color.—upper surface at first opening: Near 73B. upper surface at maturity: Near 73B. upper surface at fading: Near 73B. under surface at first opening: Near 73C. under surface at maturity: Near 73C. under surface at fading: Near 73C.

Disc florets:

Appearance.—Shiny.

Texture.—Smooth.

Average number per flower.—Approximately 3.

Shape.—Cylindric.

Apex.—Obtuse.

Average length.—Approximately 0.4 cm.

Average width.—Approximately 0.1 cm.

Color.—at first opening: Near 7C. at maturity: Near 13A. at fading: Near 15A.

Peduncle:

Length.—at terminal end (shortest): Approximately 1.6 cm. at lateral end (longest): Approximately 7 cm.

Angle to stem.—Acute.

Strength.—Moderate.

Color.—Near 137C.

Habit.—Upright.

Diameter.—Approximately 0.2 cm.

Surface texture.—Lightly pubescent.

Inflorescence bud:

Length.—Approximately 0.6 cm.

Diameter.—Approximately 0.7 cm.

Form.—Globular.

Color.—Near 137D.

Involucral bracts (phyllaries):

Appearance.—Matte.

Texture.—Lightly pubescent.

Number.—Approximately 24.

Shape.—Oblanceolate.

Margin.—Entire.

Apex.—Acute.

Length.—Approximately 0.8 cm.

Width.—Approximately 0.4 cm.

Color.—upper side: Near 141A. under side: Near 141A.

REPRODUCTIVE ORGANS

Ray florets:

Number of pistils per flower.—1.

Stigma shape.—2 branched.

Stigma color.—Near 5A.

Style color.—Near 154D.

Style length.—Approximately 0.3 cm.

Stamens.—Absent.

Disc florets (hermaphroditic):

Number of pistils per flower.—1.

Stigma shape.—Two-branched.

Stigma color.—Near 12B.

Style length.—Approximately 0.3 cm.

Style color.—Near 154C.

Number of stamens per flower.—5.

Anther shape.—Tubular.

Anther color.—Near 13A.

Pollen.—No pollen detected.

OTHER CHARACTERISTICS

Seed production: Commercially, this plant is not used or observed in a stage wherein seeds would be produced. Therefore, seed production has not been observed.

Disease resistance: Neither resistance nor susceptibility to diseases and pests has been observed in this cultivar.

Heat and cold resistance: Plants with flowers are hardy to low temperatures about -2° C. Non flowering plants are hardy in the approximate range of 3° C. to -6° C., depending upon duration of cold and amount of moisture

in the soil. With adequate water plants are hardy to a high temperature of 49° C.

What is claimed is:

1. A new and distinct cultivar of Chrysanthemum plant named 'Gedi Two Cop' as herein illustrated and described.

* * * * *

