



US00PP14618P29

(12) **United States Plant Patent**  
**Klemm**

(10) **Patent No.: US PP14,618 P2**  
(45) **Date of Patent: Mar. 23, 2004**

(54) **POINSETTIA PLANT NAMED ‘NPCW02044’**

(50) Latin Name: *Euphorbia pulcherrima*  
Varietal Denomination: **NPCW02044**

(75) Inventor: **Nils Klemm**, Stuttgart (DE)

(73) Assignee: **Klemm + Sohn GmbH + Co. KG**,  
Stuttgart (DE)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/452,068**

(22) Filed: **Jun. 2, 2003**

(51) **Int. Cl.<sup>7</sup>** ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./307**

(58) **Field of Search** ..... **Plt./307**

*Primary Examiner*—Kent Bell  
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of Poinsettia plant named  
‘NPCW02044’, characterized by its compact, uniform and  
upright plant habit; dark green-colored leaves; red-colored  
flower bracts that resist fading; early flowering habit; and  
good post-production longevity.

**1 Drawing Sheet**

**1**

Botanical classification/cultivar denomination: *Euphor-*  
*bia pulcherrima* Willd. cultivar NPCW02044.

**BACKGROUND OF THE INVENTION**

The present Invention relates to a new and distinct culti-  
var of Poinsettia plant, botanically known as *Euphorbia*  
*pulcherrima* Willd., and hereinafter referred to by the cul-  
tivar name ‘NPCW02044’.

The new Poinsettia is a product of a planned breeding  
program conducted by the Inventor in Stuttgart, Germany.  
The objective of the breeding program is to create new freely  
branching Poinsettia cultivars with upright and uniform  
plant habit and attractive flower bract coloration.

The new Poinsettia originated from a cross-pollination  
made by the Inventor in 1997 of a proprietary selection of  
*Euphorbia pulcherrima* Willd. identified as code number S  
219, not patented, as the female, or seed, parent, with a  
proprietary selection of *Euphorbia pulcherrima* Willd. iden-  
tified as code number R 82, not patented, as the male, or  
pollen, parent. The cultivar NPCW02044 was discovered  
and selected by the Inventor in 1999 as a flowering plant  
within the progeny of the stated cross-pollination in a  
controlled environment in Stuttgart, Germany. The selection  
of this plant was based on its attractive leaf and flower bract  
coloration and uniform plant habit.

Asexual reproduction of the new Poinsettia by vegetative  
terminal cuttings taken in a controlled environment in  
Stuttgart, Germany since 2000, has shown that the unique  
features of this new Poinsettia are stable and reproduced true  
to type in successive generations of asexual reproduction.

**BRIEF SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of  
‘NPCW02044’. These characteristics in combination distin-  
guish ‘NPCW02044’ as a new and distinct Poinsettia culti-  
var:

1. Compact, upright and uniform plant habit.
2. Dark green-colored leaves.
3. Red-colored flower bracts.

**2**

4. Early flowering; response time, about eight weeks.

5. Good post-production longevity.

Plants of the new Poinsettia differ from plants of the  
parent selections primarily in plant habit and flower bract  
coloration.

Plants of the new Poinsettia can be compared to plants of  
the cultivar Fiscor, disclosed in U.S. Plant Pat. No. 9,364. In  
side-by-side comparisons conducted in Stuttgart, Germany,  
plants of the new Poinsettia differed from plants of the  
cultivar Fiscor in the following characteristics:

1. Plants of the new Poinsettia were more compact than  
plants of the cultivar Fiscor.
2. Plants of the new Poinsettia were more upright and not  
as outwardly spreading as plants of the cultivar Fiscor.

**BRIEF DESCRIPTION OF THE PHOTOGRAPH**

The accompanying colored photograph illustrates the  
overall appearance of the new Poinsettia, showing the colors  
as true as it is reasonably possible to obtain in colored  
reproductions of this type. Colors in the photograph may  
differ slightly from the color values cited in the detailed  
botanical description which accurately describe the colors of  
the new Poinsettia. The photograph comprises a side per-  
spective view of a single flowering plant of ‘NPCW02044’  
grown in a container.

**DETAILED BOTANICAL DESCRIPTION**

Plants of the new Poinsettia have not been observed under  
all possible environmental conditions. The phenotype may  
vary somewhat with variations in environment such as  
temperature, daylength and light intensity, without,  
however, any variance in genotype.

The aforementioned photograph, following observations  
and measurements describe plants grown in Stuttgart, Ger-  
many under commercial practice in a glass-covered green-  
house with day temperatures ranging from 20 to 30° C.,  
night temperatures ranging from 16 to 20° C., and a maxi-  
mum light level of about 70 kilolux. Cuttings were harvested  
in August and planted into 12-cm containers when rooted  
and pinched once. Plants were flowered under natural season  
short day/long night conditions. Plants were about 15 weeks



from unrooted cuttings when the photograph and the detailed botanical description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Euphorbia pulcherrima* Willd. cultivar NPCW02044.

Parentage:

*Female, or seed, parent.*—Proprietary selection of *Euphorbia pulcherrima* Willd. identified as code number S 219, not patented.

*Male, or pollen, parent.*—Proprietary selection of *Euphorbia pulcherrima* Willd. identified as code number R 82, not patented.

Propagation:

*Type cutting.*—Vegetative terminal cuttings.

*Time to initiate roots.*—Summer: About 12 days at 22° C. Winter: About 15 days at 22° C.

*Time to produce a rooted cutting.*—Summer: About 20 days at 22° C. Winter: About 23 days at 22° C.

*Root description.*—Fibrous; freely-branching; white in color.

Plant description:

*Growth habit.*—Compact, upright and uniform plant habit; inverted triangle; top of plant rounded; low vigor.

*Plant height.*—About 20 to 25 cm.

*Plant diameter or spread.*—About 25 to 30 cm.

*Lateral branch description.*—Quantity per plant: About four to six lateral branches develop after pinching. Length: About 20 cm. Diameter: About 5 to 10 mm. Internode length: About 1 to 4 cm. Texture: Smooth, glabrous. Color: 143A overlain with 183A.

*Foliage description.*—Arrangement: Alternate, single. Quantity of leaves: About 8 to 15 per lateral branch. Length: About 4 to 10 cm. Width: About 2 to 7 cm. Shape: Ovate. Apex: Acute. Base: Obtuse. Margin: Entire with slight irregular lobing. Venation pattern: Pinnate. Texture, upper and lower surfaces: Glabrous, smooth. Color: Developing foliage, upper surface: 144B. Developing foliage, lower surface: 143C. Fully expanded foliage, upper surface: 139A. Fully expanded foliage, lower surface: 137B. Venation, upper surface: 139A. Venation, lower surface: 181A to 139D. Petiole: Length: About 2 to 4 cm. Diameter: About 2 to 3 mm. Texture: Glabrous, smooth. Color, upper and lower surfaces: 183A.

Inflorescence description:

*Inflorescence type and habit.*—Inflorescences are compound corymbs of cyathia with colored flower bracts subtending the cyathia. Inflorescences not persistent.

*Fragrance.*—None detected.

*Natural flowering season.*—Autumn/winter; flower initiation and development is induced under short day/long night conditions.

*Response time.*—Early flowering, about eight weeks.

*Post-production longevity.*—Plants of the new Poinsettia maintain good substance and bract color for about three to four weeks under interior conditions and about four to five weeks under greenhouse conditions.

*Quantity of inflorescences.*—One per lateral branch; about four to six per plant.

*Inflorescence size.*—Diameter: About 15 to 20 cm. Height (depth): About 2 to 5 cm.

*Flower bracts.*—Quantity of fully developed flower bracts: About six to eight per inflorescence. Length: About 7 to 10 cm. Width: About 4 to 6 cm. Shape: Ovate. Apex: Acute. Base: Obtuse. Margin: Entire with occasional lobing. Texture, upper and lower surfaces: Glabrous, smooth. Surface: Concave. Color: Developing and fully expanded bracts, upper surface: 45B; color becoming closer to 46B with development. Developing and fully expanded bracts, lower surface: 46C. Venation, upper and lower surfaces: Same as lamina. Bract petiole: Length: About 1 to 3 cm. Diameter: About 2 mm. Texture: Glabrous, smooth. Color, upper and lower surfaces: 183B.

*Cyathia.*—Quantity of cyathia: About four to seven per corymb. Diameter of cyathia cluster: About 1.5 to 2.5 cm. Length: About 4 to 7 mm. Diameter: About 3 to 5 mm. Shape: Ovoid. Color, immature and mature: 143C.

*Nectaries.*—Quantity of nectaries: One per cyathium. Diameter: About 2 to 3 mm. Color: 13A to 14C.

*Peduncle.*—Length: About 1 to 2 mm. Diameter: About 1 mm. Aspect: Erect. Strength: Moderately strong. Texture: Glabrous, smooth. Color: 144C.

*Reproductive organs.*—Stamens: Quantity of stamens: About five to ten per cyathium. Anther shape: Ovoid. Anther length: About 1 mm. Anther color: 9A to 12A. Amount of pollen: Moderate. Pollen color: 9A to 12A. Pistils: Quantity of pistils: One per cyathium. Pistil length: About 3 to 5 mm. Style length: About 2 to 3 mm. Stigma color: 185A.

Disease/pest resistance: Resistance to pathogens and pests common to Poinsettias has not been observed on plants grown under commercial conditions.

Temperature tolerance: Plants of the new Poinsettia have been observed to tolerate temperatures from 14 to 38° C. It is claimed:

1. A new and distinct cultivar of Poinsettia plant named 'NPCW02044', as illustrated and described.

\* \* \* \* \*

