



US00PP14679P29

(12) **United States Plant Patent**
Brown

(10) **Patent No.: US PP14,679 P2**
(45) **Date of Patent: Apr. 6, 2004**

(54) **DIASCIA PLANT NAMED ‘CODIBLIM’**

(50) Latin Name: *Diascia*×*hybrida*
Varietal Denomination: **Codiblim**

(75) Inventor: **Graham Noel Brown**, Pennant Hills
(AU)

(73) Assignee: **NuFlora International Pty. Ltd.**,
Macquarie Fields (AU)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 61 days.

(21) Appl. No.: **10/259,982**

(22) Filed: **Sep. 29, 2002**

(51) **Int. Cl.⁷** **A01H 5/00**
(52) **U.S. Cl.** **Plt./263**
(58) **Field of Search** **Plt./263**

Primary Examiner—Anne Marie Grunberg
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Diascia* plant named
‘Codiblim’, characterized by its compact and upright plant
habit; freely branching habit; freely flowering habit; white to
light red purple-colored flowers with red purple-colored
centers; and good temperature tolerance.

1 Drawing Sheet

1

Botanical classification/cultivar designation: *Diascia*×*hy-*
brida cultivar Codiblim.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of *Diascia* plant, botanically known as *Diascia*×*hybrida*,
and hereinafter referred to by the cultivar name Codiblim.

The new *Diascia* is a product of a planned breeding
program conducted by the Inventor in Cobbitty, New South
Wales, Australia. The objective of the breeding program is to
create new compact *Diascias* with numerous uniquely-
colored flowers.

The new *Diascia* originated from a cross-pollination made
by the Inventor in 1999 of the *Diascia*×*hybrida* cultivar
Codiap, disclosed in U.S. Plant Pat. No. 13,169, as the
female, or seed parent, with a proprietary *Diascia*×*hybrida*
selection identified as X99.54.9, not patented, as the male, or
pollen parent. The new *Diascia* was selected as a single plant
from the resulting progeny of the cross-pollination by the
Inventor in October, 1999, in an controlled environment in
Cobbitty, New South Wales, Australia.

Asexual reproduction of the new cultivar by terminal
cuttings taken in a controlled environment in Cobbitty, New
South Wales, Australia since October, 1999 has shown that
the unique features of this new *Diascia* are stable and
reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the cultivar Codiblim have not been observed
under all possible environmental conditions. The phenotype
may vary somewhat with variations in environment such as
temperature and light intensity without, however, any vari-
ance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Codib-
lim’. These characteristics in combination distinguish ‘Cod-
iblim’ as a new and distinct cultivar of *Diascia*:

1. Compact and upright plant habit.
2. Freely branching habit.
3. Freely flowering habit.

2

4. White to light red purple-colored flowers with red
purple-colored centers.
5. Good temperature tolerance.

Plants of the new *Diascia* differ primarily from plants of
the parents in flower color as plants of the female parent, the
cultivar Codiap, have soft orange-colored flowers and plants
of the male parent, the selection X99.54.9 have pink-colored
flowers.

Plants of the new *Diascia* can be compared to plants of the
Diascia cultivar Codiaw, disclosed in U.S. Plant Pat. No.
13,058. In side-by-side comparisons conducted in Cobbitty,
New South Wales, Australia, plants of the new *Diascia*
differed from plants of the cultivar Codiaw in the following
characteristics:

1. Plants of the new *Diascia* were shorter than plants of the
cultivar Codiaw.
2. Plant form of the new *Diascia* was fuller and not as
open as plant form of the cultivar Codiaw.
3. Plants of the new *Diascia* and the cultivar Codiaw
differed slightly in flower coloration.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the
overall appearance of the new cultivar, showing the colors as
true as it is reasonably possible to obtain in colored repro-
ductions of this type. Colors in the photographs may differ
slightly from the color values cited in the detailed botanical
description which accurately describe the colors of the new
Diascia.

The photograph at the top of the sheet comprises a side
perspective view of a typical flowering plant of ‘Codiblim’
grown in a container. The photograph at the bottom of the
sheet comprises a close-up view of typical leaves and
flowers of ‘Codiblim’.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observa-
tions and measurements describe plants grown in Encinitas,
Calif., in an outdoor nursery during the winter and spring
with day temperatures averaging 24° C., night temperatures
ranging from 12 to 19° C., and light levels about 4,000 to

6,000 foot candles. Plants were grown for about 16 weeks in 12.7-cm containers and were pinched one time.

Color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Diascia*×*hybrida* cultivar Codiblim.
Parentage:

Female parent.—*Diascia*×*hybrida* cultivar Codiap, disclosed in U.S. Plant Pat. No. 13,169.

Male parent.—*Diascia*×*hybrida* selection identified as X99.54.9, not patented.

Propagation:

Type cutting.—Terminal vegetative cuttings.

Time to initiate roots, summer and winter.—About 7 days at 20° C.

Time to produce a rooted young plant, summer and winter.—About 21 days at 20° C.

Root description.—Fine, fibrous; white in color.

Rooting habit.—Freely branching, dense.

Plant description:

Form.—Upright and compact plant habit. Freely branching with lateral branches potentially forming at every node. Moderately vigorous growth habit.

Plant height.—About 21 cm.

Plant diameter.—About 30 cm.

Lateral branches.—Length: About 21 cm. Diameter: About 2 mm. Internode length: About 1.5 to 2.5 cm. Texture: Smooth, glabrous. Color: 144A.

Foliage description.—Arrangement: Opposite; simple. Quantity per lateral branch: About 17. Length: About 2 cm. Width: About 1.3 cm. Shape: Deltoid to cordate. Apex: Acute. Base: Truncate to cordate. Margin: Slightly serrate towards the base; entire towards the apex. Texture, upper and lower surfaces: Smooth, glabrous. Venation pattern: Pinnate. Color: Young foliage, upper surface: 146A. Young foliage, lower surface: 146B. Fully expanded foliage, upper surface: 147A. Fully expanded foliage, lower surface: 147D. Venation, upper surface: 147B. Venation, lower surface: 147D. Petiole length: About 4 mm. Petiole diameter: About 2 mm. Petiole color: 144A.

Flower description:

Flower type and habit.—Solitary zygomorphic flowers arranged on terminal racemes. Five modified petals fused at base: two upper (banner) petals, two lateral petals, and one larger lower lip petal. Flowers not persistent.

Quantity.—Freely flowering; typically about 16 to 20 buds and flowers per lateral branch. Flowers face mostly outward.

Natural flowering season.—Plants typically flower during the spring in the Northern Hemisphere; flowering continuous during this period.

Flower longevity on the plant.—About four days.

Fragrance.—Not detected.

Raceme size.—Length: About 10 to 12 cm. Width: About 4.5 cm.

Flower size.—Length: About 2.2 cm. Width: About 2.4 cm. Depth: About 1.3 cm.

Flower buds (showing color).—Length: About 5 mm. Diameter: About 6 mm. Shape: Nearly spherical. Color: 155A.

Petals.—Quantity/arrangement: Five modified petals fused at base: two upper (banner) petals, two lateral petals, and one larger lower lip petal. Base of banner petals with concave yellow eyespots; lower surfaces of lateral petals modified into nectar spurs; and lower lip petal convex forming a roughly horizontal insect landing platform. Length: Banner petals: About 7 mm. Lateral petals: About 8 mm. Lower lip petal: About 1.2 cm. Width: Banner petals: About 7 mm. Lateral petals: About 9 mm. Lower lip petal: About 1.5 cm. Lateral petal spur: Length: About 1 cm. Diameter, at petal attachment: About 2 mm. Shape, all petals: Roughly spatulate. Apex, all petals: Rounded. Margin, all petals: Entire. Texture, all petals: Smooth, velvety. Color, all petals: When opening, upper and lower surfaces: 155D. Fully opened, upper surface: 155D; towards base, 67B to 67C; with development, petals become light red purple, close to 65D. Fully opened, lower surface: 155D. Nectar spurs: 74D. Eyespot on banner petals: 5A.

Sepals.—Arrangement/appearance: Single whorl of five sepals fused at base; star-shaped. Length: About 2.5 mm. Width: About 1.5 mm. Shape: Narrowly elliptic. Apex: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: Immature and mature, upper surface: 146A. Immature and mature, lower surface: 146B.

Peduncles.—Length: About 5 cm. Width: About 2 mm. Angle: Upright to about 15° from vertical. Strength: Moderately strong. Texture: Smooth, glabrous. Color: 146A.

Pedicels.—Length: About 1.3 cm. Width: About 1 mm. Angle: About 45 to 60° from the peduncle. Strength: Moderately strong; slender. Texture: Smooth, glabrous. Color: 146A.

Reproductive organs.—Stamens: Quantity per flower: Four. Anther shape: Ovoid. Anther length: Less than 1 mm. Anther color: 14A. Pollen amount: Scarce. Pollen color: 14A. Pistils: Quantity per flower: One. Pistil length: About 4 mm. Style length: About 2 mm. Style color: 145B. Stigma shape: Rounded. Stigma color: 145B. Ovary color: 145A.

Fruit/seed.—Fruit and seed production has not been observed.

Disease/pest resistance: Plants of the new *Diascia* have not been noted to be resistant to pathogens or pests common to *Diascia*.

Temperature tolerance: Plants of the new *Diascia* have been observed to tolerate temperatures from 0 to 35° C.

It is claimed:

1. A new and distinct cultivar of *Diascia* plant named ‘Codiblim’, as illustrated and described.

* * * * *

