



US00PP14781P2

(12) **United States Plant Patent**  
**Fear et al.**

(10) **Patent No.:** **US PP14,781 P2**

(45) **Date of Patent:** **May 11, 2004**

(54) **RASPBERRY PLANT NAMED ‘DRISCOLL MADONNA’**

(50) Latin Name: *Rubus idaeus L.*  
Varietal Denomination: **Driscoll Madonna**

(75) Inventors: **Carlos D. Fear**, Aptos, CA (US);  
**Richard E. Harrison**, Aptos, CA (US);  
**Fred M. Cook**, Aptos, CA (US); **Gavin Sills**, Watsonville, CA (US)

(73) Assignee: **Driscoll Strawberry Associates, Inc.**,  
Watsonville, CA (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/306,453**

(22) Filed: **Nov. 27, 2002**

(51) **Int. Cl.**<sup>7</sup> ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./204**

(58) **Field of Search** ..... **Plt./204**

*Primary Examiner*—Anne Marie Grunberg

*Assistant Examiner*—Susan B. McCormick

(74) *Attorney, Agent, or Firm*—Pennie & Edmonds LLP

(57) **ABSTRACT**

The present invention relates to a new and distinct cultivar of raspberry plant named Driscoll Madonna. The new cultivar is distinguished from other raspberry cultivars by its large fruit with excellent fruit firmness and structure. The new cultivar is distinguished from its seed parent by having better flavor. The new cultivar is distinguished from its pollen parent by producing larger, firmer fruit.

**3 Drawing Sheets**

**1**

Latin name of the genus and species of the plant claimed: The variety is botanically identified as *Rubus idaeus L.*

**BACKGROUND OF THE INVENTION**

The new cultivar of raspberry plant was developed from the hybridization of the selection of ‘R652.1’ (an unpatented variety) as the seed parent with the selection ‘R709.1’ (an unpatented variety) as the pollen parent. The parents were crossed in 1997, whereafter fruit and seed were collected to produce seedlings for field planting in Watsonville, Calif. in 1997. The new cultivar was selected from these seedlings in 1998 for its attractive and large fruit with long pedicels. The new cultivar has been asexually propagated by in vitro shoot tip culture, root sucker division and root cuttings at the Cassin Ranch in Santa Cruz county, Calif. and has been shown to maintain the desired and distinguishing characteristics after propagation over several generations.

**SUMMARY OF THE INVENTION**

The present invention provides a new and distinct cultivar of red raspberry plant named ‘Driscoll Madonna’. The cultivar is botanically identified as *Rubus idaeus L.* The ‘Driscoll Madonna’ red raspberry plant produces a primocane crop which begins in early August and continues until early November. The floricanes crop begins in late May and continues until mid to late July. The fruit of ‘Driscoll Madonna’ is notable for its consistent large size, uniform shape and long pedicels. The fruit of ‘Driscoll Madonna’ does not separate easily from its receptacle.

**BRIEF DESCRIPTION OF THE DRAWINGS**

The accompanying photographs show typical specimens of the primocane fruit, leaves and shoot of the new cultivar, in color as nearly true as it is reasonably possible to make in color illustrations of these characteristics.

FIG. 1 is a photograph of ‘Driscoll Madonna’ primocane flower and fruit in various stages of development.

FIG. 2 is a photograph of ‘Driscoll Madonna’ primocane leaves showing upper and lower surfaces.

**2**

FIG. 3 is a photograph of ‘Driscoll Madonna’ primocane shoot.

**DETAILED BOTANICAL DESCRIPTION**

The following detailed description of the new raspberry cultivar, ‘Driscoll Madonna’, is based upon observations taken of 7 to 17 month old plants and fruit grown in Watsonville, Calif. between 2001 and 2002, and is believed to apply to plants of the ‘Driscoll Madonna’ cultivar grown in similar conditions of soil and climate elsewhere.

Throughout this specification, color names beginning with a small letter signify that the name of the color, as used in common speech, is aptly descriptive. Color data followed by an alphanumeric code designates the color according to The R.H.S. Colour Chart published by The Royal Horticultural Society of London, England. Color designations, color descriptions, and other phenotypical descriptions may deviate from the stated values and descriptions depending upon variation in environmental, seasonal, climatic and cultural conditions.

Table 1 provides information on the plant and fruit characteristics of the new cultivar ‘Driscoll Madonna’ compared with characteristics of the unpatented raspberry cultivar ‘Heritage’. Observations of the cultivars were taken under similar conditions.

The new variety is particularly characterized and distinguished from other cultivars by its large fruit with excellent fruit firmness and structure. The fruit is produced on long pedicels and is difficult to release from the receptacle.

The fruit color of ‘Driscoll Madonna’ is a bright red at harvest. Fruit of ‘Driscoll Madonna’ does not separate easily from the receptacle and is of good firmness at harvest. The fruit of ‘Driscoll Madonna’ is consistent in size and shape throughout the harvest period. The average plant height is about 240 cm and the average plant spread is about 50 cm. The pigmentation of the young shoots is 144B and there were an average of 6 young shoots in the observed plants of ‘Driscoll Madonna’. The pedicel color is 144A.

The reproductive organs of 'Driscoll Madonna' are variable. The color of both surfaces of the petals is 155D and there are five petals per flower. The style color is 157D, the average number of styles per flower is about 84, the anther color is 155D, and the average number of anthers per flower is about 97. The number of petals per flower is five. The color of the seeds of 'Driscoll Madonna' is 161A, the average seed weight is about 1.2 mg, and there are an average of about 115 seeds per fruit.

The florican yield of 'Driscoll Madonna' is high relative to the variety 'Heritage'. 'Driscoll Madonna' is distinguishable from its pollen parent, selection 'R709.1', by producing larger, firmer fruit. The new cultivar is distinguished from its seed parent, selection 'R652.1', by having better flavor.

DISEASE AND STRESS RESISTANCE

Cold tolerance of the new cultivar has not been established. Post harvest fruit rot resistance is good in comparison over many selections and varieties.

TABLE 1

PLANT CHARACTERISTICS OF 'DRISCOLL MADONNA'		
	Driscoll Madonna	Heritage
<u>General</u>		
Plant size	Large	Large
Growth habit	Semi-erect	Erect
Productivity	High	Medium
Self-fruitfulness	Self-fruitful	Self-fruitful
Time of bud burst	Late	Late
<u>Primocane fruiting</u>		
Percent of cane length flowering as primocane	5-30	20-40
Percent of total yield Primocanes	44	53
<u>Number of young shoots</u>		
Young shoot pigmentation	Medium	Medium
Length (cm)	232	196
Time of shoot emergence	Late	Very late
<u>Glaucoisity (waxy bloom)</u>		
Strength	Weak	Weak
Cane Cross section	Medium	Medium
from mid cane of primocane)	Rounded to angular	Rounded
Dormant cane color	tan w/slight purple	brown to purple brown
<u>Prickles</u>		
Pigmentation	purple	green-brownish to green
Density on young shoots	Medium	Dense
Attitude of tip	Horizontal	Downward
Size	Medium	Medium
Size: Length (base to tip at 1 m height at end of season) (mm)	1	2.3
Texture	smooth	Rigid
Presence and distribution on petioles	Present irregularly distributed	Present irregularly distributed
Pubescence on canes	Absent	Absent
Internodal distance (cm) (at central 1/3 of cane)	6.0	5.3
<u>LEAVES</u>		
<u>Color</u>		
Face	147A	147A
Underside	148C	148B
Relief between veins	Medium	Very weak
Glossiness	Medium	Medium

TABLE 1-continued

PLANT CHARACTERISTICS OF 'DRISCOLL MADONNA'		
	Driscoll Madonna	Heritage
<u>Petiole length (cm)</u>		
Petiole length (cm)	6.2	7.7
<u>Stipule orientation</u>		
Stipule orientation	Erect	Erect
<u>Arrangement</u>		
Arrangement	Compound	Compound
<u>Number of leaflets</u>		
Number of leaflets	Usually 5	Sometimes 3, sometimes 5
<u>Overlapping of lateral leaflets</u>		
Overlapping of lateral leaflets	Overlapping	Free to touching
<u>Lateral leaflet: length of stalket (lower pair)</u>		
Lateral leaflet: length of stalket (lower pair)	Medium	Very short
<u>Terminal leaflet</u>		
<u>Length (cm)</u>		
Length (cm)	11.9	14.6
<u>Width (cm)</u>		
Width (cm)	8.4	7.8
<u>Shape</u>		
Shape	Ovate	Ovate
<u>Tip</u>		
Tip	Acuminate	Acuminate
<u>Base</u>		
Base	Round to cordate	Acute to rounded
<u>Margin</u>		
Margin	Doubly serrate	Doubly serrate
<u>Lateral leaflets (basal pair)</u>		
<u>Length (cm)</u>		
Length (cm)	10.9	14.7
<u>Width</u>		
Width	8.1	8.6
<u>Orientation</u>		
Orientation	Opposite	Opposite
<u>Shape</u>		
Shape	Ovate	Ovate
<u>Tip</u>		
Tip	Acuminate	Acuminate
<u>Base</u>		
Base	Round	Oblique
<u>Rachis length between terminal leaflet and adjacent lateral leaflets (cm)</u>		
Rachis length between terminal leaflet and adjacent lateral leaflets (cm)	3.8	1.5
<u>Margin</u>		
Margin	Doubly serrate	Doubly serrate
<u>FLOWERS</u>		
<u>Flowering period</u>		
<u>Primocane</u>		
Primocane	14 weeks, Late June-late September	19 weeks, Late May-late September
<u>Florican</u>		
Florican	9 weeks, Early April-mid June	10 weeks, Late March-mid June
<u>Flower diameter (cm)</u>		
Flower diameter (cm)	1.4	1.8
<u>Petal</u>		
<u>Length (cm)</u>		
Length (cm)	0.8	0.8
<u>Width (cm)</u>		
Width (cm)	0.4	0.3
<u>Pedicel coloration</u>		
Pedicel coloration	Present, medium intensity	Present, strong intensity
<u>FRUIT</u>		
<u>Harvest season</u>		
<u>Primocane</u>		
Primocane	Mid August-late Oct	Early July-early November
<u>Florican</u>		
Florican	Late May-mid July	Late May-late July
<u>Fruting lateral</u>		
<u>Length (4<sup>th</sup> lateral from tip) (cm)</u>		
Length (4 <sup>th</sup> lateral from tip) (cm)	67.8	49.8
<u>Number of fruit per lateral</u>		
Number of fruit per lateral	12.2	20.3
<u>Color</u>		
<u>Immature</u>		
Immature	46C	42C
<u>Maturing</u>		
Maturing	46A	46A
<u>Mature fruit</u>		
Mature fruit	59A	59A
<u>Glossiness</u>		
Glossiness	Medium	Medium
<u>Shape</u>		
Shape	Ovate-elliptic	Ovate
<u>Dimensions</u>		
<u>Size</u>		
Size	Large	Small
<u>Length (mm)</u>		
Length (mm)	30	17
<u>Width (mm)</u>		
Width (mm)	22	18
<u>Length:width ratio</u>		
Length:width ratio	1.36	.94
<u>Weight (g/fruit)</u>		
<u>Primocane</u>		
Primocane	7.5	3.1
<u>Florican</u>		
Florican	6.6	2.3
<u>Soluble solids (%)</u>		
Soluble solids (%)	11.5	10.8

TABLE 1-continued

<u>PLANT CHARACTERISTICS OF 'DRISCOLL MADONNA'</u>		
	Driscoll Madonna	Heritage
Titrateable acidity (% as citric acid)	1.54	1.58
Seed Weight (mg)	2.8	1.5
Number drupelets/fruit	115	72
Adherence to plug	Strong	Medium
Firmness	Medium	Firm
Yield	High	Medium

#### NUCLEIC ACID FINGERPRINTING

Distinctive patterns of polymorphism can be detected using a variety of nucleic acid analysis methods. In one

non-limiting example, molecular genetic maps can be produced using random amplified polymorphic DNA (RAPD) (Williams et al., 1990, "DNA polymorphisms amplified by arbitrary primers are useful as genetic markers", *Nucleic Acids Res.* 18(22):6531-5). Using a variety of oligonucleotide primers, alone or in combination, RAPD analysis of Driscoll Madonna and Heritage yielded DNA fragment patterns that uniquely distinguish each of these genetically distinct genotypes.

We claim:

1. A new and distinctive cultivar of raspberry plant, substantially as shown and described.

\* \* \* \* \*




FIG. 1




FIG. 2




FIG. 3