



US00PP14922P2

(12) **United States Plant Patent**  
**Noodeljik**(10) **Patent No.:** **US PP14,922 P2**  
(45) **Date of Patent:** **Jun. 22, 2004**(54) **CHrysanthemum NAMED 'HOGGAR'**(50) Latin Name: *dendranthema grandiflora*  
Varietal Denomination: Hoggar(75) Inventor: **Robert Noodeljik**, Woubrugge (NL)(73) Assignee: **Chrysanthemum Breeders  
Association, N.V. (NL)**

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/561,926**(22) Filed: **May 1, 2000**(51) Int. Cl.<sup>7</sup> ..... **A01H 5/00**(52) U.S. Cl. ..... **Plt./295**(58) Field of Search ..... **Plt./295**

Primary Examiner—Bruce R. Campell

Assistant Examiner—Annette H. Para

(74) Attorney, Agent, or Firm—Parkhurst &amp; Wendel, L.L.P.

(57) **ABSTRACT**

A Chrysanthemum plant named 'HOGGAR' characterized by its medium sized blooms with yellow ray-florets and yellow-green disc florets.

**2 Drawing Sheets****1****BACKGROUND OF THE INVENTION**

'HOGGAR' is a product of a breeding-program which had the objective of creating new chrysanthemum cultivars with a daisy type flower, a 7 week response and a medium plant height. The new plant of the present invention comprises a new and distinct cultivar of Chrysanthemum plant. 'HOGGAR' is a seedling from a cross in a breeding program maintained under the control of inventor. The female parent is #93.1028—unpatented—, the male parent is #93.0910—unpatented—, both unnamed seedlings not available to inventor for description. The new and distinct cultivar was discovered and selected as a flowering plant within the progeny of the stated cross by Rob Noodeljik in a controlled environment (greenhouse) in Rijsenhout Holland in April 1998. The first act of asexual reproduction of 'HOGGAR' was accomplished when vegetative cuttings were taken from the initial selection in July 1998 in a controlled environment in Rijsenhout Holland.

**SUMMARY OF THE INVENTION**

The present invention is a new and distinct variety of chrysanthemum bearing medium sized blooms with yellow ray-florets and yellow-green disc florets.

**BRIEF DESCRIPTION OF THE DRAWINGS**

The present invention of a new and distinct variety of chrysanthemum is shown in the accompanying drawings, the color being as nearly true as possible with color photographs of this type.

FIG. 1 shows a plant of the cultivar in full bloom.

FIG. 2 shows the various stages of bloom of the new cultivar;

FIG. 3 shows the various stages of foliage and petiole of the new cultivar.

**DESCRIPTION OF THE INVENTION**

This new variety of chrysanthemum is of the botanical classification *dendranthema grandiflora*. The observations and measurements were gathered from 64 days plants grown in a greenhouse in Rijsenhout Holland in a photo-periodic controlled crop under conditions generally used in commercial practice. The greenhouse temperatures during this crop

**2**

were at day-time between 18° C. and 25° C. and at night 20° C. The photo-periodic response time in this crop was 48 days after an average of eight long days. After this long day period to flowering growth retardants were applied 6 times in an average dose of 1.5 gram/liter water. No tests were done on disease or insect resistance or susceptibility. This new variety produces medium sized blooms with yellow ray-florets and yellow-green disc-florets blooming on the plant for 5 weeks. This new variety of chrysanthemum has been found to retain its distinctive characteristics throughout successive propagations however the phenotype may vary significantly with variations in environment such as light intensity and temperature. To show the phenotype as described 'HOGGAR' can be planted without assimilation lightning (high pressure sodium lamps) between week 50 and week 40 of the next year under greenhouse conditions in Holland. With assimilation lightning (minimum level 2500 lux) it can be planted year round under greenhouse conditions in Holland.

From cultivars known to inventor not one has a bronze underside in combination with yellow ray-florets. This is a special distinguishing characteristic of 'HOGGAR'. The most similar existing cultivar in comparison to 'HOGGAR' is 'Miramar' (U.S. Plant Pat. No. 7,469). When 'Miramar' and 'HOGGAR' are being compared the following differences are noticed: Obvious differences of 'Miramar' and 'HOGGAR' are the bud color and the bronze underside of the yellow ray-florets. In vigour, plant size and flower size 'HOGGAR' and 'Miramar' are comparable.

The following is a description of the plant and characteristics that distinguish 'HOGGAR' as a new and distinct variety.

The color designations are taken from the plant itself. Accordingly, any discrepancies between the color designations and the colors depicted in the photographs are due to photographic tolerances. The color chart used in this description is: The Royal Horticultural Society Colour Chart, edition 1995.

**Botanical Description of Cultivar 'Hoggar'****Bud:**

*Size.*—Large; cross-section 1.3 cm, height 1.0 cm.

*Outside color.*—Greyed-orange 171 A.

*Involucral bracts.*—2 rows, length 7 mm, width 3 mm.

*Involucral bracts among disc-florets.*—Not present.  
*Involucral bracts color.*—Green 138 B.  
 Bloom:  
*Type.*—Daisy.  
*Height.*—Flat.  
*Size.*—Medium.  
*Fully expanded.*—6.0–6.5 cm.  
*Number of blooms per branch.*—Approx. 4 blooms per branch.  
*Performance on the plant.*—4 weeks.  
*Seeds.*—Not produced.  
*Fragrance.*—Typical chrysanthemum.  
 Color:  
*Center of the flower (disc-florets).*—Immature yellow-green 144 B. Mature yellow-orange 14 A and yellow-green 144 B in the middle.  
*Color of upper surface of the ray-florets.*—Yellow 12 A, in spring+autumn greyed-red 171 A along the edge.  
*Color of the lower surface of the ray-florets.*—Yellow 10 A with a pattern of greyed-red 171 D.  
*Tonality from distance.*—A pot mum with medium yellow daisy flowers and bronze buds.  
*Discoloration to color.*—Yellow 10 A.  
 Ray florets:  
*Texture.*—Upper and under side smooth.  
*Number.*—26–30 (3 rows).  
*Cross-section.*—Flat.  
*Longitudinal axis of majority.*—Reflexing.  
*Length of corolla tube.*—Short.  
*Ray-floret margin.*—Entire.  
*Ray-floret length.*—3.2–3.5 cm.  
*Ray-floret width.*—1.0–1.2 cm.  
*Ratio length/width.*—Medium.  
*Shape of tip.*—Dentate.  
 Disc florets:  
*Disc diameter.*—1.2–1.4 cm.  
*Distribution of disc florets.*—Numerous clearly visible at all stages of flowering.  
*Shape.*—Tubular.  
*Color.*—Yellow-green 144 B.  
*Receptacle shape.*—Conical raised.  
 Reproductive organs:  
*Stamen (present in disc florets only).*—Yellow-green, thick, 3 mm. In length.  
*Number of stamen.*—Grown together as one.  
*Stamen color.*—Yellow-green 144 B.  
*Pollen color.*—Yellow-orange 13 A.  
*Styles (present in both ray and disc florets).*—Yellow-green, thick.  
*Style color.*—Yellow-green 144 B.  
*Style length.*—4 mm.

*Stigmas.*—Yellow-green.  
*Stigma width.*—1 mm.  
*Ovaries.*—Enclosed in calyx.  
 Plant:  
*Form.*—A pot mum meant for indoor use.  
*Growth habit.*—Spreading.  
*Growth rate.*—Rapid.  
*Height.*—23–25 cm.  
*Width.*—25–27 cm.  
*Stem color.*—Green 138 B.  
*Stem strength.*—Weak.  
*Stem brittleness.*—Present.  
*Stem anthocyanin coloration.*—Absent.  
*Length of lateral branch.*—From top to bottom 12–14 cm.  
*Lateral branch color.*—Green 138 B.  
*Lateral branch, attachment.*—Weak.  
*Branching (average number of lateral branches).*—Normal with 3–4 breaks after pinching.  
*Peduncle length.*—4 cm.  
*Peduncle color.*—Green 138 B.  
*Flowering response (photo-periodic controlled crop, not natural season).*—48 Days.  
 Foliage:  
*Color.*—Upper side yellow-green 137 A. Under side yellow-green 147 B.  
*Size.*—Medium; length 7.5 cm, width 4.0–4.5 cm.  
*Quantity (number per lateral branch).*—6–7.  
*Shape.*—Oval.  
*Texture upper side.*—Glabrous.  
*Texture under side.*—Pubescent.  
*Venation arrangement.*—Palmate.  
*Shape of the margin.*—Serrated.  
*Shape of base of sinus between lateral lobes.*—Round.  
*Margin of sinus between lateral lobes.*—Mostly diverging.  
*Shape of base.*—Acute.  
*Apex.*—Mucronate.

## Differences with the comparison varieties

	‘HOGGAR’	‘MIRAMAR’
Bud color	Greyed-oragane 171 A	Yellow 4 A
Underside color	ray-floret Yellow 10 A with a pattern or greyed-red 171 D	Yellow 9 C

## I claim:

1. A new and distinct variety of chrysanthemum plant as described and illustrated.


\* \* \* \* \*



**FIG. 1**



**FIG. 2**



**FIG. 3**