



US00PP14939P2

(12) **United States Plant Patent**
Noodelijk

(10) **Patent No.: US PP14,939 P2**
(45) **Date of Patent: Jun. 22, 2004**

(54) **CHRYSANTHEMUM PLANT NAMED
'CEFREYA'**

(50) Latin Name: *Chrysanthemum morifolium*
Varietal Denomination: **Cefreya**

(75) Inventor: **Robert Noodelijk**, Woubrugge (NL)

(73) Assignee: **Chrysanthemum Breeders Association
N.V. (NL)**

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 30 days.

(21) Appl. No.: **10/316,870**

(22) Filed: **Dec. 12, 2002**

(51) **Int. Cl.⁷** **A01H 5/00**

(52) **U.S. Cl.** **Plt./297**

(58) **Field of Search** Plt./297

(56) **References Cited**

PUBLICATIONS

UPOV-ROM Plant Variety Database, hit on 'Freya', 2003/
03.*

* cited by examiner

Primary Examiner—Anne Marie Grunberg

(74) *Attorney, Agent, or Firm*—Parkhurst & Wendel, L.L.P.

(57) **ABSTRACT**

A chrysanthemum plant named 'Cefreya' characterized by
its medium sized blooms with coral-salmon ray florets and
prolific branching, natural season flower date September
1–5; blooming for a period of 5 weeks.

3 Drawing Sheets

1

BACKGROUND OF THE INVENTION

'Cefreya' is a product of a breeding and selection program
for outdoor pot mums (garden mums) that had the objective
of creating new chrysanthemum cultivars with a decorative
type flower, a natural season flower date around September
1–5, and a blooming for a period of 5 weeks. The new plant
of the present invention comprises a new and distinct
cultivar of Chrysanthemum plant 'Cefreya', which is a
seedling resulting from the open pollination among groups
of chrysanthemum cultivars maintained under the control of
the inventor for breeding purposes. The identity of the parent
plants yielding this new chrysanthemum plant is unknown.
The new and distinct cultivar was discovered and selected as
one flowering plant by Rob Noodelijk on a cultivated field
in Rijsenhout, Holland in September 2000. The plant has
been asexually reproduced by cuttings in greenhouses at
Rijsenhout, Holland. The new cultivar has been found to
retain its distinctive characteristics through successive
propagations.

BRIEF DESCRIPTION OF THE DRAWINGS

The present invention of a new and distinct variety of
chrysanthemum is shown in the accompanying drawings,
the color being as nearly true as possible with color photo-
graphs of this type.

FIG. 1 shows a plant of the cultivar in full bloom.

FIG. 2 shows the various stages of bloom of the new
cultivar.

FIG. 3 shows the petiole and foliage of the new cultivar.

DESCRIPTION OF THE INVENTION

This new variety of chrysanthemum is of the botanical
classification *Chrysanthemum morifolium*. The observations
and measurements were gathered from plants grown out
door in Rijsenhout, Holland under natural day length and
temperature and planted week 22 in 2000 and 2001. The
natural blooming date of this crop was September 1–5 (week

2

35). The average height of the plants was 27 cms. No growth
retardants were used. No tests were done on disease or insect
resistance or susceptibility. No tests were done on cold or
drought tolerance. This new variety produces medium sized
blooms with coral-salmon ray florets and a red center
blooming for a period of 5 weeks.

From the cultivars known to inventor the most similar
existing cultivar in comparison to 'Cefreya' is 'Grenadine'
(U.S. Plant Pat. No. 5,338). When 'Grenadine' and 'Cefreya'
are being compared the following differences are noticed:
The differences of 'Grenadine' and 'Cefreya' are (1) Flower
form. Both 'Cefreya' and 'Grenadine' have a coral-salmon
decorative flower; however the ray-florets differ in form. (2)
Natural flowering date. 'Cefreya' flowers earlier.

The following is a description of the plant and character-
istics that distinguish 'Cefreya' as a new and distinct variety.
The color designations are taken from the plant itself.
Accordingly, any discrepancies between the color designa-
tions and the colors depicted in the photographs are due to
photographic tolerances. The color chart used in this
description is: The Royal Horticultural Society Colour
Chart, 1995 edition.

Table 1. Botanical Description of Cultivar
'Cefreya'

Bud:

Size.—Medium; cross-section 1.0 cm, height 1.5 cm.

Outside color.—Red 37B.

Involucral bracts.—2 rows, length 7 mm, width 3 mm.

Involucral bracts among disc-florets.—Not present.

Involucral bracts color.—Green 138B.

Bloom:

Type.—Decorative.

Height.—High; 2.5 cm.

Size.—Medium.

Fully expanded.—5.5–6.0 cm.

Number of blooms per branch.—Approx. 4–6 blooms
per branch.

Performance on the plant.—5 weeks.
Seeds (if crossed).—Produced in small quantities, ovate grey-brown 199A, 1½ mm in length.
Fragrance.—Typical chrysanthemum, slightly.

Color:
Center of the flower.—Immature red 37A. Mature red 37C.
Color of upper surface of the ray-florets.—Red 37B.
Color of the lower surface of the ray-florets.—Red 37C.
Tonality from distance.—A garden mum with coral-salmon flowers and a red disc.
Color of the upper surface of the ray-florets after aging of the plant.—Red 37C.

Ray florets:
Texture.—Upper and under side smooth.
Number.—140–150.
Cross-section.—Flat.
Longitudinal axis of majority.—Reflexing to twisted.
Length of corolla tube.—Short 0.3 cm.
Ray-floret margin.—Entire.
Ray-floret length.—2.6–2.8 cm.
Ray-floret width.—0.6 cm.
Ratio length/width.—High.
Shape of tip.—Rounded.
Disc florets.—None.
Receptacle shape.—Conical raised.

Reproductive organs:
Stamen.—Absent.
Pollen.—None.
Styles.—Thin.
Style color.—Yellow-green 144A.
Style length.—4 mm.
Stigmas.—Yellow-green 144A.
Stigma width.—1 mm.
Ovaries.—Enclosed in calyx.

Plant:
Form.—Grown as spray-type pot mum, outdoor mounded and round.
Growth habit.—Spreading.
Growth rate.—Moderate.
Height.—27 cm.

Width.—32–34 cm.
Stem color.—Green 139B.
Stem strength.—Strong.
Stem brittleness.—Present.
Stem anthocyanin coloration.—Absent.
Length of lateral branch.—From top to bottom 13–14 cm.
Lateral branch color.—Green 139B.
Lateral branch, attachment.—Flexible.
Branching (average number of lateral branches).—Prolific with 7–8 breaks after pinching.
Peduncle length.—6.0 cm.
Peduncle color.—Green 139B.
Natural season blooming date.—September 1–5.

Foliage:
Color.—Upper side green 139A. Under side green 139C.
Size.—Medium; length 5.0 cm, width 4.0 cm.
Quantity (number per lateral branch).—8–9.
Shape.—Ovate.
Texture upper side.—Glabrous.
Texture under side.—Pubescent.
Venation arrangement.—Palmate.
Shape of the margin.—Serrated.
Shape of base of sinus between lateral lobes.—Round.
Margin of sinus between lateral lobes.—Converging.
Shape of base.—Truncate.
Apex.—Cuspidate.

TABLE 2

Differences with the comparison varieties		
	‘Cefreya’	‘GRENADINE’
Longitudinal axis of ray-florets	Reflexing to twisted	Incurved
Cross section of ray-florets	Flat	Concave
Natural flowering date	September 1–5	September 17–25

I claim:

1. A new and distinct variety of chrysanthemum plant as described and illustrated.

* * * * *




FIG. 1




FIG. 2




FIG. 3