



US00PP15185P3

(12) **United States Plant Patent**  
**Hancock**

(10) **Patent No.:** **US PP15,185 P2**  
(45) **Date of Patent:** **Sep. 28, 2004**

(54) **BLUEBERRY PLANT DENOMINATED**  
**'AURORA'**

(52) **U.S. Cl.** ..... **Plt./157**

(58) **Field of Search** ..... **Plt./157**

(50) Latin Name: *Vaccinium corymbosum*  
Varietal Denomination: **Aurora**

*Primary Examiner*—Anne Marie Grunberg

*Assistant Examiner*—S B McCormick-Ewoldt

(75) Inventor: **James F. Hancock**, East Lansing, MI  
(US)

(74) *Attorney, Agent, or Firm*—Harness, Dickey & Pierce,  
P.L.C.

(73) Assignee: **Board of Trustees operating Michigan**  
**State University**, East Lansing, MI  
(US)

(57) **ABSTRACT**

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

Aurora is a new blueberry variety of *Vaccinium corymbo-  
sum*. It is a productive, very late ripening variety with high  
fresh market quality intended for areas where northern  
highbush varieties are grown successfully. Plants of Aurora  
are vigorous and upright. Canes are numerous, moderately  
branched and the fruit are well exposed. Its berries are  
moderately large, have small, dry picking scars, excellent  
powder-blue color, delicious flavor and excellent firmness.  
Aurora has a harvest season that begins about 5 days after  
the variety known as 'Elliott.'

(21) Appl. No.: **10/350,345**

(22) Filed: **Jan. 23, 2003**

(65) **Prior Publication Data**

US 2004/0148671 P1 Jul. 29, 2004

(51) **Int. Cl.**<sup>7</sup> ..... **A01H 5/00**

**2 Drawing Sheets**

**1**

Latin name and variety denomination: The present inven-  
tion relates to a new and distinct variety of *Vaccinium*  
*corymbosum*, which is hereby denominated 'Aurora.'

**CROSS-REFERENCE TO RELATED  
APPLICATIONS**

This application is related to the present inventor's United  
States Patent Applications entitled "Blueberry plant denomi-  
nated 'Liberty,'" U.S. patent application Ser. No. 10/350,  
343, filed Jan. 23, 2003, and "Blueberry plant denominated  
'Draper,'" U.S. patent application Ser. No. 10/350,354, filed  
Jan. 23, 2003. Both of these applications are also assigned  
to the assignee of the present application. The disclosures of  
the above applications are hereby incorporated herein by  
reference.

**SUMMARY OF THE INVENTION**

The present invention relates to a new and distinct variety  
ofighbush blueberry plant, denominated 'Aurora.' Aurora  
is a new blueberry variety of *Vaccinium corymbosum* from  
the Michigan State University breeding program. It is a  
productive, very late ripening variety with high fresh market  
quality that is intended for areas where northernighbush  
varieties are grown successfully. Plants of Aurora are vig-  
orous and upright. Canes are numerous, moderately  
branched and the fruit are well exposed. Its berries are  
moderately large, have small, dry picking scars, excellent  
powder-blue color, delicious flavor and excellent firmness.  
Aurora has a harvest season that begins about 5 days after  
the variety known as 'Elliott.' 'Aurora' ripens 10–14 days  
after 'Brigetta'. Blueberry growers often receive their high-  
est profits in the late season. Aurora will provide a new late  
ripening option with high quality fruit.

**2**

Emasculated flowers of the 'Brigetta'ighbush blueberry  
from Australia were pollinated in 1991 with pollen from the  
northernighbush variety Elliott. The seeds were  
germinated, grown in a greenhouse for 1 year and then field  
planted at Benton Harbor, Mich. Aurora was selected from  
a group of 54 siblings in 1997. The original selection has  
been evaluated at Benton Harbor, Mich. annually for 11  
years. Asexual reproduction took place at East Lansing,  
Mich. and Lowell, Oreg. Aurora has been propagated by  
hardwood cuttings that produced over a hundred shoots that  
were rooted in the greenhouse and then planted in the field.  
Initiation of rood development from hardwood cuttings  
takes about two to about three months. In addition, Aurora  
has been propagated by softwood cuttings that have been  
rooted. Furthermore, generation of microshoots in a green-  
house using established tissue culture methods has produced  
thousands of clones of Aurora. Initiation of root develop-  
ment from microshoots takes about three to about four  
weeks. Such methods are discussed in the following  
references, which are incorporated by reference in their  
entirety: Doran, W. L. and Bailey, J. S. "Propagation of the  
high bush blueberry by softwood cuttings," Bulletin Mas-  
sachusetts Agricultural Experiment Station; no. 410.  
Amherst, Mass. Massachusetts State College, 1943;  
Doehlert, C. A. "Propagating blueberries from hardwood  
cuttings," Circular (New Jersey Agricultural Experiment  
Station) 490. New Brunswick, N.J. New Jersey Agricultural  
Experiment Station, 1945; Doehlert, C. A. "Propagating  
blueberries from hardwood cuttings," Circular (New Jersey  
Agricultural Experiment Station) 551. New Brunswick,  
N.J.: New Jersey Agricultural Experiment Station, 1953; El  
Shiekh, A.; Wildung, D. K.; Luby, J. J.; Sargent, K. L.; Read,  
P. E. "Long term effects of propagation by tissue culture or  
softwood single node cuttings on growth habit, yield, and  
berry weight of 'Northblue' blueberry," Journal of the  
American Society for Horticultural Science. 1996, 121:2,

339 342; Galletta, G. J.; Ballington, J. R.; Daubeny, H. A.; Brennan, R. M.; Reisch, B. J.; Pratt, C.; Ferguson, A. R.; Seal, A. G.; McNeilage, M. A.; Fraser, L. G.; Harvey, C. F.; Beatson, R. A.; Hancock, J. F.; Scott, D. H.; Lawrence, F. J.; Janick, J. (ed.); Moore, J. N. "Fruit breeding. Volume II. Vine and small fruits," Department of Horticulture, Purdue University, West Lafayette, Ind. 1996 John Wiley and Sons; New York; USA; Strik, B.; Brun, C.; Ahmedullah, M.; Antonelli, A.; Askham, L.; Barney, D.; Bristow, P.; Fisher, G.; Hart, J.; Havens, D. Draper A. D. and Chandler C. K. "Accelerating highbush blueberry selection evaluation by early propagation," Journal of the American Society for Horticultural Science. 1986 111(2): 301-303; Pritts M. P. and Hancock J. F. (Eds.) "Highbush blueberry production guide," Northeast Regional Agricultural Engineering Service, Ithaca, N.Y. USA 1992.

Clones of Aurora have been tested for two years at Grand Junction, Mich., South Haven, Mich., Lacota, Mich., Corvallis, Oreg., and Lowell, Oreg. In all cases all resulting plants have stably displayed characteristics of the variety. Consistent high yields at Benton Harbor and Grand Junction, Mich. indicate that the buds and wood of Aurora are tolerant to fluctuating late fall and spring temperatures. Aurora also has excellent winter hardiness, as it has routinely been challenged with mid-winter temperatures below -20° C. Probable areas of adaptation and markets include blueberry growers in Michigan and across the USA, Canada, Argentina, Australia, Chile, France, Germany, and New Zealand.

#### BRIEF DESCRIPTION OF THE DRAWINGS

The present invention will become more fully understood from the detailed description and the accompany drawings, wherein:

FIG. 1 is a photographic print in full color showing, in the foreground, an exemplary 4-year-old Aurora blueberry plant. The plants appearing in the background are of a different variety and are not part of the Aurora blueberry plant, nor is the grass appearing in the picture part of the Aurora blueberry plant.

FIG. 2 is a photographic print in full color illustrating exemplary fruit clusters of a 4-year-old Aurora blueberry plant. Some of the fruit is not yet mature and does not show the characteristic mature color.

#### DETAILED BOTANICAL DESCRIPTION

The following is a detailed botanical description of the new and distinct variety of blueberry, its flowers, fruit and foliage, based on observations of specimens grown at Benton Harbor Mich., Grand Junction, Mich., South Haven, Mich., Lacota, Mich., Corvallis, Oreg., and Lowell, Oreg., over a two year period. The taxonomic characteristics chosen in the detailed description are standard in the practice (R E Gough, R J Hindle, and V G Shutak, "Identification of Ten Highbush Blueberry Cultivars using Morphological Characteristics," *HortScience* 11 (5): 512-4, 1976). Color descriptions, except those given in common terms, are presented in Royal Horticultural Society Colour Chart designations. In cases where the color descriptions cited from The Royal Horticultural Society Colour Chart differ from the colors shown in the drawings, the colors cited from The Royal Horticultural Society Colour Chart should be considered accurate. Any deviation from these colors in the drawings is due to failure of the photographic process to exactly

duplicate the colors of nature. In addition, fruit color designations in Table I are applicable only to mature fruit.

Aurora requires pollination for fruit development; it is self-fertile but pollen from any highbush blueberry species will initiate fruit development.

TABLE I

Aurora Characteristics	
Characteristic	Aurora
<u>Bush</u>	
Mature height	1.5 m
Mature width	1.3 m
Diameter/width ratio	1.1
Growth habit	Upright to semi-spreading
Annual renewal canes	5-6
Internode length on spring shoots	2.5 cm
Mature cane color	Greyed-Green (198A)
Mature cane length	1.4 m-1.6 m
Mature cane width	2 cm-4 cm
Fall color - new shoots	Patches of Red-Purple (60A)
<u>Foliage</u>	
Leaf shape	Elliptic - ovate
Apex shape	Acute
Base shape	Cuniate to truncate
Leaf length	2.5 cm-3.5 cm
Leaf width	1.1 cm-1.5 cm
Leaf length/width ratio	2.1
Leaf serration	Entire
Pubescence	None
Color - top	Green (137A)
Color - bottom	Green (138A)
Veination	Palmate
Vein color	Green (138B)
Petiole length	4 mm
Petiole color	Green (138B)
<u>Blossoms</u>	
Shape of corolla	Elongate-urceolata
Calyx	5 lobed
Style length	At edge of corolla
Color of open flower	Mostly white (155D)
Flower # per cluster	8-10
Flower diameter	5 mm-7 mm
<u>Reproductive Organs</u>	
Type	Berries with seeds
Seed size	1 mm x 2 mm
Number of seeds	10-50 per fruit
<u>Mature Fruit</u>	
Length	1.1 cm-1.3 cm
Width	1.4 cm-1.8 cm
Color	(156D)
Shape	Globose, uniform
Color with bloom	Violet Blue (98A)
Color without bloom	Blue (103A)
Pedicel scar size	1.44 mm
Pedicel length	2 cm-3 mm
Pedicel color	Green (143B)
Average weight	1.5 gm

In trials in Michigan and Oregon, Aurora has consistently had better color, was firmer and had a better picking scar than Elliott (Tables II and III). They also had greatly improved flavor. In several post-harvest trials, the relative fruit rot susceptibility of the Elliott and Aurora genotypes was similar and good (Table IV). Aurora had a slightly longer storage life than 'Elliott' in 2001 and 2002. The fruit are eaten fresh, frozen or processed into products like jams, jellies and yogurt.

Aurora is distinguishable from Liberty (co-pending application Ser. No. 10/350,343, filed Jan. 23, 2003) in that Aurora is first harvested 10–14 days after Liberty (mid-August vs. Late August to early September in Michigan). Aurora leaves are entire, while Liberty leaves are serrated. The bush height/width ratio of Aurora is about 1.1, while Liberty's is about 1.4.

Aurora is distinguishable from Draper (co-pending application Ser. No. 10/350,344, filed Jan. 23, 2003) in that Aurora is harvested 6–8 weeks after Draper (mid-August vs. late June to early July in Michigan). The bush height/width ratio of Aurora is about 1.1, while Draper's is about 2.2. Aurora's leaves are elliptic-ovate (leaf/length ratio is >2), while Draper's leaves are narrow elliptic (leaf/length ratio is <2).

TABLE II

Traits	Aurora	Elliott
<u>Horticultural</u>		
Date of first harvest	8/29 and 9/7	8/23 and 8/27
Fruit load	7	8
Size	7	6
Color	8	6
Picking scar	8	7
Firmness	8	7
Flavor	8	5
<u>Taxonomy</u>		
Leaf margin	entire	entire
Leaf shape	elliptic-ovate	elliptic
Leaf length/width ratio	>2	<2
Bush diameter/width ratio	1.1	1.4
Fall color on 1-year shoots	red and green	solid red
Bark texture - mature wood	rough	rough

TABLE III

Mean fruit ratings of 'Elliott' and 'Aurora' at Grand Junction, MI, Corvallis, OR and Lowell, OR in 2002. Plants were set as two-year-old rooted cuttings in 2000. Evaluations were made when the bushes were 50% ripe.

Location	Cultivar	Fruit			Picking		Flavor
		load	Size	Color	scar	Firmness	
Grand Junction	Elliott	8 <sup>Z</sup>	6	6	7	7	5
	Aurora	7	7	8	8	8	8

TABLE III-continued

Mean fruit ratings of 'Elliott' and 'Aurora' at Grand Junction, MI, Corvallis, OR and Lowell, OR in 2002. Plants were set as two-year-old rooted cuttings in 2000. Evaluations were made when the bushes were 50% ripe.

Location	Cultivar	Fruit			Picking		Flavor
		load	Size	Color	scar	Firmness	
Corvallis	Elliott	9	6	7	7	7	6
	Aurora	8	7	8	8	8	8
Lowell	Elliott	9	6	7	7	7	6
	Aurora	8	7	8	8	8	8

<sup>Z</sup>The rating scale 1–9, with 1–4 = inferior, 5–6 = acceptable, 7 = good, 8 = very good and 9 = superior.

TABLE IV

Comparison of postharvest storage life and fungal rots of 'Elliott' and 'Aurora' harvested at Benton Harbor, MI. Fruit were picked on the same day, when the bushes were 30–40% ripe.

Determinations	Elliott	Aurora
<u>Percentage of fruit rotted<sup>X</sup></u>		
2000	12%	—
2001	12%	11%
2002	10%	16%
<u>Types of fruit rots in 2002<sup>Y</sup></u>		
Alternaria	6	10
Colletotrichum	4	2
Botrytis	0	0
<u>Storage life (days)<sup>Z</sup></u>		
2000	7	—
2001	7	14
2002	7	14

<sup>X</sup>Four pints of fruit were evaluated after being held for 3 weeks at 2 C. in zip-lock storage bags. Fungal species were not determined.

<sup>Y</sup>Fifty fruit were randomly selected from 4 pints and held for ten days at room temperature.

<sup>Z</sup>Four pints of fruit were held at 2 C. in plastic zip-lock bags and examined at 7 day intervals. The containers were considered non-salable, if more than a few fruit appeared rotten or >25% were soft to the touch.

What is claimed is:

1. A new and distinct highbush blueberry plant, substantially as illustrated and described herein.

\* \* \* \* \*



*Fig. 1*



*Fig. 2*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : PP15,185 P3  
DATED : September 28, 2004  
INVENTOR(S) : James F. Hancock

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 2,

Line 12, "rood" should be -- root --.

Line 32, "Sargent, K. L.," should be -- Sargent, K. L. --

Column 3,


Line 26, "below-20" should be -- below -20° --.

Column 4,

Line 56, "3 mm" should be -- 3 cm --.

Signed and Sealed this

First Day of February, 2005

A handwritten signature in black ink on a dotted background. The signature reads "Jon W. Dudas" in a cursive style.

JON W. DUDAS

*Director of the United States Patent and Trademark Office*