



US00PP15266P2

(12) **United States Plant Patent**  
**Hoogendoorn**

(10) **Patent No.:** **US PP15,266 P2**  
(45) **Date of Patent:** **Oct. 26, 2004**

(54) **ALSTROEMERIA PLANT NAMED**  
**'ZALSAMBIA'**

(50) Latin Name: *Alstroemeria hybrida*  
Varietal Denomination: **Zalsambia**

(75) Inventor: **Cornelis Arie Hoogendoorn,**  
Aarlanderveen (NL)

(73) Assignee: **Van Zanten Plants B.V.,** Hillegom  
(NL)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/602,083**

(22) Filed: **Jun. 23, 2003**

(51) **Int. Cl.<sup>7</sup>** ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./309**

(58) **Field of Search** ..... **Plt./309**

*Primary Examiner*—Kent Bell

*Assistant Examiner*—S B McCormick-Ewoldt

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Alstroemeria* plant named  
'Zalsambia', characterized by its erect and strong flowering  
stems; vigorous growth habit; white and red purple  
bi-colored flowers; and good postproduction longevity.

**1 Drawing Sheet**

**1**

Botanical classification/cultivar designation: *Alstroemeria hybrida* cultivar *Zalsambia*.

**BACKGROUND OF THE INVENTION**

The present Invention relates to a new and distinct cultivar of *Alstroemeria* plant, botanically known as *Alstroemeria hybrida*, commercially used as a cut flower *Alstroemeria*, and hereinafter referred to by the name 'Zalsambia'.

The new *Alstroemeria* is a product of a planned breeding program conducted by the Inventor in Hillegom, The Netherlands. The objective of the breeding program was to develop new cut flower *Alstroemeria* cultivars with desirable flower and plant qualities, attractive flower colors and excellent postproduction longevity.

The new *Alstroemeria* originated from a cross-pollination made by the Inventor in June, 1996 in Hillegom, The Netherlands, of a proprietary *Alstroemeria hybrida* selection identified as 94097-2, not patented, as the female, or seed, parent with a proprietary *Alstroemeria hybrida* selection identified as 95513-5, not patented, as the male, or pollen, parent. The new *Alstroemeria* was discovered and selected by the Inventor as a flowering plant within the progeny of the stated cross-pollination in a controlled environment in Hillegom, The Netherlands in May, 1997.

Asexual reproduction of the new cultivar by root divisions taken in a controlled environment in Hillegom, The Netherlands, since September, 1997, has shown that the unique features of this new *Alstroemeria* are stable and reproduced true to type in successive generations of asexual propagation.

**SUMMARY OF THE INVENTION**

Plants of the cultivar *Zalsambia* have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Zalsambia'. These characteristics in combination distinguish 'Zalsambia' as a new and distinct cultivar:

**2**

1. Erect and strong flowering stems.
2. Vigorous growth habit.
3. White and red purple bi-colored flowers.
4. Good postproduction longevity.

Plants of the new *Alstroemeria* can be compared to plants of the female parent selection. In side-by-side comparisons conducted in Rijnsburg, The Netherlands, plants of the new *Alstroemeria* differed from plants of the female parent selection in the following characteristics:

1. Plants of the new *Alstroemeria* had shorter flowering stems than plants of the female parent selection.
2. Plants of the new *Alstroemeria* and the female parent selection differed in flower margin coloration.

Plants of the new *Alstroemeria* can be compared to plants of the male parent selection. In side-by-side comparisons conducted in Rijnsburg, The Netherlands, plants of the new *Alstroemeria* differed from plants of the male parent selection in the following characteristics:

1. Plants of the new *Alstroemeria* had longer flowering stems than plants of the male parent selection.
2. Plants of the new *Alstroemeria* and the male parent selection differed in flower margin coloration.

Plants of the new *Alstroemeria* can be compared to plants of the cultivar *Stabec*, disclosed in U.S. Plant Pat. No. 9,041. In side-by-side comparisons conducted in Rijsenhout, The Netherlands, plants of the new *Alstroemeria* differed from plants of the cultivar *Stabec* in the following characteristics:

1. Plants of the new *Alstroemeria* had smaller flowers than plants of the cultivar *Stabec*.
2. Plants of the new *Alstroemeria* had fewer flowers per inflorescence than plants of the cultivar *Stabec*.
3. Plants of the new *Alstroemeria* had white and red purple bi-colored flowers whereas plants of the cultivar *Stabec* had white and red bi-colored flowers.
4. Plants of the new *Alstroemeria* had shorter peduncles than plants of the cultivar *Stabec*.

**BRIEF DESCRIPTION OF THE PHOTOGRAPH**

The accompanying colored photograph illustrates the overall appearance of the new *Alstroemeria*, showing the colors as true as it is reasonably possible to obtain in colored



reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Alstroemeria*.

The photograph comprises a side perspective view of a typical flowering stem of 'Zalsambia'.

#### DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph, following observations and measurements describe plants of the new *Alstroemeria* grown in Rijsenhout, The Netherlands in a glass-covered greenhouse in ground beds. During the production of the plants, day temperatures ranged from 15 to 25° C., night temperatures ranged from 10 to 15° C. and light levels averaged 5,000 lux. Plants used for the photograph and description were about one year old. The photograph and the description were taken during August and September, 2002. Color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Alstroemeria hybrida* cultivar Zalsambia.

Parentage:

*Female parent*.—Proprietary *Alstroemeria hybrida* selection identified as 94097-2, not patented.

*Male parent*.—Proprietary selection of *Alstroemeria hybrida* identified as 95513-5, not patented.

Propagation:

*Type*.—By root divisions.

*Root description*.—Fibrous, fleshy, thick; white, close to 155D, in color.

*Rooting habit*.—Freely branching.

*Rhizomes*.—Shape: Elongate; rounded. Length: About 10 to 30 cm. Diameter: About 3 to 10 mm. Texture: Smooth. Color: Close to 155D.

Plant description:

*Plant habit*.—Upright; freely branching, bushy appearance. Vigorous growth habit.

*Time from planting to harvest of cut flowers*.—About 80 to 90 days.

*Number of flowering stems produced per year*.—About 204 to 220.

*Plant height*.—About 125 to 175 cm.

*Plant diameter (spread)*.—About 25 to 30 cm.

*Flowering stem description*.—Aspect: Erect. Length: About 150 cm. Diameter: About 4.75 to 10 mm. Internode length: About 1 to 6 cm. Strength: Strong. Texture: Glabrous. Color: Close to 146A.

*Foliage description*.—Leaves asymmetrical; sessile. Length: About 15 to 20 cm. Width: About 4.5 to 5 cm. Shape: Lanceolate. Apex: Acute. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Venation pattern: Parallel. Color: Developing and fully developed foliage, upper surface: Close to 137A; moderately glossy. Developing and fully developed foliage, lower surface: Close to 137C. Venation: Upper surface, close to 137A; lower surface, close to 137C.

Flower description:

*Flower type and habit*.—Single cup-shaped flowers arranged in compound umbels. Perianth segments separate. Freely and continuously flowering. Flowers not persistent.

*Natural flowering season*.—Flowering continuous during the spring in The Netherlands.

*Fragrance*.—None detected.

*Flower longevity on the plant*.—About four weeks.

*Flower longevity as a cut flower*.—About 12 to 16 days.

*Flower buds (showing color)*.—Length: About 2.5 to 4.5 cm. Diameter: About 1 to 2 cm. Shape: Roughly ovoid. Color: Greenish.

*Umbel length*.—About 17 to 22 cm.

*Umbel diameter*.—About 15 to 25 cm.

*Number of flowers per umbel*.—About 6 to 10.

*Flower length (height)*.—About 6 to 7 cm.

*Flower diameter*.—About 5 to 6 cm.

*Flower depth*.—About 5.5 to 6.5 cm.

*Perianth*.—Arrangement: Six arranged in two whorls, each whorl with two lateral and one median segments. Size, inner perianth segments: Length: Laterals, about 6.5 to 7 cm; median, 5 to 5.5 cm. Width: Laterals, about 1.5 to 2 cm; median, about 2 cm. Size, outer perianth segments: Length: Laterals, about 5.5 to 6 cm; median, about 6 to 6.5 cm. Width: Laterals, about 3.5 to 4 cm; median, about 3 to 3.5 cm. Shape, inner perianth, all segments: Oblanceolate. Shape, outer perianth, all segments: Obovate. Apex, inner perianth, all segments: Acute. Apex, outer perianth, all segments: Bracket-shaped. Base, inner and outer perianths, all segments: Attenuate. Margin, inner and outer perianths, all segments: Entire; weakly undulate. Texture, inner and outer perianths, all segments: Smooth, glabrous; velvety. Color, inner perianth, lateral tepals: When opening and fully opened, upper surface: 155A; towards the apex, 63A; towards the base, 69C; spots and stripes, close to 187A yellow splotch close to 5C. When opening and fully opened, lower surface: 155A; towards the apex, 63A; towards the base, 69C. Color, inner perianth, median tepal: When opening and fully opened, upper and lower surfaces: 155D; towards the apex, 63A. Color, outer perianth, lateral tepals: When opening and fully opened, upper surface: 69B; towards the apex and margins, 63A; apex tip, close to 146A. When opening and fully opened, lower surface: 69B; towards the apex and margins, 63B; apex tip, close to 146A. Color, outer perianth, median tepal: When opening and fully opened, upper and lower surfaces: 69B; towards the apex and margins, 63A to 63B; apex tip, close to 146A.

*Peduncles*.—Length: About 7 to 10 cm. Diameter: About 3 to 4 cm. Strength: Strong. Angle: About 20 to 30° from vertical. Texture: Smooth, glabrous. Color: Close to 137A.

*Pedicels*.—Length: About 3 to 5 cm. Diameter: About 2 to 3 mm. Strength: Strong. Angle: About 20 to 30° from vertical. Texture: Smooth, glabrous. Color: Close to 137A.

*Reproductive organs*.—Stamens: Quantity per flower: Six. Anther shape: Elliptical. Anther length: About 8 mm. Anther diameter: About 3 mm. Anther color: Close to 197B. Pollen amount: Abundant. Pollen color: Close to 202A to 202B. Pistils: Quantity per flower: One. Style length: About 3.5 to 4 cm. Style color: White to pink. Ovary color: Close to 144A.

*Fruit*.—Shape: Globular. Color: Brownish.

*Disease/pest resistance*: Plants of the new *Alstroemeria* have not been observed to be resistant to pathogens and pests common to *Alstroemerias*.

*Temperature tolerance*: Plants of the new *Alstroemeria* have been observed to tolerate temperatures from -5 to 40° C. It is claimed:

1. A new and distinct cultivar of *Alstroemeria* plant named 'Zalsambia', as illustrated and described.



