



US00PP15393P2

(12) **United States Plant Patent**
Chou et al.(10) **Patent No.:** US PP15,393 P2
(45) **Date of Patent:** Nov. 30, 2004

- (54) **IMPATIENS PLANT NAMED 'BALFUSINRED'**
- (50) Latin Name: *Impatiens walleriana* × *auricoma*
Varietal Denomination: Balfusinred
- (75) Inventors: **Tau San Chou**, Batavia, IL (US);
Kristin Barry, Gainesville, FL (US)
- (73) Assignee: **Ball Horticultural Company**, West Chicago, IL (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **10/741,994**

- (22) Filed: **Dec. 19, 2003**
- (51) Int. Cl.⁷ **A01H 5/00**
- (52) U.S. Cl. **Plt./317**
- (58) Field of Search **Plt./317**

Primary Examiner—Kent Bell

(74) Attorney, Agent, or Firm—Wood, Phillips, Katz, Clark & Mortimer

(57) **ABSTRACT**

A new and distinct cultivar of *Impatiens* plant named 'Balfusinred', characterized by its single, cupped flowers, medium green-colored foliage, upright and mounded habit, and excellent basal branching.

1 Drawing Sheet

1

Latin name of genus and species of plant claimed: *Impatiens walleriana* × *auricoma*.
Variety denomination: 'Balfusinred'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct *Impatiens* plant botanically known as *Impatiens walleriana* × *auricoma* and hereinafter referred to by the cultivar name 'Balfusinred'.

The new cultivar was developed by the inventors in a controlled breeding program during March 1999 at Elburn, Ill. The objective of the breeding program was to develop *Impatiens* cultivars with numerous flowers, new and unique flower shape and colors, excellent basal branching and upright compact to moderate growth habit.

The female (seed) parent of 'Balfusinred' was the proprietary *Impatiens walleriana* selection designated '12660' (not patented) characterized by its upright growth habit, white-colored flowers, and medium green-colored foliage. The male (pollen) parent of 'Balfusinred' was the proprietary *Impatiens auricoma* selection designated '10449-1-2' (not patented) characterized by its upright growth habit, yellow-colored flowers, and dark green-colored foliage. 'Balfusinred' was discovered and selected as a single flowering plant within the progeny of the above stated cross in February of 2000 at Elburn, Ill. and initially designated PAS '1-08'.

Asexual reproduction of the new cultivar by shoot tip or stem cuttings taken since February 2000 at Elburn, Ill. and West Chicago, Ill. has demonstrated that the new cultivar reproduces true to type with all the characteristics, as herein described, firmly fixed and retained through successive generations of such asexual propagation.

SUMMARY OF THE INVENTION

The new cultivar has not been observed under all possible environmental conditions to date. Accordingly, it is possible that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length without, however, any variance in genotype.

It was repeatedly found that the cultivar of the present invention:

2

1. Exhibits single, slightly cupped, flowers,
2. Forms medium green-colored foliage,
3. Exhibits an upright and mounded growth habit, and
4. Exhibits a good basal branching character.

Plants of the new cultivar differ from plants of the female parent primarily in flower color, flower shape and leaf shape and from plants of the male parent primarily in flower color, flower shape, leaf size and branching habit. Plants of the new cultivar can be compared to 'Balfusheat' disclosed in U.S. Plant patent application Ser. No. 10/741,696. 'Balfusheat' and 'Balfusinred' were derived from crossing the same two cultivars, however, they exhibit very different flower colors.

Plants of the new cultivar can be compared to plants of '96-009-4' (U.S. Plant Pat. No. 10,973). In a side-by-side comparison conducted in West Chicago, Ill., plants of the new cultivar differed from plants of '96-009-4' primarily in the following characteristics:

1. Plants of the new cultivar have a more compact growth habit than plants of '96-009-4', and
2. Plants of the new cultivar have larger flowers than plants of '96-009-4'.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it is reasonably possible to make the same in color illustrations of this type, typical flower and foliage characteristics of the new cultivar. Colors in the photographs differ from the color values cited in the detailed description, which accurately describe the colors of 'Balfusinred'. The plants were grown for 13 weeks in a greenhouse at West Chicago, Ill.

FIG. 1 illustrates a side view of the overall growth and flowering habit of 'Balfusinred'.

FIG. 2 illustrates a close-up view of a single flower of 'Balfusinred'.

DETAILED DESCRIPTION

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 1995 edition, except where general color terms of ordinary significance are used. The

color values were determined on Sep. 3, 2003 between 9:00 and 10:45 a.m. under natural light conditions.

The following measurements and comparisons describe plants produced from shoot tip or stem (with two nodes) cuttings taken from stock plants and grown under greenhouse conditions comparable to those used in commercial practice in West Chicago, Ill. The plants were grown in 10 cm pots for 15 weeks utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 72° F. during the day and approximately 65° F. during the night. Greenhouse light levels were maintained at approximately 4,000–6,000 footcandles during the day.

Botanical classification: *Impatiens wallerianaxauricoma* cultivar ‘Balfusinred’.

Parentage:

Female (seed) parent.—Proprietary *Impatiens walleriana* selection designated ‘12660’ (not patented).

Male (pollen) parent.—Proprietary *Impatiens auricoma* selection designated ‘10449-1-2’ (not patented).

Propagation:

Type cutting.—Shoot tip or stem (with two nodes).

Time to initiate roots.—Approximately 7–14 days.

Time to develop a rooted cutting.—Approximately 21 days.

Root description.—Fine, fibrous.

Rooting habit.—Freely branching.

Plant description:

Habit of growth.—Moderately vigorous with good basal branching. A mature plant, 15 weeks after the planting of a rooted cutting, commonly measures approximately 23.7 cm in height and approximately 38.5 cm in width.

Plant form.—Upright and mounded.

Lateral branches.—Quantity: Approximately 4. Length: Approximately 17.1 cm. Diameter: Approximately 8 mm. Texture: Glabrous. Color: 145A with faint overlay of 45A. Internode length: Approximately 3.3 cm.

Foliage.—Type: Simple. Arrangement of lower leaves: Alternate. Arrangement of upper leaves: Whorls. Shape: Elliptic. Apex: Acuminate/cuspidate. Base: Attenuate. Margin: Crenate with ciliation. Texture of upper and lower surface: Glabrous. Venation pattern: Pinnate, arcuate. Size of mature foliage: Length: Approximately 7.2 cm. Width: Approximately 3.9 cm. Color of mature foliage: Upper surface is 137A with veins and mid-vein of 145A. Lower surface is 138B with veins and mid-vein of 145A. Petiole: Diameter: Approximately 2 mm. Length: Approximately 2.7 cm with 2 mm stipules along edges. Texture: Upper and lower surface: Glabrous. Color: Upper and lower surface: 145A with overlay of 46A.

Flower description:

Flower type.—Single, not fragrant with self-cleaning petals.

Flowering habit.—Freely flowering with flowers positioned above the foliage and facing outward.

Natural flowering season.—Year round in greenhouse environment. Flowering is continuous from spring until fall in the garden.

Flower longevity on plant.—Approximately 5 days.

Quantity of flowers.—Approximately 8 flowers and 7 buds per stem at any one time.

Mature flower buds (just before opening).—Shape: Ovate. Length: Approximately 1.2 cm. Diameter: Approximately 9 mm. Texture: Glabrous. Color: 38B.

Flower shape.—Oval, cupped when first open becoming more flat with age.

Flower size.—Length: Approximately 3.1 cm. Width: Approximately 2.5 cm. Depth: Approximately 1.3 cm.

Petals.—Number: Five per flower. Texture: Glabrous. Appearance: Iridescent.

Superior petal.—Length: Approximately 1.2 cm. Width: Approximately 1.2 cm. Shape: Cucullate. Margin: Entire. Apex: Emarginate. Base: Truncate. Color of upper surface: 55B with overlay of 37A. Color of lower surface: 39C.

Lateral petals.—Length: 1.3 cm. Width: 1.1 cm. Shape: Obovate. Margin: Entire. Apex: Obtuse. Base: Attenuate. Color of upper surface: Margin at lower edge is 55A, rest of margin is 37A. There is central spot of 33C and base of N155B. Color of lower surface: 29B.

Lower petals.—Length: 1.8 cm. Width: 1.1 cm. Shape: Obovate. Margin: Entire. Apex: Obtuse. Base: Attenuate. Color of upper surface: 55A with center of petal having a small area of N30B. Color of lower surface: Background of 6D with marginal overlay of 55A and central spot of 33B.

Peduncles.—Strength: Strong. Angle to stem: Acute. Length: Approximately 2.1 cm. Diameter: Approximately 1 mm. Texture: Glabrous. Color: 144C.

Calyx.—Consists of 3 sepals. Lateral sepals: Length: Approximately 9 mm. Width: Approximately 6 mm. Texture: Glabrous. Color of both surfaces: 154D with apex of 144B. Lower sepal: Length: Approximately 1.3 cm. Width: Approximately 1 cm. Texture: Glabrous. Color of both surfaces: 16C with apex of 1447B. Base of lower sepal is modified to form spur.

Spur.—Quantity: One per flower. Double tipped. Length: Approximately 1.3 cm. Diameter at flower: Approximately 2 mm. Diameter of tips: Approximately 1 mm. Aspect: Moderately curved downward. Color: 150D.

Reproductive organs.—Androecium: Five stamens, anthers are fused together forming one organ that surrounds the pistil. Filament color: 8B. Anther shape: Hooded. Anther length: Approximately 2 mm. Anther color: N78D. Pollen: None observed. Gynoecium: Pistil number: One per flower. Pistil length: 6 mm. Stigma shape: Five pointed star. Stigma length: 0.5 mm. Stigma color: Colorless, translucent. Style length: 1.5. Style color: Colorless, translucent. Ovary length: 4 mm. Ovary diameter: 2 mm. Ovary texture: Glabrous. Ovary color: N144C.

Seed and fruit development: Neither seed nor fruit production has been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Impatiens* has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Impatiens* plant named ‘Balfusinred’ substantially, as herein shown and described, which:

1. Exhibits single, cupped, flowers,
2. Forms medium green-colored foliage,
3. Exhibits a good basal branching character, and
4. Exhibits an upright and mounded growth habit.

* * * * *



Figure 1




Figure 2