



US00PP15434P2

(12) **United States Plant Patent**  
**Westhoff**

(10) **Patent No.:** **US PP15,434 P2**  
(45) **Date of Patent:** **Dec. 14, 2004**

(54) **LOBELIA PLANT NAMED 'WESLOLAV'**

PP13,264 P2 \* 11/2002 Heims ..... Plt./263  
PP14,419 P2 \* 12/2003 Heims ..... Plt./263

(50) Latin Name: *Lobelia erinus*  
Varietal Denomination: **Weslolav**

**OTHER PUBLICATIONS**

(75) Inventor: **Heinrich Westhoff**, Südlohn (DE)

UPOV ROM GTITM Computer Database, GTI Jouve  
Retrieval Software 2004/02 Citation for 'Weslolav'.\*

(73) Assignee: **Josef & Heinrich Westhoff**  
**Gartenbau - Spezialkulturen**, Südlohn  
(DE)

\* cited by examiner

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

*Primary Examiner*—Bruce R. Campell  
*Assistant Examiner*—W C Haas  
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(21) Appl. No.: **10/805,580**

(57) **ABSTRACT**

(22) Filed: **Mar. 20, 2004**

(51) **Int. Cl.**<sup>7</sup> ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./263**

(58) **Field of Search** ..... **Plt./263**

A new and distinct cultivar of *Lobelia* plant named  
'Weslolav', characterized by its cascading plant habit; freely  
branching habit with short internodes; dense and bushy plant  
form; continuously and freely flowering habit; densely  
pubescent leaves; light lavender-colored flowers; and rela-  
tively tolerant to high temperatures.

(56) **References Cited**

**1 Drawing Sheet**

**U.S. PATENT DOCUMENTS**

PP13,253 P2 \* 11/2002 Cosner ..... Plt./263

**1**

**2**

Botanical classification/cultivar designation: *Lobelia eri-  
nus* cultivar **Weslolav**.

'Weslolav'. These characteristics in combination distinguish  
'Weslolav' as a new and distinct cultivar:

**BACKGROUND OF THE INVENTION**

The present Invention relates to a new and distinct culti-  
var of *Lobelia* plant, botanically known as *Lobelia erinus*,  
and hereinafter referred to by the name 'Weslolav'.

5

1. Cascading plant habit.
2. Freely branching habit with short internodes; bushy and  
dense plant form.
3. Continuously and freely flowering habit.
4. Densely pubescent stems and leaves.
5. Light lavender-colored flowers.
6. Relatively tolerant to high temperatures.

The new *Lobelia* is a product of a planned breeding  
program conducted by the Inventor in Südlohn, Germany.  
The objective of the breeding program was to develop new  
*Lobelia* cultivars with continuous flowering, interesting  
flower colors and high temperature tolerance.

10

Plants of the new *Lobelia* have larger flowers than the  
female parent, a proprietary selection identified as code  
number 01P050. In addition, plants of the new *Lobelia* and  
the female parent selection differ in flower color. Plants of  
the new *Lobelia* have foliage with dense pubescence com-  
pared to the smooth foliage of plants of the male parent, the  
cultivar **Weslobigblue**. In addition, plants of the new *Lobelia*  
have thicker stems than plants of the cultivar **Weslobigblue**  
and differ in flower color.

The new *Lobelia* originated from a cross-pollination made  
by the Inventor in 2002 of a proprietary selection of *Lobelia  
erinus* identified as code number 01P050, not patented, as  
the female, or seed, parent with the *Lobelia erinus* cultivar  
**Weslobigblue**, disclosed in U.S. Plant Pat. No. 12,634, as the  
male, or pollen, parent. The new *Lobelia* was discovered and  
selected by the Inventor from within the resultant progeny  
from the above-mentioned cross-pollination in a controlled  
environment in Südlohn, Germany in 2002.

15

Plants of the new *Lobelia* differ primarily from plants of  
the cultivar **Weslowhite**, disclosed in U.S. Plant patent  
application Ser. No. 10/805,583 primarily in flower color.

Asexual reproduction since 2002 of the new cultivar by  
terminal cuttings in a controlled environment in Südlohn,  
Germany, has shown that the unique features of this new  
*Lobelia* are stable and reproduced true to type in successive  
generations.

20

Plants of the cultivar **Weslolav** can be compared to the  
cultivar **Weslowei**, disclosed in U.S. Plant Pat. No. 12,708.

**SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of

25

However, in side-by-side comparisons conducted in  
Südlohn, Germany, plants of the cultivar **Weslolav** and the  
cultivar **Weslowei** differed in the following characteristics:

30

1. Plants of the new *Lobelia* had thinner stems than plants  
of the cultivar **Weslowei**.
2. Plants of the new *Lobelia* had foliage and stems were  
more pubescent than plants of the cultivar **Weslowei**.
3. Plants of the new *Lobelia* flowered more continuously  
than plants of the cultivar **Weslowei**.



4. Plants of the new *Lobelia* and the cultivar Weslowei differed in flower coloration.
5. Plants of the new *Lobelia* were more tolerant to high temperatures than plants of the cultivar Weslowei.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description, which accurately describe the actual colors of the new *Lobelia*.

The photograph at the top of the sheet comprises a close-up view of typical flowers of 'Weslolav'.

The photograph at the bottom of the sheet comprises side view of a typical plant of 'Weslolav' grown in a container.

#### DETAILED BOTANICAL DESCRIPTION

Plants of the cultivar Weslolav have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype.

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used. Plants used for the description were grown in a glass-covered greenhouse and conditions that closely approximate commercial production conditions during the spring and summer in Südlohn, Germany. Plants used for the above-mentioned photographs and following description were grown as one plant per 12-cm container or three plants per 25-cm hanging basket container. During the production of the plants, day temperatures ranged from 20 to 25° C. and night temperatures ranged from 16 to 18° C. Plants were pinched once during the production period by removing about 1 to 2 cm of the uppermost apical growing tip. Plants were about 20 weeks from planting when the photographs and description were taken.

Botanical classification: *Lobelia erinus* cultivar Weslolav.

Parentage:

*Female parent*.—Proprietary selection of *Lobelia erinus* identified as code number 01P050, not patented.

*Male parent*.—*Lobelia erinus* cultivar Weslobigblue, disclosed in U.S. Plant Pat. No. 12,634.

Propagation:

*Type cutting*.—Terminal vegetative cuttings.

*Time to initiate roots*.—About 18 to 21 days at 20° C.

*Time to develop roots*.—About 20 to 28 days at 20° C.

*Root description*.—Fine, fibrous and well-branched.

Plant description:

*Plant form/habit*.—Cascading flowering plants with lavender-colored flowers. Lateral shoots outwardly spreading; plants uniform with dense foliage. Foliage and stems are very pubescent. Freely branching with lateral branches forming at every node; dense and bushy plant habit. Pinching plants will enhance branching. Vigorous growth habit.

*Usage*.—Appropriate for hanging baskets, window boxes and patio containers.

*Plant height (soil level to top of plant plane)*.—About 18.2 cm.

*Plant length (soil level to lateral branches apices)*.—About 65 cm.

*Plant diameter*.—About 40 to 50 cm.

*Lateral branch description*.—Length: About 15 cm. Diameter: About 1.2 mm. Internode length: About 1.4 cm. Texture: Densely pubescent. Color: 147B.

*Foliage description*.—Arrangement: Alternate; simple. Basal leaves: Length: About 4.4 cm. Width: About 3 cm. Shape: Nearly round. Apex: Obtuse. Base: Attenuate. Margin: Nearly entire. Petiole length: About 1.5 mm. Mid-plant leaves: Length: About 3.4 cm. Width: About 2.5 cm. Shape: Elliptic to round. Apex: Rounded. Base: Attenuate. Margin: Slightly crenate. Petiole length: About 1 cm. Apical leaves: Length: About 2 cm. Width: About 5 mm. Shape: Linear. Apex: Acute. Base: Attenuate. Margin: Entire. Petiole length: Petioles not observed. Texture, all leaves, upper and lower surfaces: Densely pubescent. Color, all leaves: Developing foliage, upper surface: 146A to 146B. Developing foliage, lower surface: 146B. Fully developed foliage, upper surface: 147A; venation, 147A. Fully developed foliage, lower surface: 147C; venation, 147C.

Flower description:

*Flower type and habit*.—Flowers arranged singly at lateral apices. Flowers held mostly outwardly. Flowers persistent. Older flowers are overgrown by new flowers and foliage. Freely and continuously flowering. Flowers not fragrant.

*Flower shape*.—Tubular with three larger lower petals and two upright petals.

*Natural flowering season*.—Spring until frost in the autumn.

*Flower longevity on the plant*.—Longevity of individual flowers is highly dependent on weather conditions; typically three to ten days.

*Flower size*.—Diameter: About 2.1 cm. Depth (height): About 2.1 cm. Tube length: About 8 mm. Throat diameter, distal end: About 4.3 mm. Tube diameter, proximal end: About 2.8 mm.

*Flower buds*.—Length: About 1.4 cm. Diameter: About 2.5 mm. Shape: Oblong. Color: Base and center, 145D; towards the apex, close to 145C.

*Petals*.—Arrangement: Single whorl of five petals, fused; three larger lower petals and two smaller upper petals. Three lower petals: Shape: Obovate. Length, above throat: About 1.1 cm. Width: About 7 mm. Two upper petals: Shape: Oblanceolate. Length, above throat: About 6.3 mm. Width: About 2 mm. Upper and lower petals: Apex: Cuspidate. Margin: Entire. Texture, upper and lower surfaces: Smooth, satiny. Color: When opening, upper surface: 85B to 85C. When opening, lower surface: 85C to 85D. Fully opened, upper surface: 85C to 85D. Fully opened, lower surface: 85D. Throat: 76C to 76D; stripes, 144B, and spots, 82A. Tube: 76A to 76C; stripes, 144B.

*Sepals*.—Arrangement: Single whorl of five sepals, star-shaped calyx. Length: About 7 mm. Width: About 1.1 mm. Shape: Elliptic. Apex: Acute. Margin: Entire. Texture, upper and lower surfaces: Densely pubescent. Color, upper and lower surfaces: 147B.

*Peduncles*.—Strength: Flexible. Length: About 2.6 cm. Diameter: About 6.5 mm. Texture: Densely pubescent. Color: 137A.

## 5

*Reproductive organs.*—Stamens: Quantity per flower: About five, fused. Anther length: About 2.5 mm. Anther diameter: About 1.7 mm. Anther texture: Pubescent. Anther color: 92A to 84D. Pollen amount: Moderate. Pollen color: 7A. Pistils: Quantity per flower: One. Pistil length: About 1 cm. Stigma shape: Two-parted, ovate. Stigma texture: Pubescent. Stigma color, immature: 79A. Stigma color, mature: 79D. Style length: About 5 mm. Style color: 144A. Ovary color: 137B. Seed/fruit: Seed and fruit production have not been observed.

## 6

Disease/pest resistance: Plants of the new *Lobelia* have not been noted to be resistant to pathogens and pests common to *Lobelia*.

Temperature tolerance: Plants of the new *Lobelia* have been observed to tolerate temperatures ranging from 2 to 30° C.

It is claimed:

1. A new and distinct cultivar of *Lobelia* plant named 'Weslolv', as illustrated and described.

\* \* \* \* \*



