



US00PP15552P2

(12) **United States Plant Patent**
Ballington

(10) **Patent No.: US PP15,552 P2**
(45) **Date of Patent: Feb. 8, 2005**

(54) **STRAWBERRY PLANT NAMED 'BISH'**

(50) Latin Name: *Fragaria×ananassa*
Varietal Denomination: **Bish**

(75) Inventor: **James R. Ballington**, Raleigh, NC
(US)

(73) Assignee: **North Carolina State University**,
Raleigh, NC (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/685,163**

(22) Filed: **Oct. 14, 2003**

(51) **Int. Cl.**⁷ **A01H 5/00**

(52) **U.S. Cl.** **Plt./208**

(58) **Field of Search** **Plt./208**

(56) **References Cited**
PUBLICATIONS

Declaration of James R. Ballington, Ph.D. under 37 C.F.R.
Section 1.132; Jun. 14, 2004.

Release Announcement for Strawberry 'Bish,' Southeast
Strawberry Expo; Nov. 7–8, 2002; Greenville, North Caro-
lina.

Primary Examiner—Anne Marie Grunberg

Assistant Examiner—Annette H Para

(74) *Attorney, Agent, or Firm*—Myers Bigel Sibley &
Sajovec, P.A.

(57) **ABSTRACT**

A new and distinct annual hill culture adapted variety of
Fragaria×ananassa Duch. plant, designated 'Bish', is char-
acterized by being significantly more resistant to anthra-
cnose fruit rot than the current standard annual hill varieties
'Chandler' and 'Camarosa', while being equal to one or both
these varieties for most other economically important fruit
and plant characters. 'Bish' appears best adapted from the
Southern Appalachians up through the Middle Atlantic
regions of the United States.

2 Drawing Sheets

1

Latin name of the genus and species: The Latin name of
the novel variety disclosed herein is *Fragaria×ananassa*
Duch.

Varietal Denomination: The inventive variety of straw-
berry disclosed herein has been given the variety denomi-
nation 'Bish'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct variety
of Junebearing or short-day strawberry (*Fragaria×ananassa*
Duch.), which has been named 'Bish'. This variety is the
result of a cross between 'FL 87-210' (unpatented) and
'Delmarvel' (unpatented) made in 1993 in Raleigh, N.C. as
part of a strawberry breeding program. Plants were first
germinated in a greenhouse in Raleigh, N.C. in 1993. In the
spring of 1994, germinated seedlings were transferred into
seedling trays and allowed to grow over the summer.
Subsequently, the seedlings were transplanted into the field
in Reidsville, N.C. in the fall of 1994. 'Bish' was discovered
in May 1995 in a cultivated field in Reidsville, N.C. and
originally designated 'NCR 95-08' during the testing period.

'Bish' was first asexually reproduced by runners (i.e.,
stolons) and planted in Fletcher, N.C. in fall 1995.
Subsequently, 'Bish' has also been asexually propagated by
tissue culture micropropagation from runner meristems. The
combination of traits disclosed herein that characterize
'Bish' have been retained true to type through successive
cycles of asexual propagation.

SUMMARY OF THE INVENTION

'Bish' is a new and distinct variety of strawberry plant of
the short-day type that is adapted to annual hill culture. It is
significantly more resistant to anthracnose fruit rot than the
current standard annual hill varieties 'Chandler' (U.S. Plant
Pat. No. 5,262) and 'Camarosa' (U.S. Plant Pat. No. 8,708),

2

while being equivalent to both of these varieties for most
other economically important fruit and plant characteristics.
Fruit skin color of 'Bish' is dark red and flesh color light red.
The calyx is medium in size and reflexed. Petiole pubes-
cence on 'Bish' is more dense than on 'Chandler', and
basipetal in direction, compared with perpendicular to acro-
petal in the latter variety. 'Bish' is also characterized by the
presence of two prominent leafy petiole bracts on mid-tier
leaves.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows fruit shape and color of 'Bish' strawberry.

FIG. 2 shows internal flesh color of 'Bish' strawberry.

FIG. 3 shows a typical leaf of 'Bish' strawberry.

**DETAILED BOTANICAL DESCRIPTION OF
THE VARIETY**

The following is a detailed botanical description of a new
and distinct variety of *Fragaria×ananassa* Duch. known as
'Bish'. The observations described below are from mature
plants grown in test plots established in an annual hill
strawberry production system on raised beds covered with
black plastic mulch and subsurface drip irrigation.

Asexual propagules derived from the original source were
established in an observation plot at Fletcher, N.C., in fall
1995, and in replicated trials at Reidsville and Fletcher,
N.C., from 1996–2001; at Castle Hayne, N.C., in 1997 and
1999; and at Beltsville, Md., from 1998–2000. Plants were
established in double offset rows spaced 12 inches apart on
the beds with plants also spaced 12" apart within rows. The
North Carolina test plots were established in a split plot
design and the Beltsville plots in a randomized complete
block design. All plots were established in late summer/early
fall and data collected the following spring. Data from North

Carolina locations in 1997–2003 and the Beltsville location from 1999–2001 are presented in Tables 1–30 below.

Those skilled in the art will appreciate that certain characteristics of the variety will vary with older or, conversely, with younger plants. ‘Bish’ has not been observed under all possible environmental conditions. Where dimensions, sizes, colors and other characteristics are given, it is to be understood that such characteristics are approximations or averages set forth as accurately as practicable. The phenotype of the variety may differ from the descriptions herein with variations in the environment such as season, temperature, light intensity, day length, cultural conditions, and the like. Color notations are based on The Royal Horticultural Society Colour Chart, The Royal Horticultural Society, London, 1995 edition.

‘Bish’ is a typical short-day strawberry variety with respect to seasonal growth cycle with a production pattern similar to ‘Chandler’, the most widely grown cultivar in annual hill strawberry culture in North Carolina at the present time. The five-plant observation plot of ‘Bish’ established at Fletcher, N.C., in fall 1995, yielded 1.04 kg of fruit per plant in 1996. On this basis, ‘Bish’ was chosen for further observation and testing, and was established in the first replicated trials in fall, 1996.

Technical Description of the Variety:

The performance of ‘Bish’ for yield and fruit size in replicated trials in North Carolina is summarized in Tables 1–14 below. Performance at Beltsville, Md., is summarized in Table 15. Fruit quality characteristics are summarized in Tables 16–26. Technical descriptive data are included in Tables 27–30.

In addition to overall performance on a site, planting date was also a variable, since ideal planting date varies by location and often also varies among varieties. In 2001 and 2002, “original” (OR) source plants of ‘Bish’ were also compared to meristemmed and virus-tested source plants. In the tables and description below, the latter are referred to as: (99) tissue-culture propagated in 1999; (00) tissue-culture (TC) propagated in 2000; and (MD) remeristemmed and virus-tested by the Maryland Department of Agriculture.

Yield and Fruit Size (weight): The performance of ‘Bish’ for yield and fruit size in replicated trials in North Carolina is summarized in Tables 1–14 below. Table 1 provides the performance of ‘Bish’ in annual hill culture at the Upper Piedmont Research Station, Reidsville, N.C. in 1997. The yield and size were based on four, five-plant plots each across two planting dates.

TABLE 1

Selection	Planting Date			
	Sep. 17, 1996		Sep. 24, 1996	
	Marketable Yield (g/plant)	Size (g/berry)	Marketable Yield (g/plant)	Size (g/berry)
‘Bish’	136*	17.1A	235*	16.0*
‘Camarosa’	189*	22.2A	161*	20.0*
‘Chandler’	151*	17.1A	180*	18.0*
‘Delmarvel’ ¹	181*	14.7B	91*	17.0*
‘NCS 93-05’ ²	143*	20.5A	177*	20.8*

^{1,2}Unpatented

*Indicates no significant differences observed.

Mean separation was by a Duncan’s Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 2 discloses the performance of ‘Bish’ in annual hill culture at the Mountain Horticultural Crops Research Station, Fletcher, N.C. in 1997. The marketable yield and size were based on four, five-plant plots across two planting dates.

TABLE 2

Selection	Planting Date				% Athracnose ¹
	Aug. 15, 1996		Aug. 22, 1996		
	Market. Yield (g/plant)	Size (g/berry)	Market. Yield (g/plant)	Size (g/berry)	
‘Bish’	551*	15.2B	675A	17.0B	0.0
‘Apollo’ ²	609*	15.3B	585ABC	16.0B	0.5
‘Chandler’	597*	14.9B	652AB	16.3B	5.0
‘Jewel’ ³	536*	12.2B	499ABC	15.3B	3.4
‘NCS 93-05’	521*	24.5A	447BC	26.8A	0.6

¹Percent by weight of fruit infected by strawberry anthracnose caused by *Colletotrichum acutatum*.

²Unpatented

³United States PP5,897

*Indicates no significant differences observed.

Mean separation was by a Duncan’s Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 3 provides the performance of ‘Bish’ in annual hill culture at the Horticultural Crops Research Station, Castle Hayne, N.C. in 1998. The yield and size were based on four, 10-plant replicates planted Oct. 15, 1997.

TABLE 3

Selection	Marketable Yield (g/plant)	Fruit Size (g/plant)	% Athracnose ¹
‘Bish’	210A	17C	4.8A
‘Camarosa’	201A	22B	38.4C
‘Chandler’	35B	18C	42.7C
‘NCS 93-05’	86B	25A	23.6B

¹Percent by weight of fruit infected by strawberry anthracnose caused by *Colletotrichum acutatum*.

Mean separation was by a Duncan’s Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 4 provides the performance of ‘Bish’ in annual hill culture at the Upper Piedmont Research Station, Reidsville, N.C. in 1998. The yield and size were based on four, 10-plant replicates planted Sep. 15, 1997.

TABLE 4

Selection	Marketable Yield (g/plant)	Fruit Size (g/plant)	% Athracnose ¹
‘Bish’	219A	14.7B	2.3A
‘Chandler’	190A	13.9B	22.3C
‘NCS 93-05’	132B	20.1A	11.3B

¹Percent by weight of fruit infected by strawberry anthracnose caused by *Colletotrichum acutatum*.

Mean separation was by a Duncan’s Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 5 provides the performance of 'Bish' in annual hill culture in Fletcher, N.C. in 1998. The yield and size were based on four, 10-plant replicates planted Aug. 15, 1997.

TABLE 5

Selection	Marketable Yield (g/plant)	Fruit Size (g/plant)	% Anthracnose ¹
'Bish'	385A	14.9*	0.04A
'Chandler'	327B	14.2*	34.0C
'NCS 93-05'	274C	18.1*	12.6B

¹Percent by weight of fruit infected by strawberry anthracnose caused by *Colletotrichum acutatum*.

*Indicates no significant differences observed.

Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 6 provides the performance of 'Bish' in annual hill culture at the Upper Piedmont Research Station, Reidsville, N.C. in 1999. The yield and size were based on three, 10-plant replicates for each of the three planting dates.

TABLE 6

Selection	Planting Date					
	Sep. 10, 1998			Sep. 17, 1998		
	Mkt. Yld. ¹	Size ²	% Anth. ³	Mkt. Yld. ¹	Size ²	% Anth. ³
'Bish'	277B	12.7C	0.0A	207C	12.3B	0.0A
'Camarosa'	316A	14.4B	4.0B	400A	17.5A	1.0B
'Chandler'	308AB	12.8C	4.2B	281B	12.2B	1.8B
'NCS 93-05'	168C	17.5A	0.0A	211C	17.9A	0.0A

Selection	Planting Date Sep. 23, 1998		
	Mkt. Yld. ¹	Size ²	% Anth. ³
'Bish'	246B	13.5B	0.3B
'Camarosa'	281A	15.9A	0.0A
'Chandler'	230B	12.6B	0.7C
'NCS 93-05'	149C	17.9A	0.0A

¹Mkt. Yld. is marketable yield expressed as g/plant.

²Size is expressed as g/berry.

³% Anth. is percent by weight of fruit infected by strawberry anthracnose caused by *Colletotrichum acutatum*.

Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 7 provides the performance of 'Bish' in annual hill culture at Fletcher, N.C. in 1999. The yield and size were based on three, 10-plant replicates for each of the three planting dates.

TABLE 7

Selection	Planting Date					
	Aug. 17, 1998			Aug. 24, 1998		
	Mkt. Yld. ¹	Size ²	% Anth. ³	Mkt. Yld. ¹	Size ²	% Anth. ³
'Bish'	580A	15C	0.1A	562A	15C	0.1A
'Camarosa'	445B	18B	30B	378B	18B	34C
'Chandler'	88C	14D	50C	498B	14D	6B
'NCS 93-05'	547AB	21A	2.6A	413C	22A	0.1A

TABLE 7-continued

Selection	Planting Date Aug. 31, 1998		
	Mkt. Yld. ¹	Size ²	% Anth. ³
'Bish'	622A	16B	0.1A
'Camarosa'	—	—	—
'Chandler'	467B	13C	13B
'NCS 93-05'	408C	21A	0.1A

¹Mkt. Yld. is marketable yield expressed as g/plant.

²Size is expressed as g/berry.

³% Anth. is percent by weight of fruit infected by strawberry anthracnose caused by *Colletotrichum acutatum*.

Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 8 provides the performance of 'Bish' in annual hill culture at Castle Hayne, N.C. in 2000. The yield, size, and % anthracnose values were averages of three, 20-plant plots on each of three planting dates.

TABLE 8

Selection	Planting Date					
	Oct. 14, 1999			Oct. 20, 1999		
	Mkt. Yld. ¹	Size ²	% Anth. ³	Mkt. Yld. ¹	Size ²	% Anth. ³
'Bish'	333A	13B	0	200A	19C	1.1B
'Camarosa'	—	—	—	132BC	22BC	41.9D
'Chandler'	174C	18A	0	80D	21BC	51.4E
'Gaviota' ⁴	—	—	—	199A	23AB	2.2B
'Gem Star' ⁵	—	—	—	113C	27A	33.0C
'Sw Charlie' ⁶	216B	16A	0	152B	19C	0.0A

Selection	Planting Date Oct. 27, 1999		
	Mkt. Yld. ¹	Size ²	% Anth. ³
'Bish'	205A	18B	0.0A
'Camarosa'	42D	22AB	50.6E
'Chandler'	177BC	19AB	35.6D
'Gaviota' ⁴	188AB	23A	3.9B
'Gem Star' ⁵	181BC	23A	19.0C
'Sw Charlie' ⁶	163C	20AB	0.0A

¹Mkt. Yld. is marketable yield expressed as g/plant.

²Size is expressed as g/berry.

³% Anth. is percent by weight of fruit infected by strawberry anthracnose caused by *Colletotrichum acutatum*.

⁴United States PP10,461

⁵United States PP12,377

⁶United States PP8,729

Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 9 discloses the performance of 'Bish' in annual hill culture in Reidsville, N.C. in 2000. The yield and size were based on three, 10-plant replicates each on each planting date.

TABLE 9

Selection	Planting Date					
	Sep. 15, 1999		Sep. 22, 1999			
	Market-able Yield (g/plant)	Size (g/berry)	% Anthracnose ¹	Market-able Yield (g/plant)	Size (g/berry)	% Anthracnose ¹
'Bish'	327A	17B	0A	209AB	14C	0A
'Camarosa'	303A	20B	26C	147B	17BC	12C

TABLE 9-continued

Selection	Planting Date					
	Sep. 15, 1999			Sep. 22, 1999		
	Market-able Yield (g/plant)	Size (g/berry)	% An-thrac-nose ¹	Market-able Yield (g/plant)	Size (g/berry)	% An-thrac-nose ¹
'Chandler'	257AB	17B	6B	241A	19B	10C
'Gaviota'	135C	21AB	3B	145B	21AB	0A
'Gem Star'	216B	24A	11BC	232A	24A	5B

¹Percent by weight of fruit infected by strawberry anthracnose caused by *Colletotrichum acutatum*. Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 10 provides the performance of 'Bish' in annual hill culture at Fletcher, N.C. in 2000. The yield, size, and % anthracnose values were averages of three, 20-plant plots on each of the three planting dates.

TABLE 10

Selection	Planting Date					
	Aug. 4, 1999			Aug. 11, 1999		
	Mkt. Yld. ¹	Size ²	% Anth. ³	Mkt. Yld. ¹	Size ²	% Anth. ³
'Bish'	701B	16*	0.1A	639B	15B	0.1A
'Chandler'	781A	16*	0.5B	758A	16A	0.2B
'Gaviota'	—	—	—	—	—	—
'Gem Star'	—	—	—	—	—	—

Selection	Planting Date Aug. 18, 1999		
	Mkt. Yld. ¹	Size ²	% Anth. ³
	'Bish'	585B	16BC
'Chandler'	705A	15C	0.8B
'Gaviota'	308D	17B	0.2A
'Gem Star'	543C	22A	2.0C

¹Mkt. Yld. is marketable yield expressed as g/plant.

²Size is expressed as g/berry.

³% Anth. is percent by weight of fruit infected by strawberry anthracnose caused by *Colletotrichum acutatum*.

*Indicates no significant differences observed.

Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 11 provides the performance of 'Bish' in annual hill culture at the Upper Piedmont Research Station, Reidsville, N.C. in 2001. The yield was reduced due to a cool fall resulting in reduced daughter crown development. The yield and size were based on three replicates of 10 plants each on each planting date.

TABLE 11

Selection	Planting Date			
	Sep. 22, 2000		Sep. 29, 2000	
	Marketable Yield (g/plant)	Size (g/berry)	Marketable Yield (g/plant)	Size (g/berry)
'Bish'(Original)	154B	20*	93B	16B
'Bish'(99-TC)	201A	17*	96B	18A

TABLE 11-continued

Selection	Planting Date			
	Sep. 22, 2000		Sep. 29, 2000	
	Marketable Yield (g/plant)	Size (g/berry)	Marketable Yield (g/plant)	Size (g/berry)
'Bish'(00-TC)	125B	18*	—	—
'Camarosa'	231A	21*	118AB	20A
'Chandler'	208A	18*	146A	19A

*Indicates no significant differences observed.

Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 12 discloses the performance of 'Bish' in annual hill culture in Fletcher, N.C. in 2001. The yield was reduced due to a cool fall resulting in reduced daughter crown development. The yield and size were based on three, 10-plant replicates for each planting date.

TABLE 12

Selection	Planting Date			
	Aug. 23, 2000		Aug. 30, 2000	
	Marketable Yield (g/plant)	Size (g/berry)	Marketable Yield (g/plant)	Size (g/berry)
'Bish'(Original)	330C	16C	420B	16B
'Bish'(99-TC)	420B	16C	392B	18AB
'Bish'(00-TC)	391B	16C	357BC	15BC
'Camarosa'	537A	18AB	408B	19AB
'Chandler'	540A	18AB	542A	17B
'NCS 93-05'	255D	20A	231D	22A

Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$.

Table 13 provides the performance of 'Bish' in annual hill culture at the Upper Piedmont Research Station, Reidsville, N.C. in 2002. The yield and size were averaged across three, 10-plant replicates for each of two planting dates (Sep. 5, 2001 and Sep. 12, 2001).

TABLE 13

Selection	Marketable Yield (g/plant)	Fruit Size (g/plant)	% Anthracnose ¹
'Bish'(Original)	231*	14.2AB	0.0A
'Bish'(99-TC)	260*	12.8BC	0.0A
'Bish'(00-TC)	257*	13.2BC	0.0A
'Camarosa'	446*	15.1A	3.5C
'Chandler'	440*	11.4D	0.3B

¹Percent by weight of fruit infected by strawberry anthracnose caused by *Colletotrichum acutatum*.

*Indicates no significant differences observed.

Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Table 14 discloses the performance of 'Bish' in annual hill culture in Fletcher, N.C. in 2002. The yield, size and % anthracnose were based on three, 10-plant replicates each on each planting date.

TABLE 14

Selection	Planting Date					
	Aug. 15, 2001			Aug. 22, 2001		
	Market- able Yield (g/plant)	Size (g/ berry)	% An- thrac- nose ¹	Market- able Yield (g/plant)	Size (g/berry)	% An- thrac- nose ¹
'Bish'(OR)	455B	12D	0.0A	390C	13B	0.0A
'Bish'(MD)	382C	13C	0.0A	488B	12BC	0.0A
'Camarosa'	467B	14C	0.1B	520AB	14B	0.0A
'Chandler'	512A	15B	0.0A	548A	13B	0.1B
'Gem Star'	461B	17A	2.2C	559A	18A	0.4C

¹Percent by weight of fruit infected by strawberry antracnose caused by *Colletotrichum acutatum*. Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

The performance of 'Bish' in annual hill culture at Beltsville, Md. between 1999 and 2001 is summarized in Table 15. 'Bish' was an outstanding performer in replicated trials at Beltsville, Md. The yield and size were based on results from replicated trials involving four, five-plant plots on raised beds with plastic mulch and drip irrigation, but without preplant fumigation.

TABLE 15

Selection	1999		2000		2001 ¹	
	Yield/ plant (g/ plant)	Size (g/ berry)	Yield/ plant (g/ plant)	Size (g/ berry)	Yield/ plant (g/ plant)	Size (g/ berry)
'Bish'	750A	13.7BC	620*	13.4*	530AB	14.2B
'Allstar' ²	880A	13.5BC	700*	13.9*	680A	14.9B
'Chandler'	720AB	11.7C	550*	12.8*	480AB	15.5B
'North- easter' ³	790A	13.1C	580*	15.0*	450ABC	17.2AB
'NCS 93-05'	520B	16.3A	—	—	250C	20.1A

¹Yields were reduced due to a cool fall, resulting in reduced daughter crown development.

^{2,3}Unpatented

*Indicates no significant differences observed.

Mean separation was by a Duncan's Multiple Range Test with $p \leq 0.05$ and values designated with different letters are statistically different.

Overall, 'Bish' was generally equivalent to 'Chandler' for yield and fruit size in North Carolina plots. This same pattern was observed when compared with 'Camarosa' for yield. However, 'Camarosa' typically produced larger fruit than 'Bish'. Yields across locations in North Carolina were generally higher at Fletcher; the plant also appeared to be well-adapted at Beltsville, Md. Therefore, the preferred region of adaptation for 'Bish' appears to be the Southern Appalachians up through the Middle Atlantic States. A consistent pattern for yield differences by planting date was not observed. Thus, it is contemplated that planting dates suitable for 'Chandler' will also be suitable for 'Bish'. Further, there was no consistent differences among plant sources, i.e., original vs. 99 vs. 00 vs. MD (see Tables 11–14). Accordingly, 'Bish' is genetically stable when propagated by micropropagation.

Fruit Characteristics: The performance of 'Bish' for fruit appearance (attractiveness), firmness, symmetry (shape and uniformity), skin toughness (resistance to abrasion) and flavor in subjective evaluations is summarized in Tables 16–26. The values provided in Tables 16–26 were based on a subjective 0–90 scale where less than 60 was unacceptable,

60–69 was acceptable, 70–79 was good, and 80 and above was superior.

Table 16 discloses fruit characteristics of 'Bish' grown at the Horticultural Crops Research Station, Castle Hayne, N.C. in 1998. The averages were of ratings across four, 10-plant plots with separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 16

Selection	Appear- ance	Firm- ness	Sym- metry	Skin Color	Flesh Color	Skin Toughness	Flavor
'Bish'	77AB	77BC	76*	77*	73B	77B	76B
'Camarosa'	77AB	81A	75*	78*	76A	81A	71C
'Chandler'	75B	76C	75*	77*	77A	77B	76B
'NCS 93-05'	79A	79AB	76*	79*	79A	78AB	79A

*Indicates no significant differences observed.

Values designated with different letters are statistically different.

Table 17 shows fruit characteristics of 'Bish' grown at the Upper Piedmont Research Station, Reidsville, N.C. in 1998. The averages were of ratings across four, 10-plant plots with separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 17

Selection	Appear- ance	Firm- ness	Sym- metry	Skin Color	Flesh Color	Skin Toughness	Flavor
'Bish'	74B	74*	74B	74B	73*	72.5B	75.0B
'Chandler'	75B	75*	75AB	75A	75*	75.0A	75.0B
'NCS 93-05'	78A	74*	77A	78A	75*	72.5B	77.5A

*Indicates no significant differences observed.

Values designated with different letters are statistically different.

Table 18 discloses fruit characteristics of 'Bish' grown at the Mountain Horticultural Crops Research Station, Fletcher, N.C. in 1998. The averages were of ratings across four, 10-plant plots with separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 18

Selection	Appear- ance	Firm- ness	Sym- metry	Skin Color	Flesh Color	Skin Toughness	Flavor
'Bish'	78*	79*	78*	79*	71B	76*	76*
'Chandler'	75*	76*	74*	76*	76A	76*	72*
'NCS 93-05'	75*	81*	75*	79*	76A	80*	75*

*Indicates no significant differences observed.

Values designated with different letters are statistically different.

Table 19 discloses fruit characteristics of 'Bish' grown at the Upper Piedmont Research station, Reidsville, N.C. in 1999. Values were of averages of three replicates for a September 17 planting date with mean separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 19

Selection	Appear- ance	Firm- ness	Sym- metry	Skin Color	Flesh Color	Skin Tough- ness	Flavor
'Bish'	76.3AB	80.8B	75.0AB	79.3*	72.5C	80.0B	77.1A
'Cama- rosa'	75.2B	85.5A	73.3B	78.8*	78.2A	84.3A	68.7B
'Chand- ler'	71.7C	77.5C	70.0C	80.0*	75.0BC	76.7C	76.3A

TABLE 19-continued

Selection	Appearance	Firmness	Symmetry	Skin Color	Flesh Color	Skin Toughness	Flavor
'NCS 93-05'	78.3A	80.8B	76.5A	79.2*	75.3B	80.5B	75.3A

*Indicates no significant differences observed.
Values designated with different letters are statistically different.

Table 20 discloses fruit characteristics of 'Bish' grown at Fletcher, N.C. in 1999. Values were of averages of three replicates across each of three planting dates (Aug. 17, 1998; Aug. 24, 1998; and Aug. 30, 1998) with mean separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 20

Selection	Appearance	Firmness	Symmetry	Skin Color	Flesh Color	Skin Toughness	Flavor
'Bish'	77B	80B	77A	79*	72B	80B	79A
'Camarosa'	74C	84A	73B	79*	74A	83A	69C
'Chandler'	71D	77C	71C	79*	75A	77C	75B
'NCS 93-05'	80A	80B	79A	80*	74A	80B	78A

*Indicates no significant differences observed.
Values designated with different letters are statistically different.

Table 21 provides fruit characteristics of 'Bish' grown at Castle Hayne, N.C. in 2000. Values were of averages of four, 20-plant plots with mean separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 21

Selection	Appearance	Firmness	Symmetry	Skin Color	Flesh Color	Skin Toughness	Flavor
'Bish'	75B	76B	76A	78A	74B	74CD	75A
'Camarosa'	74B	80A	74B	76AB	76A	80A	72BC
'Chandler'	69C	75B	72C	75BC	76A	75C	74AB
'Gaviota'	76A	76B	75A	78A	76A	72E	75A
'Sw. Charlie'	76A	73C	74B	74C	70C	73DE	71C
'NCC 87-45'	70C	75B	74B	69D	69C	76B	71C

¹Unpatented
Values designated with different letters are statistically different.

Table 22 provides fruit characteristics of 'Bish' grown at Reidsville, N.C. in 2000. Values were of averages of three replicates of a Sep. 5, 1999 planting date with mean separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 22

Selection	Appearance	Firmness	Symmetry	Skin Color	Flesh Color	Skin Toughness	Flavor
'Bish'	76AB	76B	76A	79A	77AB	74B	78A
'Camarosa'	72C	80A	73B	78A	78A	80A	75B
'Chandler'	72C	75B	73B	76B	75B	75B	75B
'Gaviota'	78A	75B	75AB	80A	76AB	75B	75B
'Gem Star'	75BC	75B	75AB	75B	70C	75B	75B

Values designated with different letters are statistically different.

Table 23 provides fruit characteristics of 'Bish' grown at Fletcher, N.C. in 2000. Values were of averages of three, 20-plant replicates of an Aug. 18, 1999 planting date with mean separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 23

Selection	Appearance	Firmness	Symmetry	Skin Color	Flesh Color	Skin Toughness	Flavor
'Bish'	76B	79A	75B	80A	72*	78A	73AB
'Chandler'	70D	70B	70C	75B	75*	70B	73AB
'Gaviota'	80A	80A	80A	80A	73*	78A	75A
'Gem Star'	73C	78A	75B	72C	70*	78A	70B

*Indicates no significant differences observed.
Values designated with different letters are statistically different.

Table 24 provides fruit characteristics of 'Bish' grown at Fletcher, N.C. in 2001. Values were of averages of three replicates of two planting dates (Aug. 14, 2000 and Aug. 22, 2000) with mean separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 24

Selection	Appearance	Firmness	Symmetry	Skin Color	Flesh Color	Skin Toughness	Flavor
'Bish'	73AB	75*	73A	72*	73*	75A	70*
(original)							
'Bish'	70B	75*	72AB	72*	75*	70B	72*
(99-TC)							
'Bish'	75A	75*	73A	73*	75*	75A	72*
(00-TC)							
'Camarosa'	75A	75*	70B	72*	77*	75A	70*
'Chandler'	73AB	75*	72AB	75*	77*	73AB	68*
'NCS 93-05'	75A	75*	73A	72*	77*	75A	75*

*Indicates no significant differences observed.
Values designated with different letters are statistically different.

Table 25 provides fruit characteristics of 'Bish' grown at the Upper Piedmont Research Station, Reidsville, N.C. in 2002. Values were of averages across Sep. 5, 2001 and Sep. 12, 2001 planting dates with mean separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 25

Selection	Appearance	Firmness	Symmetry	Skin Color	Flesh Color	Skin Toughness	Flavor
'Bish'	77A	77B	75A	77A	71B	78B	79A
(original)							
'Bish'	77A	78B	75A	77A	71B	78B	78A
(99-TC)							
'Bish'	75A	77B	74A	77A	71B	78B	77A
(00-TC)							
'Camarosa'	72B	84A	71B	76AB	77A	82A	71C
'Chandler'	70B	75C	70B	75B	75A	74C	74B

Values designated with different letters are statistically different.

Table 26 provides fruit characteristics of 'Bish' grown at Fletcher, N.C. in 2002. Values were of averages across Aug. 15, 2001 and Aug. 22, 2001 planting dates with mean separation by Duncan's Multiple Range Test with $p \leq 0.05$.

TABLE 26

Selection	Appearance	Firmness	Symmetry	Skin Color	Flesh Color	Skin Toughness	Flavor
'Bish' (OR)	78A	76B	75A	80A	72A	77BC	77A
'Bish' (MD)	77A	77B	75A	80A	72A	78BC	78A
'Camarosa'	73B	80A	70BC	78AB	74A	85A	68C
'Chandler'	72B	72C	70BC	78AB	74A	75C	74AB
'Gem Star'	75AB	76B	72AB	70C	64B	79B	72C

Values designated with different letters are statistically different.

'Bish' was equal to or better than 'Chandler' for fruit appearance, firmness, symmetry, skin toughness and flavor. Further, 'Bish' was superior to 'Camarosa' for flavor, and equal to or better than 'Camarosa' for appearance and symmetry. 'Camarosa' fruit are extremely firm and also have excellent skin toughness. 'Bish' fruit were neither as firm nor as tough as 'Camarosa', but were satisfactory for both of these traits. In these subjective evaluations 'Bish' was equal to both 'Chandler' and 'Camarosa' for skin color, and poorer than these varieties for flesh color, but still satisfactory for the latter trait. The skin of 'Bish' was uniformly glossy and dark red, but slightly uneven in color (FIG. 1). The skin color on the lighter side of the fruit matches Royal Horticultural Society (RHS) color chip 53A, and the darker side chip 187A. Flesh color of 'Bish' fruit matches RHS color chip 33A, and the central vascular cylinder was whitish (FIG. 2). Hollow centers occur in 80% of the berries. Where these occur, the mean diameter is 5 mm. The range is from 2–10(14) mm (observed May 2004 in Raleigh, N.C.). The achenes on the surface were yellow and slightly indented (sunken), so that the outer surface of the achene is even with the surface of the receptacle tissue between the achenes. As observed in May 2004, in Raleigh, N.C., the color of the achenes on the exposed side of the fruit is RHS color chip 53A to 53B. Achenes on the side away from the sun correspond to R.H.S. color chip 1C in color.

Subjective observations indicate that the shape of the fruit of 'Bish' ranges primarily from blunt conic (with some primary fruit) to conic (FIG. 1). The calyx is medium in size and reflexed (FIG. 1). Fruit shape of 'Bish' was more uniform than 'Chandler' across primary, secondary, and higher order fruits (Table 27).

TABLE 27

Selection		Length/Width Ratio		
		Primary	Secondary	Tertiary, etc.
'Chandler'	Mean	1.22	1.12	1.29
	Range	1.00–1.44	1.00–1.39	1.00–1.63
'Bish'	Mean	1.19	1.18	1.19
	Range	1.00–1.34	1.05–1.24	1.07–1.41

Fruit shape characteristics were from plants grown in Fletcher, NC in 2002.

Flowers and Flowering Characteristics: Both primary and secondary flowers of 'Bish' were slightly larger than 'Chandler' (Table 28).

TABLE 28

Selection		Calyx Diameter (mm)		Corolla Diameter (mm)	Sepal Number
		Outer	Inner		
'Chandler'	<u>Primary</u>				
	Mean	30	31	31	13
	Range	27–34	28–34	27–35	13–14
	<u>Secondary</u>				
'Bish'	<u>Primary</u>				
	Mean	25	27	30	13
	Range	23–27	24–29	28–32	11–15
	<u>Secondary</u>				
	<u>Secondary</u>				
	Mean	32	34	34	11
	Range	30–35	31–40	31–40	10–12
	<u>Secondary</u>				
	Mean	28	30	31	11
	Range	25–30	28–32	30–33	10–12

TABLE 28-continued

Selection	Petal Number	Petal Length (mm)	Petal Width (mm)	
'Chandler'	<u>Primary</u>			
	Mean	7	12	12
	Range	6–7	9–14	8–13
	<u>Secondary</u>			
'Bish'	<u>Primary</u>			
	Mean	6	10	11
	Range	5–7	9–13	9–13
	<u>Secondary</u>			
	<u>Secondary</u>			
	Mean	6	13	11
	Range	5–6	8–16	7–14
	<u>Secondary</u>			
	Mean	5	12	11
	Range	5–6	9–14	8–12

Flower measurements were from plants grown in Reidsville, NC in 2003.

For both 'Bish' and 'Chandler', the diameter of the outer calyx of primary flowers is generally equal to that of the corolla, and slightly smaller than the corolla on secondary flowers. The basal half of the calyx is fused (calyx tube) and adherent to the fruit. The apical half of the calyx lobes are free and typically range from clasping to perpendicular to slightly reflexed in pose on each fruit. The calyx is inserted beneath the fruit, with the calyx tube forming a concave indentation in the base of the fruit. Sepal number averaged 11 for 'Bish' and 13 for 'Chandler'. Mean petal number was also slightly less on both primary and secondary petals for 'Bish'. Petal length for 'Bish' was somewhat greater than width, while for 'Chandler' they were essentially the same. The petal color of flowers produced by 'Bish' is white, corresponding closest to RHS color chip 155C, however the petals are whiter than RHS 155C or any of the other RHS color chips. The average date of first flowering for 'Bish' was three days earlier than for 'Chandler', while the range in first flowering dates was very similar for both varieties (Table 29).

TABLE 29

Selection	Date of First Flowering		Date of First Leaf Emergence	
	Mean	Range	Mean	Range
'Chandler'	March 13	March 7–22	March 10	March 7–15
'Bish'	March 10	March 5–22	March 15	March 10–21

Observations were of plants grown in Reidsville, NC in 2003.

First leaf emergence in spring was, on average, five days later than the date of first flowering for 'Bish', while average first leaf emergence was three days earlier than first flowering for 'Chandler'. For 'Bish', inflorescence height is generally equal to foliage canopy height.

As observed in May 2004 in Raleigh N.C., the mean fruiting truss length was 21 cm, with a range of (18)19–24 (27) cm. The average number of crowns was 4 (range: 3–5).

Plants and Foliage: Fruiting plants of 'Bish' are generally of equal vigor to 'Chandler'. Mean canopy height was equal for the two varieties, although the range of variation in canopy heights was greater for 'Chandler' (Table 30).

TABLE 30

Character	Variety	
	'Chandler'	'Bish'
Canopy Height (cm)	Mean	26
	Range	22–30
<u>Mid-tier Leaflet</u>		
Length (mm)	Mean	67
	Range	55–80
Width (mm)	Mean	62
	Range	55–72
<u>Mid-tier Leaf</u>		
Length (mm)	Mean	87
	Range	70–105
Width (mm)	Mean	119
	Range	95–166
Leaflets/Leaf	3 (rarely 4)	3
Leaf Convexity	Mostly Flat-Convex Some Flat	Mainly convex
<u>Leaf Serrations</u>		
Number	Many	Many
Shape	Semi-Pointed	Semi-Pointed
Leaf Pubescence	Moderate	Moderate
Petiole Pubescence		
Density	Moderate to Heavy	Heavy
Direction	Acropetal to Perpendicular	Mostly Basipetal

Observations were of plants grown in Reidsville, NC in 2003.

Individual leaflets of 'Bish' were very similar in size to those of 'Chandler'. Leaflets of 'Bish' tend to be broadly elliptic-ovate, while those of 'Chandler' are generally broadly elliptic. Leaves of 'Bish' are similar in length and somewhat broader than 'Chandler'. 'Bish' leaves are generally convex (FIG. 3) while those of 'Chandler' are mostly flat-convex. Pubescence on leaves of 'Bish' is moderate, while petiole pubescence is heavy and mostly basipetal in direction. 'Bish' leaves generally show a slight puckering between the veins. The color of the adaxial surface of the leaves of 'Bish' matches RHS color chip 137B, and the abaxial surface corresponds to RHS color chip 147C. One notable feature of 'Bish' is the presence of two leafy petiole bracts (FIG. 3). These structures occur consistently on mid-tier leaves. They are typically quite prominent with a

mean length of 43 mm and a range of 20–55 mm. Similar petiole bracts also occur on the variety 'Sweet Charlie'. However, 'Sweet Charlie' can be distinguished from 'Bish' by having lighter green and more strongly convex leaves. Petiole bracts only occur occasionally on 'Chandler' and 'Camarosa' and have been observed to be mostly rudimentary when they do occur.

'Bish' is more similar in overall morphological appearance to its 'Delmarvel' parent than to any other current variety. The primary morphological difference between 'Bish' and 'Delmarvel' is the absence of petiole bracts on the latter variety. In addition, 'Delmarvel' is not adapted to annual hill culture, which 'Bish' is.

Disease Resistance: One of the major disadvantages of 'Chandler' and 'Camarosa' varieties in annual hill culture throughout the Southeastern region of the United States is extreme susceptibility to anthracnose fruit rot caused by *Colletotrichum acutatum*. With the industry-wide adoption of these two varieties, anthracnose fruit rot is now the primary disease limiting strawberry production in this region. Field reaction of 'Bish', 'Chandler' and 'Camarosa' to anthracnose fruit rot is presented in Tables 1–14. The extent of anthracnose fruit rot varied widely from year to year and site to site. This variation is due both to the level of inoculum available and weather. Warm, humid and especially rainy weather is highly favorable to disease development. Consistently, 'Bish' was significantly more resistant to anthracnose fruit rot than either 'Chandler' or 'Camarosa'. Throughout the test period, 'Bish' never exceeded 4.8%, by weight, of fruit infected, while both 'Chandler' and 'Camarosa' had up to 50% infection. For annual hill culture strawberries in the Upper South/Middle Atlantic regions, carry-over plantings for a second year is an important consideration, and anthracnose susceptibility has been the main deterrent to this practice. The anthracnose resistance of 'Bish' makes it adaptable to carry-over in these regions. 'Bish' is at least moderately resistant to other fungal leaf scorch (*Diplocarpon earliana*), leaf blight (*Phomopsis obscurans*) and powdery mildew (*Spaerotheca macularis*).

That which is claimed is:

1. A new and distinct variety of *Fragaria xananassa* Duch. plant named 'Bish', substantially as described and illustrated herein.

* * * * *




FIG. 1




FIG. 2




FIG. 3