

(12) **United States Plant Patent**  
**Bessho et al.**

(10) **Patent No.:** **US PP15,614 P2**  
(45) **Date of Patent:** **Mar. 1, 2005**

(54) **CALIBRACHOA PLANT NAMED**  
**'KAKEGAWA S64'**

(50) Latin Name: *Calibrachoa* species  
Varietal Denomination: **Takegawa S64**

(75) Inventors: **Masao Bessho**, Kakegawa (JP);  
**Toshimi Ohga**, Kikugawa (JP)

(73) Assignee: **Sakata Seed Corporation**, Yokohama  
(JP)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 36 days.

(21) Appl. No.: **10/801,908**

(22) Filed: **Mar. 16, 2004**

(51) **Int. Cl.**<sup>7</sup> ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./263**

(58) **Field of Search** ..... **Plt./263, 356**

*Primary Examiner*—Kent Bell

(74) *Attorney, Agent, or Firm*—Jondle & Associates P.C.

(57) **ABSTRACT**

'Takegawa S64' is a new variety of *Calibrachoa* plant. This  
new variety has lighter purple petal lobes and dark green  
foliage.

**1 Drawing Sheet**

**1**

Genus/species: *Calibrachoa* species.  
Varietal denomination: 'Takegawa S64'.

**BACKGROUND OF THE INVENTION**

'Takegawa S64' originated from a hybridization made in  
November 1998 in Kakegawa, Japan. The female parent was  
a *Calibrachoa* 'Liracashower Blue' (U.S. Plant Pat. No.  
9,885). The male parent was a *Calibrachoa* breeding line  
with deep rose colored flowers and mounding habit and  
short internode length known as 97-1176 (not patented).

In February 1999, F<sub>1</sub> seed from this cross was sown and  
later transplanted outdoors in Kakegawa, Japan. The F<sub>1</sub>  
plants were blue, rose or pink in flower color and ranged  
from semi-creeping to compact in habit. Three, single-plant  
selections were made from the F<sub>1</sub> plants based on their blue  
flower color and intercrossed to create an F<sub>2</sub> generation. In  
August 1999, seed from the F<sub>2</sub> generation was sown and  
later transplanted outdoors. F<sub>2</sub> plants ranged from extra  
compact to mounding in habit and all flowers were violet-  
blue in flower color. Two, single-plant selections were made  
from the F<sub>2</sub> generation based on their extra compact habit  
and vegetatively propagated. In August 2002, the two selec-  
tions were evaluated in 9 cm hanging pots in a greenhouse  
as well as in an open field. One selection was chosen based  
on the trial.

The selection was further evaluated from new vegetative  
plants in Salinas, Calif. during 2003. The selection was  
subsequently named 'Takegawa S64'. 'Takegawa S64' was  
asexually reproduced by stem cuttings in Salinas, Calif. and  
was determined to reproduce true to type in successive  
generations of asexual propagation.

**DESCRIPTION OF PHOTOGRAPH**

This new *Calibrachoa* plant is illustrated by the accom-  
panying photograph which shows blooms and foliage of the  
plant in full color. The colors shown being as true as can be  
reasonably obtained by conventional photographic proce-  
dures.

FIG. 1 shows a close-up view of flowers.

FIG. 2 shows the entire plant.

**2**

**DESCRIPTION OF THE GENUS CALIBRACHOA**  
**LLAVE & LEX**

The genus *Petunia* was originally established in 1803 by  
A. L. Jussieu, who described both *P. parviflora* and *P.*  
*nyctaginiflora* as type species. Using a non-horticultural  
system that selected the first mentioned species as the type  
species (lectotype), N. L. Britton and H. A. Brown declared  
*P. parviflora* as the type species for *Petunia* in 1913.

During the 1980's and 1990, H. J. Wijsman published a  
series of articles regarding the ancestry of *P. hybrida*, the  
Garden *Petunia*, and the inter-relationship of several species  
classified as *Petunia*. These studies discovered that *P.*  
*hybrida* and its ancestral species, *P. nyctaginiflora* (= *P.*  
*axillaris*) and *P. violacea* (= *P. integrifolia*), possessed 14  
pairs of chromosomes while several other species, including  
*P. parviflora*, possessed 18 pairs of chromosomes. Since *P.*  
*parviflora* was the lectotype species for the *Petunia* genus,  
Wijsman and J. H. de Jong proposed transferring the 14  
chromosome species to the genus *Stimoryne*. Horticulturists  
opposed reclassifying the Garden *Petunia* and in 1986,  
Wijsman proposed the alternative of making *P. nyctagini-*  
*flora* the lectotype species for *Petunia* and transferring the  
18 chromosome species to another genus. The I. N. G.  
Committee adopted this proposal. By 1990, Wijsman had  
transferred several species, including *P. parviflora* (= *C.*  
*parviflora*) to *Calibrachoa*, originally established by Llave  
and Lexarza in 1825. *Calibrachoa parviflora* (= *C. mexicana*  
Llave & Lexarza) is now the type species for the genus  
*Calibrachoa*.

Classification of the current *Petunia* and *Calibrachoa*  
species is still in progress. New species are also being  
identified. Consequently, a proper description has not been  
written for the *Calibrachoa* genus. *Calibrachoa* can,  
however, be distinguished from *Petunia* based on the higher  
chromosome number, chromosome morphology, plant  
branching habit and type of flower bud aestivation. Whereas  
*Petunia* species bear a flower peduncle and one new stem  
from a node, *Calibrachoa* bear a flower peduncle and three  
stems. *Petunia* species have a cochlear corolla bud, a single  
outermost petal covers the other four, radially folded and  
terminally contorted petals. *Calibrachoa* flower buds are flat  
with all five petals linearly folded and the two lower petals  
forming a cover around the three other petals and fused  
together.



In U.S. Plant Pat. No. 9,885, the Genus for 'Liracashower Blue' was identified as *Petunia*. Since that time, as discussed above, the inventors have learned that the Genus *Petunia* had been split by the I. N. G., and this particular variety, because of its chromosome number and bud aestivation is more accurately characterized as a member of the *Calibrachoa* Genus.

#### ENVIRONMENTAL CONDITIONS FOR PLANT GROWTH

The terminal 1.0 to 1.5 inches of an actively growing stem was excised. The vegetative cuttings were propagated in five to six weeks. The base of the cuttings were dipped for 1 to 2 seconds in a 1:9 solution of DIP 'N GROW (1 DIP 'N GROW: 9 water) root-inducing solution immediately prior to sticking into the cell trays. Cuttings were stuck into plastic cell trays having 98 cells and containing a moistened peat moss-based growing medium. The cuttings were misted with water from overhead for 10 seconds every 30 minutes until sufficient roots were formed.

Rooted cuttings were transplanted and grown in 20 cm diameter plastic pots in a glass greenhouse located in Salinas, Calif. Pots contained a peat moss-based growing medium. Soluble fertilizer containing 20% nitrogen, 10% phosphorus and 20% potassium was applied once a day or every other day by overhead irrigation. Pots were top-dressed with a dry, slow release fertilizer containing 20% nitrogen, 10% phosphorus and 18% potassium. The typical average air temperature was 24° C.

#### DETAILED DESCRIPTION OF THE NEW PLANT

Data below collected on plants four months from rooted cutting and transplanted into 20.0 cm diameter pots. Color references are to the R.H.S. Colour Chart of The Royal Horticultural Society of London (R.H.S.). The following traits and characteristics describe the new variety.

##### Classification:

*Family*.—Solanaceae.

*Species*.—*Calibrachoa* sp.

*Common names*.—*Calibrachoa*.

##### Parentage:

*Female parent*.—*Calibrachoa* 'Liracashower Blue' (U.S. Plant Pat. No. 9,885).

*Male parent*.—Breeding line 97-1176 (not patented).

##### Growth:

*Habit*.—Decumbent.

*Height*.—13.0–15.0 cm.

*Spread*.—40.0–45.0 cm for one plant when grown in a 20.0 cm diameter pot.

*Life cycle*.—Perennial.

*Form*.—Branching, dense, compact (shorter internode lengths).

*Time to produce a rooted cutting*.—6 weeks.

*Time to bloom from propagation*.—10 weeks.

*Flowering requirements*.—Will flower so long as day length is greater than 12 hours and temperatures exceed 13° C.

*Resistance/susceptibility*.—Excellent resistance to rain, heat and drought. Will not tolerate temperature below 10° C. Plants are susceptible to Botrytis, powdery mildew, various stem and root rots and certain viruses, like Tobacco Mosaic Virus and Impatiens Necrotic Spotted Virus. Plants can be infested

with aphids, leafminer, whitefly and various Lepidoptera.

##### Stems:

*Stem color*.—RHS 144B (yellow-green).

*Anthocyanin color*.—RHS N77A (purple).

*Pubescence*.—Heavy.

*Pubescence color*.—RHS N155A (white).

*Stem description*.—Round.

*Stem length*.—2.5 cm–3.0 cm.

*Stem diameter*.—2.0 mm.

*Internode length*.—0.5–1.0 cm.

##### Leaves:

*Leaf arrangement*.—Alternate.

*Leaf shape*.—Elliptical.

*Leaf tip*.—Mucronate.

*Leaf base*.—Decurrent.

*Leaf margin*.—Entire.

*Leaf surface*.—Rough, dull.

*Leaf length*.—1.0 cm.

*Leaf width*.—0.5 cm.

*Leaf color*.—Upper: RHS 137A (green); Lower: RHS 138B (green).

*Leaf surface pubescence*.—Slight.

*Leaf surface pubescence color*.—RHS N155B (white).

*Petiole length*.—2.0 mm.

*Petiole color*.—RHS 138B (green).

*Venation*.—Pinnate.

##### Flowers:

*Inflorescence type*.—Solitary.

*Flowering habit*.—Indeterminate.

*Duration of flower life*.—5 days.

*Shape*.—The flowers are funnel shaped with five fissures.

*Flower depth*.—2.0 cm–2.5 cm.

*Floral tube length*.—0.8 cm–1.0 cm.

*Floral tube diameter*.—0.4 cm–0.5 cm.

*Flower diameter*.—2.0 cm.

*Calyx*.—5 sepals, free.

*Sepal shape*.—Lanceolate.

*Sepal apex*.—Mucronate.

*Sepal margin*.—Entire.

*Sepal color*.—RHS 143A (green).

*Bud shape*.—Ovate.

*Bud length*.—1.5 cm.

*Bud diameter*.—0.4 cm–0.5 cm.

*Bud surface*.—Pubescent.

*Bud color*.—RHS 68A (violet).

*Peduncle length*.—1.0 cm.

*Peduncle color*.—RHS 144C (yellow-green).

*Ovary*.—Superior.

*Pistil number*.—1.

*Stigma color*.—RHS 150C (yellow).

*Style color*.—RHS 149B (yellow-green).

*Corolla*.—5 petals, fused.

*Petal shape*.—Spatulate.

*Petal apex*.—Truncate.

*Petal margin*.—Entire.

*Petal pubescence*.—Glabrous.

*Petal size*.—1.0 cm×2.0 cm.

*Petal color*.—Lobes: upper RHS 86A (violet) with RHS 83A (violet) veins; lower about lighter than RHS 86A (violet) with RHS 83A (violet) veins; Corolla tube: inner RHS 14D (yellow-orange) with RHS 83A (violet) veins; outer RHS 10D (yellow) with RHS 83A (violet) veins.

*Stamen number.*—5, free.  
*Stamen color.*—RHS 150C (yellow-green).  
*Pollen color.*—RHS 9B (yellow).  
*Fragrance.*—Absent.  
*Seed production.*—None.

COMPARISON WITH MOST SIMILAR VARIETY

‘Kakegawa S64’ is a distinct variety of *Calibrachoa* owing to its short internode length, which leads to a compact growth habit, and lighter purple petal lobe color. ‘Kakegawa S64’ is most similar to the variety ‘Liracashower Blue’ (U.S. Plant Pat. No. 9,885); however, there are differences as shown in Table 1 below.

TABLE 1

	‘Kakegawa S64’	‘Liracashower Blue’
Internode Length	0.5–1.0 cm	1.5–2.0 cm
Petal Color Upper	RHS 86A (violet) is the upper petal color. It has more blue in it when compared to ‘Liracashower Blue’. The corolla tube base is RHS 14D (yellow-orange)	RHS 88A (violet) is the upper petal color, that is more purple compared to ‘Kakegawa S64’. Fades slightly into RHS 88B (violet) with aging.

TABLE 1-continued

‘Kakegawa S64’	‘Liracashower Blue’
with veins running through of RHS 83A (violet).	Throat shows pale white to cream to light green yellow tones.

Some differences between ‘Kakegawa S64’ and its parental lines are shown in Table 2 below.

TABLE 2

Characteristic	‘Kakegawa S64’	Male (97-1176)	Female (‘Liracashower Blue’)
Petal Color	RHS 86A (violet)	Deep rose	RHS 88A (violet)
Habit	Decumbent	Mounding	Decumbent
Internode Length	0.5–1.0 cm	0.5–1.0 cm	1.5–2.0 cm

What is claimed is:

1. A new and distinct cultivar of *Calibrachoa* plant as shown and described herein.

\* \* \* \* \*





Fig.1



Fig.2