



US00PP15675P2

(12) **United States Plant Patent**  
**Sakazaki**

(10) **Patent No.:** **US PP15,675 P2**  
(45) **Date of Patent:** **Mar. 15, 2005**

(54) **VERBENA PLANT NAMED 'USBENAS24'**

(50) Latin Name: *Verbena hybrida*  
Varietal Denomination: **USBENAS24**

(75) Inventor: **Ushio Sakazaki**, Shiga (JP)

(73) Assignee: **Plant 21 LLC**, San Marco, CA (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/835,382**

(22) Filed: **Apr. 29, 2004**

(51) **Int. Cl.**<sup>7</sup> ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./308**

(58) **Field of Search** ..... **Plt./308**

*Primary Examiner*—Anne Marie Grunberg

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Verbena* plant named 'USBENAS24', characterized by its low and outwardly spreading plant habit; vigorous growth habit; numerous large pink-colored flowers; and tolerance to Powdery Mildew.

**1 Drawing Sheet**

**1**

Botanical classification/cultivar designation: *Verbena hybrida* cultivar USBENAS24.

**BACKGROUND OF THE INVENTION**

The present Invention relates to a new and distinct cultivar of *Verbena* plant, botanically known as *Verbena hybrida*, and hereinafter referred to by the name 'USBENAS24'.

The new *Verbena* is a product of a planned breeding program conducted by the Inventor in Hikone, Shiga, Japan. The objective of the breeding program is to develop new disease-resistant and high temperature-tolerant *Verbena* cultivars with semi-upright plant habit and attractive flower coloration.

The new *Verbena* originated from a cross-pollination made by the Inventor on May 25, 1999 of an unidentified selection of *Verbena hybrida*, not patented, as the female, or seed, parent with a proprietary selection of *Verbena hybrida* identified as code number VJS9-3, not patented, as the male, or pollen, parent. The cultivar USBENAS24 was discovered and selected by the Inventor as a flowering plant within the progeny from the aforementioned cross-pollination in a controlled environment in Bonsall, Calif. on Jun. 10, 2000.

Asexual reproduction of the new cultivar by cuttings at Bonsall, Calif., since Jul. 1, 2000, has shown that the unique features of this new *Verbena* are stable and reproduced true to type in successive generations of asexual reproduction.

**SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'USBENAS24'. These characteristics in combination distinguish 'USBENAS24' as a new and distinct cultivar:

1. Low and outwardly spreading plant habit.
2. Vigorous growth habit.
3. Numerous large pink-colored flowers.
4. Tolerant to Powdery Mildew.

In side-by-side comparisons conducted in Hikone, Shiga, Japan, plants of the new *Verbena* differed from plants of the female parent selection in the following characteristics:

1. Plants of the new *Verbena* were more vigorous than plants of the female parent selection.

**2**

2. Plants of the new *Verbena* had larger flowers than plants of the female parent selection.

3. Plants of the new *Verbena* flowered earlier than plants of the female parent selection.

5 In side-by-side comparisons conducted in Hikone, Shiga, Japan, plants of the new *Verbena* differed from plants of the male parent selection in the following characteristics:

1. Plants of the new *Verbena* were more upright than plants of the male parent selection.

10 2. Plants of the new *Verbena* flowered earlier than plants of the male parent selection.

15 The new *Verbena* can be compared to the cultivar, Tapien Pink, not patented. However, in side-by-side comparisons conducted in Bonsall, Calif., plants of the new *Verbena* differed from plants of the cultivar Tapien Pink in the following characteristics:

1. Plants of the new *Verbena* were more upright than plants of the cultivar Tapien Pink.

20 2. Plants of the new *Verbena* had taller inflorescences than plants of the cultivar Tapien Pink.

3. Plants of the new *Verbena* flowered earlier than plants of the cultivar Tapien Pink.

25 **BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

30 The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Verbena*.

35 The photograph at the top of the sheet comprises a side perspective view of a typical plant of 'USBENAS24' grown in a container.

The photograph at the bottom of the sheet is a close-up view of typical flowers of 'USBENAS24'.

40 **DETAILED BOTANICAL DESCRIPTION**

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary signifi-



cance are used. Plants grown in 15-cm containers were used for the aforementioned photographs and following description. Plants had been growing for about ten weeks when the photographs and the description were taken. Plants were grown under conditions which closely approximate commercial production conditions during the autumn in Bonsall, Calif. in an outdoor nursery. During the production period, day temperatures ranged from about 18 to about 35° C. and night temperatures ranged from about 16 to about 21° C.

Botanical classification: *Verbena hybrida* cultivar USBENAS24.

Parentage:

*Female parent.*—Unidentified selection of *Verbena hybrida*, not patented.

*Male parent.*—Proprietary selection of *Verbena hybrida* identified as code number VJS9-3, not patented.

Propagation:

*Type cutting.*—Vegetative tip cuttings.

*Time to initiate roots.*—About 7 days at 18° C.

*Time to develop roots.*—About 21 days at 18° C.

*Root description.*—Fibrous; whitish in color.

*Rooting habit.*—Freely branching.

Plant description:

*General appearance.*—Low and outwardly spreading plant habit; eventually somewhat trailing plant habit.

*Growth and branching habit.*—Freely basal branching; about seven lateral branches develop per plant. Pinching, that is, removal of the terminal apices, enhances branching with lateral branches potentially forming at every node. Vigorous growth habit; moderate growth rate.

*Plant height.*—About 15 cm.

*Plant diameter or spread.*—About 37 cm.

*Lateral branch description.*—Length: About 27 cm. Diameter: About 2 mm. Internode length: About 3 cm. Texture: Pubescent. Color: 146A.

*Foliage description.*—Arrangement: Opposite, simple. Length: About 2.8 cm. Width: About 3 cm. Shape: Deltoid. Apex: Acute. Base: Attenuate. Margin: Deeply dissected. Texture, upper and lower surfaces: Pubescent; slightly coarse. Venation pattern: Pinnate. Color: Developing foliage, upper surface: 146A. Developing foliage, lower surface: 146B. Fully expanded foliage, upper surface: 147A. Fully expanded foliage, lower surface: 147B. Venation, upper surface: 147B. Venation, lower surface: 147C. Petiole: Length: About 1.5 cm. Diameter: About 1.5 mm. Texture, upper and lower surfaces: Pubescent. Color, upper and lower surfaces: 146B.

Flower description:

*Flower type and habit.*—Single upright salverform flowers arranged on hemispherical corymbs. Freely

flowering with about 46 flowers and flower buds per inflorescence. Inflorescences positioned above and beyond the foliage. Flowers last about five to seven days under greenhouse conditions. Corollas not persistent. Flowers not fragrant.

*Flowering season.*—In the garden, flowering is continuous from spring until fall.

*Inflorescence height.*—About 4.5 cm.

*Inflorescence diameter.*—About 5 cm.

*Flower size.*—Diameter: About 1.5 cm. Height (depth): About 2.5 cm.

*Flower buds.*—Length: About 1 cm. Diameter: About 2 mm. Shape: Elongate. Color: 75C.

*Petals.*—Quantity/arrangement: Five per flower fused at base. Lobe length: About 8 mm. Lobe width: About 6 mm. Shape: Cordate. Apex: Emarginate. Margin: Entire. Texture, upper and lower petal surfaces: Glabrous, smooth; velvety. Texture, throat and tube: Pubescent. Color: When opening, upper surface: 75A. When opening, lower surface: 75D. Fully opened, upper surface: Towards the margins, 75D; towards the throat, 75A. Fully opened, lower surface: 75D. Throat: Close to 155D. Tube: Close to 157B.

*Sepals.*—Quantity/arrangement: Five, fused into an elongated tube. Length: About 9 mm. Diameter: Less than 1 mm. Shape: Linear. Apex: Acuminate. Margin: Entire. Texture, upper and lower surfaces: Pubescent. Color, upper surface: 146B. Color, lower surface: 146A.

*Peduncles.*—Length: About 6.5 cm. Diameter: About 1.5 mm. Strength: Strong. Angle to stem: Horizontal to curving upright. Texture: Pubescent. Color: 146B.

*Reproductive organs.*—Stamens: Quantity: Four per flower. Anther shape: Oval. Anther length: Less than 1 mm. Anther color: 150A. Pollen amount: Scarce. Pollen color: 150A. Pistils: Quantity: One per flower. Pistil length: About 1.8 cm. Style length: About 1.5 cm. Style color: 145C to 145D. Stigma shape: Bi-parted. Stigma color: 145A. Ovary color: 144A.

*Seed/fruit.*—Seed and fruit production has not been observed.

*Disease/pest resistance:* Plants of the new *Verbena* have been observed to be tolerant to Powdery Mildew. Plants of the new *Verbena* have not been observed to be resistant to other pathogens and pests common to *Verbena*.

*Temperature tolerance:* Plants of the new *Verbena* have been observed to be tolerant to temperatures ranging from -4 to 40° C.

It is claimed:

1. A new and distinct cultivar of *Verbena* plant named 'USBENAS24', as illustrated and described.

\* \* \* \* \*



