

US00PP15719P2

(12) **United States Plant Patent**
Boeder

(10) **Patent No.: US PP15,719 P2**
(45) **Date of Patent: Apr. 12, 2005**

(54) **CHRYSANTHEMUM PLANT NAMED ‘AGAPI’**

(50) Latin Name: *Chrysanthemum morifolium*
Varietal Denomination: **Agapi**

(75) Inventor: **Mark Roland Boeder**, The Hague
(NL)

(73) Assignee: **Chrysanthemum Breeders**
Association, N.V. (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/756,362**

(22) Filed: **Jan. 14, 2004**

(51) **Int. Cl.⁷ A01H 5/00**

(52) **U.S. Cl. Plt./287**

(58) **Field of Search Plt./287**

Primary Examiner—Anne Marie Grunberg

(74) *Attorney, Agent, or Firm*—Parkhurst & Wendel, L.L.P.

(57) **ABSTRACT**

A *chrysanthemum* plant named ‘Agapi’ characterized by its
medium sized blooms with pink ray florets and prolific
branching; natural season flower date September 8–12;
blooming for a period of 5 weeks.

2 Drawing Sheets

1

BACKGROUND OF THE INVENTION

‘Agapi’ is a product of a breeding and selection program
for outdoor pot mums (garden mums) which had the objec-
tive of creating new *chrysanthemum* cultivars with a deco-
rative type flower, a natural season flower date around
September 8–12; blooming for a period of 5 weeks. The new
plant of the present invention comprises a new and distinct
cultivar of *Chrysanthemum* plant ‘Agapi’ is a seedling
resulting from the open pollination among groups of *chry-*
santhemum cultivars maintained under the control of the
inventor for breeding purposes. The new and distinct culti-
var was discovered and selected as one flowering plant by
Mark Roland Boeder on a cultivated field in Rijsenhout
Holland in September 2001. The female parent is identified
by the code 95.3002, not patented. The female parent has
similar blooms to ‘Agapi’ but differs in the center, being less
deeply colored than in ‘Agapi’. The plant has been asexually
reproduced by cuttings in greenhouses at Rijsenhout Hol-
land. The new cultivar has been found to retain its distinctive
characteristics through successive propagations.

BRIEF DESCRIPTION OF THE DRAWINGS

The present invention of a new and distinct variety of
chrysanthemum is shown in the accompanying drawings, the
color being as nearly true as possible with color photographs
of this type.

FIG. 1 shows a plant of the cultivar in full bloom.

FIG. 2 shows the various stages of bloom and foliage of
the new cultivar.

DESCRIPTION OF THE INVENTION

This new variety of *chrysanthemum* is of the botanical
classification *Chrysanthemum morifolium*. The observations
and measurements were gathered from plants grown out
door in Rijsenhout, Holland under natural day length and
temperature and planted in week 23 in 2002. The natural
blooming date of this crop was September 8–13 (week 37).
The average height of the plants was 35 cms. No growth
retardants were used. No tests were done on disease or insect
resistance or susceptibility. No tests were done on cold or
drought tolerance. This new variety produces medium sized
blooms with pink ray florets and a yellow center blooming

2

for a period of 5 weeks. From the cultivars known to
inventor the most similar existing cultivar in comparison to
‘Agapi’ is ‘Lynn’ (U.S. Plant Pat. No. 8,171). When ‘Lynn’
and ‘Agapi’ are being compared the following differences
are noticed: The differences of ‘Lynn’ and ‘Agapi’ are (1)
Color of center of flower. The center of flowers of ‘Lynn’ is
purple, while that of ‘Agapi’ is yellow. (2) Plant vigor. The
vigor of ‘Lynn’ is low, while that of ‘Agapi’ is high.

The following is a description of the plant and character-
istics that distinguish ‘Agapi’ as a new and distinct variety.
The color designations are taken from the plant itself.
Accordingly, any discrepancies between the color designa-
tions and the colors depicted in the photographs are due to
photographic tolerances. The color chart used in this
description is: The Royal Horticultural Society Colour
Chart, edition 1995.

TABLE 1

Botanical Description of cultivar ‘Agapi’	
<u>Bud</u>	
Size	small; cross-section 0.6 cm, height 0.7 cm
Outside Color	Greyed-yellow 160A
Involucral bracts	2 rows, length 7 mm, width 3 mm
Involucral bracts among disc-florets	Not present
Involucral bracts color	Green 138B
<u>Bloom</u>	
Type	Decorative
Size	Medium
Fully Expanded	5 cm
Peduncle length	8.5 cm
Peduncle color	Yellow-green 147D
Number of blooms per branch	Approx. 9 blooms per branch
Performance on the plant	5 weeks
Seeds	Produced in small quantities, ovate grey-brown 199A, 1½ mm in length.
Fragrance	Typical <i>chrysanthemum</i> , slightly
Color	
Center of the flower	Immature Red 51C Mature Yellow 14A The center of the mature flowers is colored by the visible styles from the open disc florets.

TABLE 1-continued

Botanical Description of cultivar ‘Agapi’	
Color of upper surface of the ray-florets	Purple 75A
Color of the lower surface of the ray-florets	Purple 75C
Tonality from Distance	A garden mum with pink flowers and a yellow disc
Color of the uppersurface of the ray-florets after aging of the plant	Purple 75B
<u>Ray florets</u>	
Texture	Upper and under side smooth
Number	150–170
Cross-section	flat
Longitudinal axis of majority	straight
Length of corolla tube	0.5 cm
Ray-floret margin	Entire
Ray-floret length	3 cm
Ray-floret width	0.5 cm
Ratio length/width	high
Shape of tip	rounded
<u>Disc florets</u>	
Disc diameter	1 cm
Distribution of disc florets	Few, only present in mature stage
Shape	Tubular
Color	Yellow-green 154D
Receptacle shape	Domed raised
<u>Reproductive Organs</u>	
Stamen	Present in disc florets only
Stamen color	Yellow-green 144A
Pollen	Produced in small quantity
Pollen color	Yellow 7A
Styles	Thick
Style color	Yellow 14A
Style Length	4 mm
Stigma color	Yellow-green 144A
Stigma Width	1 mm
Ovaries	Enclosed in calyx
<u>Plant</u>	
Form	A outdoor mounded and round garden mum
Growth habit	Spreading
Growth rate	High
Height	35 cm
Width	55 cm
Stem Color	Green 138C
Stem Strength	Weak
Stem Brittleness	Medium

TABLE 1-continued

Botanical Description of cultivar ‘Agapi’	
Stem Anthocyanin Coloration	Absent
Internode length	3 cm
Length of lateral branch	From top to bottom 17 cm
Lateral branch color	Green 138C
Lateral branch, attachment	Medium strength
Branching (average number of lateral branches)	Good with 6–7 breaks after pinching
Natural season blooming date	SEPTEMBER 8–12
<u>Foliage</u>	
Leaf color	Upper side Green 139B Under side Green 138B
Color midvein	Upper side Yellow-green 147D Under side Yellow-green 148D
Size	Medium; length 5.5 cm, width 4 cm
Quantity (number per lateral branch)	20
Shape	Cordiform
Texture upper side	Glabrous
Texture under side	Pubescent
Venation arrangement	Palmate
Shape of the margin	Serrated
Shape of Base of Sinus Between Lateral Lobes	Rounded
Margin of Sinus Between Lateral Lobes	Diverging
Shape of Base	Obtuse
Apex	Mucronate
Petiole length	1.5 cm
Petiole color	Yellow-green 147D

TABLE 2

<u>Differences with the comparison varieties</u>		
	‘Agapi’	‘Lynn’
Color center of flower	Yellow 14A	Red-purple 70A
Vigor	High	Low

I claim:

1. A new and distinct variety of *chrysanthemum* plant as described and illustrated.

* * * * *




FIG. 1




FIG. 2