



US00PP15813P2

(12) **United States Plant Patent**
Kjaer-Larsen

(10) **Patent No.:** **US PP15,813 P2**
(45) **Date of Patent:** **Jun. 28, 2005**

(54) **BEGONIA PLANT NAMED 'BBEATRICE'**

(50) Latin Name: *Begonia*×*hiemalis*
Varietal Denomination: **Bbeatrice**

(75) Inventor: **Henrik Kjaer-Larsen**, Odense (DK)

(73) Assignee: **Begonia Breeders Association B.V.**,
Aalsmeer (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/859,501**

(22) Filed: **Jun. 1, 2004**

(51) **Int. Cl.**⁷ **A01H 5/00**

(52) **U.S. Cl.** **Plt./348**

(58) **Field of Search** **Plt./348**

Primary Examiner—Kent Bell

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Begonia* plant named
'Bbeatrice', characterized by its compact, upright and
mounded plant habit; double flowers with salmon pink-
colored tepals; and excellent flower longevity.

1 Drawing Sheet

1

Botanical classification/cultivar designation: *Begonia*×*hiemalis* cultivar Bbeatrice.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of *Begonia* plant, botanically known as *Begonia*×
hiemalis, commercially known as Elatior Begonia, and
hereinafter referred to by the name 'Bbeatrice'.

The new *Begonia* was discovered and selected by the
Inventor in a controlled environment in Odense, Denmark in
March, 1999, as a naturally-occurring whole plant mutation
of *Begonia*×*hiemalis* 'Netja', not patented. The new *Bego-*
nia was observed as a single plant in a group of flowering
plants of the parent cultivar.

Asexual reproduction of the new *Begonia* by cuttings in
a controlled environment in Odense, Denmark since 1999,
has shown that the unique features of this new *Begonia* are
stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

The cultivar 'Bbeatrice' has not been observed under all
possible environmental conditions. The phenotype may vary
somewhat with variations in environment such as
temperature, daylength and light intensity, without,
however, any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of 'Bbeat-
rice'. These characteristics in combination distinguish
'Bbeatrice' as a new and distinct *Begonia*:

1. Compact, upright and mounded plant habit.
2. Double flowers with salmon pink-colored tepals.
3. Excellent flower longevity.

Plants of the new *Begonia* are most similar to plants of the
parent, the cultivar Netja. Plants of the new *Begonia* differ
primarily from plants of the cultivar Netja in flower color as
tepals of plants of the cultivar Bbeatrice are salmon pink in
color whereas tepals of plants of the cultivar Netja are light
pink in color.

Plants of the new *Begonia* are similar in flower color to
plants of the cultivar Tilly, not patented. However flower
color of plants of the new *Begonia* is more intense salmon
pink than flower color of plants of the cultivar Tilly.

2

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the
overall appearance of the new *Begonia*, showing the colors
as true as it is reasonably possible to obtain in colored
reproductions of this type. Colors in the photograph may
differ slightly from the color values cited in the detailed
botanical description which accurately describe the colors of
the new *Begonia*. The photograph comprises a side perspec-
tive view of a typical flowering plant of 'Bbeatrice'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to
The Royal Horticultural Society Colour Chart, 1995 Edition,
except where general terms of ordinary dictionary signifi-
cance are used. The aforementioned photograph and follow-
ing observations and measurements describe plants grown in
Aalsmeer, The Netherlands during the spring and summer,
and grown under conditions typical of commercial practice
in a glass-covered greenhouse. During the production of the
plants, day and night temperatures ranged from 15 to 20° C.
and light levels were about 18,000 lux. Plants used for the
photograph and the description were grown in 13-cm con-
tainers and were about three months old when the photo-
graph and description were taken.

Botanical classification: *Begonia*×*hiemalis* cultivar Bbeat-
rice.

Commercial classification: Elatior Begonia.

Parentage: Naturally-occurring whole plant mutation of
Begonia×*hiemalis* cultivar Netja, not patented.

Propagation:

Type.—By cuttings.

Time to initiate roots.—About two weeks at tempera-
tures of 20° C.

Time to develop roots.—About five weeks at tempera-
tures of 20° C.

Root description.—Fine, fibrous and well-branched.
Plants of the new *Begonia* have not been observed to
form tubers.

Plant description:

Plant form and habit.—Compact, upright and mounded
plant habit; freely branching with about five or six
basal branches per plant. Moderately vigorous. Veg-

etative shoots are formed at basal nodes and flowering shoots are formed at upper nodes.

Plant height.—About 20 cm.

Plant width.—About 20 to 25 cm.

Leaves.—Arrangement: Alternate, simple. Length: About 9 cm. Width: About 6 to 7 cm. Shape: Ovate; asymmetrical. Apex: Acuminate. Base: Oblique. Margin: Doubly crenate. Texture, upper and lower surfaces: Glabrous, smooth; leathery. Venation pattern: Palmate. Color: Developing and fully expanded leaves, upper surface: Close to 137A. Developing and fully expanded leaves, lower surface: Close to 137C. Venation, upper surface: Close to 137A. Venation, lower surface: Close to 137C. Petiole length: About 4 to 6 cm. Petiole diameter: About 4 mm. Petiole color, upper and lower surfaces: Close to 137A. Stipule quantity: Two per leaf. Stipule size: About 1 cm by 1 cm. Stipule texture, upper and lower surfaces: Smooth, glabrous. Stipule color, upper and lower surfaces: Close to 137C.

Flower description:

Flowering habit.—Double flowers with about 10 to 12 tepals per flower; flowers arranged in axillary cymes. Numerous cymes in flower simultaneously; about 25 to 30 open flowers per plant. Flowers positioned above and beyond the foliage. Flowers not fragrant.

Natural flowering season.—Under natural daylight conditions, plants flower from spring until the fall. Flower initiation and development is induced by long day/short night conditions. Flowering continuous under photoinductive conditions.

Flower longevity.—Individual flowers last about four to five weeks on the plant.

Flowers.—Shape: Oval; double. Orientation: Flat to cupped. Diameter: About 4 to 5 cm. Depth (height): About 1 cm.

Flower buds.—Length: About 1 to 1.5 cm. Diameter: About 2 cm. Color: Close to 38B.

Tepals.—Arrangement: Rosette. Shape: Obovate to rounded. Apex: Rounded. Base: Obtuse. Margin: Entire. Length: About 2 to 3 cm. Width: About 3 to 4 cm. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening and fully opened, upper surface: 37B; towards the margins, 36A. When opening and fully opened, lower surface: 38B.

Flower bracts.—Quantity: Two per flower. Arrangement: Opposite. Length: About 1 cm. Width: About 1 cm. Shape: Rounded. Apex: Rounded. Base: Cuneate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: 146D.

Peduncles.—Angle: About 30 to 45° from vertical. Length: About 4 to 5 cm. Diameter: About 3 to 4 mm. Strength: Strong. Texture: Smooth. Color: 146D overlain with 39B.

Pedicels.—Angle: About 30 to 45° from vertical. Length: About 2 to 3 cm. Diameter: About 2 to 3 mm. Strength: Strong. Texture: Smooth. Color: 146D overlain with 39B.

Reproductive organs.—Stamens: None observed. Pistils: None observed.

Seed/fruit.—Seed and fruit production has not been observed as reproductive organs are not formed.

Disease/pest resistance: Plants of the new *Begonia* have been observed to be resistant to Powdery Mildew. Plants of the new *Begonia* have not been observed to be resistant to pests and other pathogens common to *Begonia*.

It is claimed:

1. A new and distinct cultivar of *Begonia* plant named 'Bbeatrice', as illustrated and described.

* * * * *

