



US00PP15855P2

(12) United States Plant Patent
Boeder(10) Patent No.: US PP15,855 P2
(45) Date of Patent: Jul. 12, 2005(54) CHrysanthemum PLANT NAMED
'ATHENA'(50) Latin Name: *Chrysanthemum morifolium*
Varietal Denomination: Athena(75) Inventor: **Mark Roland Boeder**, The Hague
(NL)(73) Assignee: **Chrysanthemum Breeders
Association, N.V.** (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 59 days.

(21) Appl. No.: 10/756,363

(22) Filed: Jan. 14, 2004

(51) Int. Cl.⁷ A01H 5/00

(52) U.S. Cl. Plt./290

(58) Field of Search Plt./290

Primary Examiner—Anne Marie Grunberg

Assistant Examiner—Annette H Para

(74) Attorney, Agent, or Firm—Parkhurst & Wendel, L.L.P.

(57) ABSTRACT

A *chrysanthemum* plant named 'Athena' characterized by its medium sized blooms with pink ray florets and prolific branching; natural season flower date September 15–20; blooming for a period of 5 weeks.

2 Drawing Sheets

1

BACKGROUND OF THE INVENTION

'Athena' is a product of a breeding and selection program for outdoor pot mums (garden mums) which had the objective of creating new *chrysanthemum* cultivars with a decorative type flower, a natural season flower date around September 15–20; blooming for a period of 5 weeks. The new plant of the present invention comprises a new and distinct cultivar of *Chrysanthemum* plant 'Athena' is a seedling resulting from the open pollination among groups of *chrysanthemum* cultivars maintained under the control of the inventor for breeding purposes. The new and distinct cultivar was discovered and selected as one flowering plant by Mark Roland Boeder on a cultivated field in Rijenhout Holland in September 2001. The plant has been asexually reproduced by cuttings in greenhouses at Rijenhout Holland. The new cultivar has been found to retain its distinctive characteristics through successive propagations.

BRIEF DESCRIPTION OF THE DRAWINGS

The present invention of a new and distinct variety of *chrysanthemum* is shown in the accompanying drawings, the color being as nearly true as possible with color photographs of this type.

FIG. 1 shows a plant of the cultivar in full bloom.

FIG. 2 shows the various stages of bloom and foliage of the new cultivar.

DESCRIPTION OF THE INVENTION

This new variety of *chrysanthemum* is of the botanical classification *Chrysanthemum morifolium*. The observations and measurements were gathered from plants grown out door in Rijenhout, Holland under natural day length and temperature and planted in week 23 in 2002. The natural blooming date of this crop was September 15–20 (week 38). The average height of the plants was 35 cms. No growth retardants were used. No tests were done on disease or insect resistance or susceptibility. No tests were done on cold or drought tolerance. This new variety produces medium sized blooms with pink ray florets and a salmon center blooming for a period of 5 weeks.

2

From the cultivars known to inventor the most similar existing cultivar in comparison to 'Athena' is 'Cefreya' (U.S. Plant Pat. No. 14,939). When 'Cefreya' and 'Athena' are being compared the following differences are noticed:
5 The differences of 'Cefreya' and 'Athena' are (1) Flower colour. The ray-florets of 'Cefreya' are salmon colored, while those of 'Athena' are pink. (2) Longitudinal axis of majority of ray-florets. The ray-florets of 'Cefreya' show a reflexing to twisted curvature, while those of 'Athena' are straight.

The following is a description of the plant and characteristics that distinguish 'Athena' as a new and distinct variety.

The color designations are taken from the plant itself.
15 Accordingly, any discrepancies between the color designations and the colors depicted in the photographs are due to photographic tolerances. The color chart used in this description is: The Royal Horticultural Society Colour Chart, edition 1995.

TABLE 1

Botanical Description of cultivar 'Athena'

| Bud | |
|--------------------------------------|---|
| Size | small; cross-section 1 cm, height 0.5 cm |
| Outside Color | Yellow-green 148D |
| Involucral bracts | 2 rows, length 7 mm, width 3 mm |
| Involucral bracts among disc-florets | Not present |
| Involucral bracts color | Green 138B |
| Bloom | |
| Type | Decorative |
| Size | Medium |
| Fully Expanded | 5 cm |
| Peduncle length | 7 cm |
| Peduncle color | Green 138C |
| Number of blooms per branch | Approx. 9 blooms per branch |
| Performance on the plant | 5 weeks |
| Seeds | Produced in small quantities, ovate grey-brown 199A, 1½ mm in length. |
| Fragrance | Typical <i>chrysanthemum</i> , slightly |
| Color | |
| Center of the flower | Immature Orange-red 31D |
| | Mature Orange-red 31D |

US PP15,855 P2

3

TABLE 1-continued

| Botanical Description of cultivar ‘Athena’ | |
|--|--|
| Color of upper surface of the ray-florets | Red 55C |
| Color of the lower surface of the ray-florets | Red 38D |
| Tonality from Distance | A garden mum with pink flowers and a salmon disc |
| Color of upper surface of ray-florets after aging of the plant | Red 55C |
| <u>Ray florets</u> | |
| Texture | Upper and under side smooth |
| Number | 150–170 |
| Cross-section | Flat |
| Longitudinal axis of majority | Straight |
| Length of corolla tube | 0.5 cm |
| Ray-floret margin | Entire |
| Ray-floret length | 3 cm |
| Ray-floret width | 0.8 cm |
| Ratio length/width | medium |
| Shape of tip | Rounded |
| Disc florets | Absent |
| Receptacle shape | Domed raised |
| <u>Reproductive Organs</u> | |
| Stamen | Absent |
| Pollen | Not produced |
| Styles | Thin |
| Style color | Yellow 13A |
| Style Length | 3 mm |
| Stigma color | Yellow-green 144A |
| Stigma Width | 1 mm |
| Ovaries | Enclosed in calyx |
| <u>Plant</u> | |
| Form | A garden mum outdoor mounded and round |
| Growth habit | Round and spreading |
| Growth rate | Intermediate |
| Height | 35 cm |
| Width | 40 cm |
| Stem Color | Green 138C |
| Stem Strength | Weak |
| Stem Brittleness | Brittle |
| Stem Anthocyanin Coloration | Absent |
| Internode length | 2 cm |
| Length of lateral branch | From top to bottom 16 cm |

4

TABLE 1-continued

| Botanical Description of cultivar ‘Athena’ | |
|--|--|
| Lateral branch color | Green 138B |
| Lateral branch, attachment | Weak |
| Branching (average number of lateral branches) | Good with 7–8 breaks after pinching |
| Natural season blooming date | September 15–20 |
| <u>Foliage</u> | |
| Leaf color | Upper side Green 137A Under side Green 137D |
| Color midvein | Upper side Yellow-green 148D Under side Yellow-green 147D |
| Size | Small; length 4 cm, width 3 cm |
| Quantity (number per lateral branch) | 20 |
| Shape | Cordiform |
| Texture upper side | Glabrous |
| Texture under side | Pubescent |
| Venation arrangement | Palmate |
| Shape of the margin | Serrated |
| Shape of Base of Sinus Between Lateral Lobes | Rounded |
| Margin of Sinus Between Lateral Lobes | Diverging |
| Shape of Base | Obtuse |
| Apex | Mucronate |
| Petiole length | 1 cm |
| Petiole color | Yellow-green 148D |

TABLE 2

Differences with the comparison variety

| | ‘Athena’ | ‘Cefreya’ |
|----------------------------------|----------|----------------------|
| Color ray-florets | Red 55C | Red 37B |
| Longitudinal axis of ray-florets | Straight | Reflexing to twisted |

I claim:

1. A new and distinct variety of *chrysanthemum* plant as described and illustrated.

* * * * *




FIG. 1




FIG. 2