



US00PP15922P3

(12) **United States Plant Patent**
Strope

(10) **Patent No.:** **US PP15,922 P3**
(45) **Date of Patent:** **Aug. 16, 2005**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED
'BALCEBPIEL'**

(50) Latin Name: *Impatiens*×*hawkeri*
Varietal Denomination: **Balcebpiel**

(75) Inventor: **Kerry Strope**, Jefferson City, MO (US)

(73) Assignee: **Ball Horticultural Company**, West
Chicago, IL (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 90 days.

(21) Appl. No.: **10/736,736**

(22) Filed: **Dec. 16, 2003**

(65) **Prior Publication Data**

US 2005/0132455 P1 Jun. 16, 2005

(51) **Int. Cl.**⁷ **A01H 5/00**

(52) **U.S. Cl.** **Plt./318**

(58) **Field of Search** **Plt./318**

(56) **References Cited**

U.S. PATENT DOCUMENTS

PP11,415 P * 6/2000 Trees Plt./318
PP13,698 P2 * 4/2003 Hofmann Plt./318
PP14,864 P2 * 6/2004 Hofmann Plt./318
PP15,223 P2 * 10/2004 Jorna Plt./318

OTHER PUBLICATIONS

UPOV ROM GTITM Computer Database, GTI Jouve
Retrieval Software 2004/04 Citations for Balcebpiel.*

* cited by examiner

Primary Examiner—Kent Bell

Assistant Examiner—W. C. Haas

(74) *Attorney, Agent, or Firm*—Wood, Phillips, Katz, Clark
& Mortimer

(57) **ABSTRACT**

A new and distinct cultivar of New Guinea *Impatiens* plant
named 'Balcebpiel' characterized by its large pink-colored
flowers, medium green-colored foliage, medium upright,
mounded growth habit, and excellent basal branching.

1 Drawing Sheet

1

Latin name of genius and species of plant claimed:
Impatiens hawkeri.

Variety denomination: 'Balcebpiel'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct New
Guinea *Impatiens* plant botanically known as *Impatiens*
hawkeri and hereinafter referred to by the cultivar name
'Balcebpiel'.

The new cultivar was developed by the inventor in a
controlled breeding program during June 2000 at Arroyo
Grande, Calif. The objective of the breeding program was to
develop New Guinea *Impatiens* cultivars with numerous
large flowers, excellent basal branching and upright compact
to moderate growth habits.

The female (seed) parent of 'Balcebpiel' was the propri-
etary *Impatiens hawkeri* selection designated '1585-4' (not
patented), characterized by its upright growth habit, light
pink-colored flowers and medium green-colored foliage.
The male (pollen) parent of 'Balcebpiel' was proprietary
Impatiens hawkeri selection designated 'BFP-2565' (not
patented), characterized by its pink-colored flowers and dark
green-colored foliage. 'Balcebpiel' was discovered and
selected as a single flowering plant within the progeny of the
above stated cross-pollination in January of 2001 at Arroyo
Grande, Calif. and was initially designated '3183-2'.

Asexual reproduction of the new cultivar by terminal stem
cuttings taken since January 2001 at West Chicago, Ill. has
demonstrated that the new cultivar reproduces true to type
with all the characteristics, as herein described, firmly fixed
and retained through successive generations of such asexual
propagation.

2

SUMMARY OF THE INVENTION

It was repeatedly found that the cultivar of the present
invention:

- 5 1. Exhibits large round pink-colored flowers,
- 2. Forms medium green-colored foliage,
- 3. Exhibits a good basal branching character, and
- 4. Exhibits a medium, upright mounded growth habit.

10 Plants of the new cultivar differ from plants of the female
parent primarily in flower color and from plants of the male
parent primarily in foliage color.

15 Of the many commercially available New Guinea *Impa-
tiens* cultivars known to the inventor, 'Balcebpiel' is most
similar to 'BFP-844' (U.S. Plant Pat. No. 11,415). However,
in side-by-side comparisons conducted in West Chicago, Ill.,
plants of the new cultivar differed from plants of 'BFP-844'
in the following characteristics.

- 20 1. Plants of the new cultivar have more branches than
plants of 'BFP-844',
- 2. Plants of the new cultivar have shorter branches than
plants of 'BFP-844', and
- 25 3. Plants of the new cultivar have larger flowers than
plants of 'BFP-844'.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

30 The accompanying photographs show, as nearly true as it
is reasonably possible to make the same in color illustrations
of this type, typical flower and foliage characteristics of the
new cultivar. Colors in the photographs differ slightly from
the color values cited in the detailed description, which
accurately describes the colors of 'Balcebpiel'. The plants
were grown from rooted cuttings for 14 weeks in a green-
35 house at West Chicago, Ill.

FIG. 1 illustrates a side view of the overall growth and flowering habit of 'Balcebpel'.

FIG. 2 illustrates a close-up view of a single flower of 'Balcebpel'.

DETAILED DESCRIPTION

The new cultivar has not been observed under all possible environmental conditions to date. Accordingly, it is possible that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 1995 edition, except where general color terms of ordinary significance are used. The color values were determined on Sep. 11, 2003 between 1:00 and 1:45 p.m. under natural daylight conditions.

The following measurements and comparisons describe plants produced from cuttings taken from stock plants and grown under greenhouse conditions comparable to those used in commercial practice. The plants were grown in 10 cm pots for 14 weeks utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 72° F. during the day and approximately 65° F. during the night. Greenhouse light levels were maintained at approximately 4,000–7,000 footcandles during the day.

Classification:

Botanical.—*Impatiens hawkeri* cultivar 'Balcebpel'.

Parentage:

Female (seed) parent.—Proprietary *Impatiens hawkeri* selection designated '1585-4' (not patented).

Male (pollen) parent.—Proprietary *Impatiens hawkeri* cultivar 'BFP-2565' (not patented).

Propagation:

Type cutting.—Terminal stem.

Time to initiate roots.—Approximately 7–14 days.

Time to develop a rooted cutting.—Approximately 21 days.

Root description.—Fine, fibrous.

Rooting habit.—Freely branching.

Plant description:

Habit of growth.—Compact with good basal branching.

A mature plant, 14 weeks after the planting of a rooted cutting, commonly measures approximately 15 cm from soil level to top of plant plane and approximately 24 cm in width (plant spread).

Plant form.—Upright and mounded.

Branches.—Quantity per plant: Approximately 2.

Length: Approximately 7.4 cm. Diameter: Approximately 9 mm. Texture: Glabrous. Color: 144A. Internode length: Approximately 1.9 cm.

Foliage.—Type: Simple. Arrangement of lower leaves: Alternate. Arrangement of upper leaves: Whorls. Shape: Elliptic. Apex: Acuminate. Base: Attenuate. Margin: Serrulate, ciliate. Texture of upper and lower surfaces: Slightly rugose. Pubescence of upper surface: Moderately papillate. Pubescence on lower surface: None. Venation pattern: Pinnate. Size of mature foliage: Length: Approximately 9.7 cm. Width: Approximately 3.7 cm. Color of mature foliage: Upper surface: Slightly darker than 137A, with veins and mid-vein of 145C. Lower surface: 147B with veins and mid-vein of 145B. Petiole: Length: Approximately 1 cm. Diameter: Approximately 3.4

mm. Texture: Upper and lower surfaces: Glabrous. Color: 145B.

Flower description:

Flower type.—Single, not fragrant with self-cleaning petals.

Flowering habit.—Freely flowering with flowers positioned above the foliage and facing outward or upward.

Natural flowering season.—Year round in greenhouse environment. Flowering is continuous from spring until fall in the garden.

Flower longevity on plant.—Approximately 10 days.

Quantity of flowers and buds per stem.—Approximately 1 fully open flower and 7 buds at any one time.

Mature flower buds (just before opening).—Shape: Ovoid. Length: Approximately 1.8 cm. Diameter: Approximately 1 cm. Color of petals: Darker than 54A.

Flower shape.—Round, flat to slightly cupped.

Flower size.—Diameter: Approximately 6.6 cm. Depth: Approximately 1.7 cm.

Petals.—Number: Five per flower. Shape: Obovate. Margin: Entire. Apex: Emarginate. Base: Superior petal has truncate base, other four petals have attenuate bases. Texture of upper and lower surfaces: Glabrous. Appearance: Iridescent. Size: Superior petal length: Approximately 2.7 cm. Superior petal width: 4.3 cm. Lateral petal length: 3.2 cm. Lateral petal width: 2.8 cm. Lower petal length: 3.5 cm. Lower petal width: 3.1 cm. Color of upper surface of all petals: 47D with venation of 47B and base darker than N57A. Color of lower surfaces of all petals: 49D with venation of 47B, mid-vein of 59C and base of 47C. Petal color fades with age to 54A.

Calyx.—Quantity of sepals: Three, with lower sepal modified into a spur. Sepal shape: Ovate. Sepal apex: Acuminate. Sepal texture of both surfaces: Glabrous. Lateral sepal length: Approximately 1.4 cm. Lateral sepal width: Approximately 6 mm. Lateral sepal color of both surfaces: 145B. Lower sepal length: Approximately 1.5 cm. Lower sepal width: Approximately 7 mm. Lower sepal color of both surfaces: 55D with venation and margin of 55A.

Spur.—Quantity: One per flower. Length: Approximately 5.1 cm. Diameter at flower: 3 mm. Diameter at tip: 1 mm. Aspect: Curved downward. Color: 146C overlaid with 59C.

Peduncles.—Strength: Strong. Angle to stem: Acute. Length: Approximately 4.6 cm. Diameter: 2 mm. Texture: Glabrous. Color: 145B.

Reproductive organs.—Androecium: Five stamens, anthers are fused together forming one organ that surrounds the pistil. Filaments are free. Generally, the anthers shed pollen prior to the stigma becoming receptive. Filament color: 61B. Anther shape: Oval. Anther length: Approximately 3 mm. Anther color: Colorless and translucent. Pollen amount: Moderate. Pollen color: 8D. Gynoecium: Pistil number: One per flower. Pistil length: Approximately 5 mm. Stigma length: 0.5 mm. Stigma color: Colorless and translucent. Style length: 0.5 mm. Style color: 144D and translucent. Ovary length: 4 mm. Ovary diameter: 2 mm. Ovary color: 144A.

Seed and fruit development: Neither seed nor fruit production has been observed.

5

Disease and pest resistance: Resistance to pathogens and pests common to *Impatiens* has not been observed.

What is claimed is:

1. A new and distinct cultivar of New Guinea *Impatiens* plant named 'Balcebpel' substantially as herein shown and described, which:

6

1. Exhibits large round pink-colored flowers,
2. Forms medium green-colored foliage,
3. Exhibits a good basal branching character, and
4. Exhibits a medium upright, mounded grown habit.

* * * * *




Figure 1




Figure 2