



US00PP16702P2

(12) **United States Plant Patent**
Robb

(10) **Patent No.:** **US PP16,702 P2**
(45) **Date of Patent:** **Jun. 20, 2006**

(54) **LAVENDER PLANT NAMED ‘COCDAP’**

(56) **References Cited**

(50) Latin Name: *Lavandula stoechas*
Varietal Denomination: **Cocdap**

U.S. PATENT DOCUMENTS

PP14,205 P2 * 10/2003 Cherry Plt./226

(76) Inventor: **John Robb**, RMB 2117 Greta Road,
Kulnurra NSW 2250 (AU)

* cited by examiner

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 47 days.

Primary Examiner—Kent Bell
Assistant Examiner—W. C. Haas
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(21) Appl. No.: **11/113,391**

(57) **ABSTRACT**

(22) Filed: **Apr. 23, 2005**

A new and distinct cultivar of Lavender plant named
‘Cocdap’, characterized by its compact, upright, outwardly
spreading and mounded plant habit; freely branching habit,
dense and bushy plant form; early flowering habit; and
violet-colored flowers with pink-colored terminal flower
bracts.

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./226**

(58) **Field of Classification Search** Plt./226
See application file for complete search history.

1 Drawing Sheet

1

2

Botanical designation: *Lavandula stoechas*.
Cultivar denomination: ‘Cocdap’.

SUMMARY OF THE INVENTION

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application is co-pending with the following
related application: U.S. Plant patent application Ser. No.
11/113,400, Lavender Plant Named ‘Coccap’.

Plants of the cultivar ‘Cocdap’ have not been observed
under all possible environmental conditions. The phenotype
may vary somewhat with variations in environment such as
temperature and light intensity without, however, any vari-
ance in genotype.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of Lavender plant, botanically known as *Lavandula*
stoechas, and hereinafter referred to by the name ‘Cocdap’.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Cocdap’.
These characteristics in combination distinguish ‘Cocdap’ as
a new and distinct cultivar:

The new Lavender is a product of a planned breeding
program conducted by the Inventor in Kulnura, New South
Wales, Australia. The objective of the breeding program was
to create new compact, mounded and freely branching
Lavender cultivars with attractive flowers and good garden
performance.

1. Compact, upright, outwardly spreading and mounded
plant habit.
2. Freely branching habit, dense and bushy plant form.
3. Early flowering habit.
4. Violet-colored flowers with pink-colored terminal
flower bracts.

The new Lavender originated from an open-pollination in
1999 of a proprietary selection of *Lavandula stoechas*
identified as code number 00-37, not patented, as the female,
or seed, parent with an unknown selection of *Lavandula*
stoechas, as the male, or pollen, parent. The new Lavender
was discovered and selected by the Inventor as a single
flowering plant within the progeny of the stated open-
pollination grown in a controlled environment in Kulnura,
New South Wales, Australia in September, 2000.

Plants of the new Lavender differ from plants of the
female parent selection in the following characteristics:

1. Plants of the new Lavender are more uniform in plant
habit than plants of the female parent selection.
2. Plants of the new Lavender and the female parent
selection differ in flower coloration.
3. Plants of the new Lavender and the female parent
selection differ in flower bract coloration.

Asexual reproduction of the new cultivar by terminal
cuttings at Kulnura, New South Wales, Australia, since
2000, has shown that the unique features of this new
Lavender are stable and reproduced true to type in succes-
sive generations.

Plants of the new Lavender differ primarily from plants of
the cultivar ‘Coccap’, disclosed in U.S. Plant patent appli-
cation Ser. No. 11/113,400, in flower and terminal flower
bract coloration.

Plants of the new Lavender can be compared to plants of
the Lavender cultivar Kew Red, not patented. In side-by-
side comparisons conducted in Kulnura, New South Wales,
Australia, plants of the new Lavender differed from plants of
the cultivar Kew Red primarily in flower and terminal bract
coloration, inflorescence size, terminal flower bract size and
peduncle length.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the
overall appearance of the new cultivar, showing the colors as

true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the actual colors of the new Lavender.

The photograph at the top of the sheet comprises a side perspective view of a typical flowering plant of 'Cocdap', grown in a container.

The photograph at the bottom of the sheet comprises a close-up view of typical inflorescences of 'Cocdap'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1999 Edition, except where general terms of ordinary dictionary significance are used. Plants used for the aforementioned photographs and following description were grown under conditions which closely approximate commercial production conditions during the spring in a glass-covered greenhouse in Kulnura, New South Wales, Australia for about four to six months in 15-cm containers.

Botanical classification: *Lavandula stoechas* cultivar 'Cocdap'.

Parentage:

Female, or seed, parent.—Proprietary selection of *Lavandula stoechas* identified as code number 00-37, not patented.

Male, or pollen, parent.—Unknown selection of *Lavandula stoechas*, not patented.

Propagation:

Type cutting.—Terminal vegetative cuttings.

Time to initiate roots, summer.—About one week at 25° C.

Time to initiate roots, winter.—About two weeks at 20° C.

Time to produce a rooted young plant, summer.—About six weeks at 25° C.

Time to produce a rooted young plant, winter.—About seven weeks at 20° C.

Root description.—Fine, fibrous.

Rooting habit.—Freely branching.

Plant description:

Form.—Perennial plant; compact, upright, outwardly spreading and mounded plant form. Freely branching habit with lateral branches potentially at every node; dense and bushy plant habit; vigorous growth habit. Flowers in verticillasters on crowded spikes with showy terminal flower bracts.

Plant height.—About 70 cm.

Plant width.—About 70 cm.

Lateral branch description.—Length: About 15 cm to 20 cm. Diameter: About 2 mm. Internode length: About 1.5 cm to 2 cm. Strength: Strong. Aspect: Mostly upright to outwardly spreading. Texture, immature: Pubescent. Texture, mature: Woody. Color, immature: 144C. Color, mature: 141C.

Foliage description.—Arrangement: Opposite, simple; sessile. Length: About 2 cm to 2.5 cm. Width: About 3 mm to 4 mm. Shape: Linear. Apex: Mucronate. Base: Attenuate, clasping. Margin: Entire. Texture, upper and lower surfaces: Pubescent. Fragrance: Very aromatic, pungent. Venation pattern: Pinnate; reticulate. Color: Developing and fully expanded foliage, upper surface: 137A. Developing and fully expanded foliage, lower surface: 137C. Venation, upper and lower surfaces: 137A.

Flower description:

Flower arrangement and shape.—Small single flowers in compact verticillasters on crowded spikes. Freely flowering, about four to six whorls each with about eight to twelve open flowers and flower buds per spike; flowers tubular with five lobes; inflorescences with showy terminal bracts.

Natural flowering season.—Continuous throughout the Spring.

Time to flowering.—Early flowering, plants begin to flower about three months after planting.

Flower longevity on the plant.—Individual inflorescences last about two weeks on the plant and individual flowers last about three days on the plant. Flowers, not persistent; terminal flower bracts, persistent.

Flower buds.—Length: About 4 mm. Diameter: About 1 mm to 2 mm. Shape: Linear to ovoid. Color: 87B.

Inflorescence size.—Height: About 4 cm to 5 cm. Diameter: About 1.5 cm.

Flowers.—Diameter: About 4 mm. Depth (height): About 5 mm.

Petals.—Arrangement: Five, fused into a tube. Length, lobes: About 1 mm. Width, lobes: About 1 mm. Tube length: About 2 mm. Shape: Roughly spatulate. Apex: Rounded. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; velvety. Color: When opening, upper and lower surfaces: 87B. Fully opened, upper and lower surfaces: 87B.

Terminal flower bracts.—Arrangement: About four in a single whorl at inflorescence apex. Length: About 4 cm to 5 cm. Width: About 1.1 cm to 1.5 cm. Shape: Ligulate. Apex: Mucronate. Base: Obtuse. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; velvety. Color, immature, upper and lower surfaces: 80C. Color, mature, upper and lower surfaces: 80C; venation, 81B.

Flower bracts.—Arrangement: Each whorl of flowers subtended by a flower bract. Length: About 6 mm. Width: About 8 mm. Shape: Broadly ovate. Apex: Mucronate. Base: Obtuse. Margin: Entire. Texture, upper and lower surfaces: Smooth. Color, upper and lower surfaces: 147C; venation, 147C.

Calyx.—Arrangement: Five sepals fused into a tube. Length: About 3 mm. Width: About 5 mm. Sepal apex: Acute. Color, immature and mature, upper and lower surfaces: Close to 138B.

Peduncle.—Strength: Strong. Length: About 6 cm to 7 cm. Diameter: About 2 mm. Aspect: Mostly upright. Color: 134A.

Reproductive organs.—Stamens: Quantity per flower: Four. Anther shape: Oval. Anther length: Less than 1 mm. Anther color: Yellow. Pollen amount: Scarce. Pollen color: Yellow. Pistils: Quantity per flower: One. Pistil length: About 5 mm. Stigma shape: Rounded. Stigma color: White. Style length: About 4 mm. Style color: White. Ovary color: Close to 144B.

Seed/fruit.—Seed and fruit production has not been observed.

Disease/pest resistance: Plants of the new Lavender have not been noted to be resistant to pathogens and pests common to Lavender.

Weather tolerance: Plants of the new Lavender have exhibited good tolerance to rain and wind and have been observed to tolerate temperatures from -2° C. to 40° C. It is claimed:

1. A new and distinct cultivar of Lavender plant named 'Cocdap', as illustrated and described.

* * * * *

