

**(12) United States Plant Patent**
Vandenberg**(10) Patent No.: US PP16,813 P2****(45) Date of Patent: Jul. 11, 2006****(54) CHRYSANTHEMUM PLANT NAMED**
'YOLACE'**(50) Latin Name: *Chrysanthemum*×*morifolium***
Varietal Denomination: Yolace**(75) Inventor: Cornelis P. Vandenberg, Fort Myers,**
FL (US)**(73) Assignee: Yoder Brothers Inc., Barberton, OH**
(US)**(*) Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 5 days.**(21) Appl. No.: 11/122,815****(22) Filed: May 5, 2005****(51) Int. Cl.**
A01H 5/00 (2006.01)**(52) U.S. Cl. Plt./288****(58) Field of Classification Search Plt./288**
See application file for complete search history.*Primary Examiner*—Kent Bell*Assistant Examiner*—Annette H Para**(74) Attorney, Agent, or Firm**—C. A. Whealy**(57) ABSTRACT**

A new and distinct cultivar of *Chrysanthemum* plant named 'Yolace', characterized by its upright cut *Chrysanthemum* that is usually grown as a natural spray; dark green-colored foliage; freely flowering habit; uniform inflorescence form and development; decorative-type inflorescences; quill to spoon-shaped and white-colored ray florets; response time about eleven weeks; strong peduncles; and good postproduction longevity.

2 Drawing Sheets**1**Botanical designation: *Chrysanthemum*×*morifolium*.
Cultivar denomination: 'Yolace'.**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of *Chrysanthemum* plant, botanically known as *Chrysanthemum*×*morifolium* and hereinafter referred to by the name 'Yolace'.

The new *Chrysanthemum* is a product of a planned breeding program conducted by the Inventor in Salinas, Calif. and Alva, Fla. The objective of the breeding program is to create new cut *Chrysanthemum* cultivars having inflorescences with desirable colors and good inflorescence form and substance.

The new *Chrysanthemum* originated from a cross-pollination made by the Inventor in December, 1999, in Salinas, Calif., of the *Chrysanthemum*×*morifolium* cultivar Watch, disclosed in U.S. Plant Pat. No. 13,463, as the female, or seed, parent with a proprietary selection of *Chrysanthemum*×*morifolium* identified as code number E620, not patented, as the male, or pollen, parent.

The cultivar Yolace was discovered and selected by the Inventor as a flowering plant within the progeny of the stated cross-pollination in a controlled environment in Alva, Fla., in March, 2001. The selection of this plant was based on its desirable inflorescence color and good inflorescence form and substance.

Asexual reproduction of the new *Chrysanthemum* by terminal cuttings in a controlled environment in Alva, Fla. since May, 2001, has shown that the unique features of this new *Chrysanthemum* are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the cultivar Yolace have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as

2

temperature, daylength and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Yolace'. These characteristics in combination distinguish 'Yolace' as a new and distinct cultivar:

1. Upright cut *Chrysanthemum* that is usually grown as a natural spray.
2. Dark green-colored foliage.
3. Freely flowering habit, about nine to ten inflorescences per flowering stem.
4. Uniform inflorescence form and development.
5. Decorative-type inflorescences that are about 8.3 cm in diameter.
6. Attractive quill or spoon-shaped and white-colored ray florets.
7. Response time about 11.5 weeks.
8. Strong peduncles.
9. Good postproduction longevity with inflorescences and foliage maintaining good substance and color for about two weeks in an interior environment.

Compared to plants of the female parent, the cultivar Watch, plants of the new *Chrysanthemum* have smaller inflorescences. In addition, plants of the new *Chrysanthemum* and the cultivar Watch differ in ray floret coloration as plants of the cultivar Watch have yellow-colored ray florets.

Compared to plants of the male parent selection, plants of the new *Chrysanthemum* have smaller inflorescences. In addition, plants of the new *Chrysanthemum* typically produce few to no disc florets whereas plants of the male parent selection produce numerous disc florets.

Plants of the new *Chrysanthemum* can be compared to plants of the *Chrysanthemum*×*morifolium* cultivar Starburst, not patented. In side-by-side comparisons conducted in Alva, Fla., plants of the new *Chrysanthemum* differed from plants of the cultivar Starburst in the following characteristics:

1. Plants of the new *Chrysanthemum* had stronger flowering stems than plants of the cultivar Starburst.
2. Plants of the new *Chrysanthemum* flowered about one week earlier than plants of the cultivar Starburst.
3. Plants of the new *Chrysanthemum* were more freely flowering than plants of the cultivar Starburst.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Chrysanthemum*, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Chrysanthemum*.

The photograph on the first sheet comprises a side perspective view of a typical flowering stem of 'Yolace' grown as a natural spray.

The photograph on the second sheet comprises a close-up view of typical inflorescences of 'Yolace'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. The aforementioned photographs and following observations and measurements describe plants grown in Salinas, Calif., under conditions which approximate commercial practice in a polyethylene-covered greenhouse. Two-week old rooted cuttings were planted on Nov. 3, 2004 and received four weeks of long day/short nights followed by short day/long nights until flowering. Plants were grown as single-stem natural spray cut *Chrysanthemums*. During the production time, the following environmental conditions were measured: day temperatures, 24 to 27° C.; night temperatures, 10 to 16° C.; and light levels, 2,000 to 4,000 foot-candles. Measurements and numerical values represent averages for six to ten typical flowering stems and were taken about twelve weeks after the start of short days.

Botanical classification: *Chrysanthemum* × *morifolium* cultivar Yolace.

Commercial classification: Decorative-type cut *Chrysanthemum*.

Parentage:

Female or seed parent.—*Chrysanthemum* × *morifolium* cultivar Watch, disclosed in U.S. Plant Pat. No. 13,463.

Male or pollen parent.—Proprietary selection of *Chrysanthemum* × *morifolium* identified as code number E620, not patented.

Propagation:

Type.—Terminal tip cuttings.

Time to rooting.—About 10 to 14 days with soil temperatures of 18 to 21° C.

Root description.—Fine, fibrous; white in color.

Rooting habit.—Freely branching.

Plant description:

Appearance.—Herbaceous decorative-type cut flower that is typically grown as a natural spray.

Flowering stem description.—Aspect: Erect. Length: About 107 cm. Spray diameter: About 17 cm. Stem diameter: About 7.5 mm. Internode length: About 3.7

cm. Texture: Pubescent; longitudinally ridged. Color: Close to 146A.

Foliage description.—Arrangement: Alternate. Length: About 8.1 cm. Width: About 5.2 cm. Apex: Cuspidate. Base: Attenuate to truncate. Margin: Palmately lobed; sinuses mostly divergent. Texture: Upper and lower surfaces, pubescent and leathery; veins prominent on lower surface. Color: Developing and fully expanded foliage, upper surface: Close to 147A. Developing and fully expanded foliage, lower surface: Close to 147B. Venation, upper surface: Close to 147A. Venation, lower surface: Close to 147B to 146A. Petiole: Length: About 2.1 cm. Diameter: About 3.5 mm. Texture, upper and lower surfaces: Pubescent. Color, upper and lower surfaces: 146A to 146B.

Flowering description:

Appearance.—Decorative-type inflorescence form with elongated oblong-shaped ray florets. Inflorescences borne on terminals, arising from leaf axils. Ray florets develop acropetally on a capitulum.

Flowering response.—Under natural conditions, plant flower in the autumn/winter in the Northern Hemisphere. At other times of the year, inflorescence initiation and development can be induced under short day/long night conditions (at least 13.5 hours of darkness). Plants exposed to three to four weeks of long day/short night conditions after planting followed by photoinductive short day/long night conditions flower about 11.5 weeks later when grown as a natural spray. Inflorescences uniform in form and development.

Postproduction longevity.—In an interior environment, inflorescences and foliage will maintain good color and substance for about two weeks in an interior environment.

Quantity of inflorescences.—Freely flowering habit, about nine to ten inflorescences per stem develop.

Fragrance.—None detected.

Inflorescence size.—Diameter: About 7.3 cm. Depth (height): About 2.5 cm. Disc diameter: No disc florets observed. Receptacle diameter: About 6.5 mm. Receptacle height: About 7 mm.

Inflorescence buds.—Height: About 6 mm. Diameter: About 7 mm. Shape: Oblate. Color: Close to 146A to 147A.

Ray florets.—Shape: Quilled or spoon-shaped. Length: About 3.6 cm. Width, apex: About 4.5 mm. Width, tube: About 3 mm. Corolla tube length: About 2.75 cm. Apex: Emarginate, retuse or acute. Base: Fused. Texture: Smooth, glabrous; satiny; longitudinally ridged. Aspect: Initially erect; when mature, mostly straight and about 45° from vertical, eventually perpendicular to the peduncle. Number of ray florets per inflorescence: About 225 in numerous whorls. Color: When opening, upper and lower surfaces: Close to 144A to 144B. Fully opened, upper and lower surfaces: Close to 155D.

Phyllaries.—Quantity per inflorescence: About 26 in about two or three whorls. Length: About 8 mm. Width: About 3.5 mm. Shape: Lanceolate. Apex: Acute. Base: Truncate. Margin: Entire. Texture, upper surface: Smooth, waxy. Texture, lower surface: Pubescent. Color, upper surface: Close to 146A. Color, lower surface: Close to 146A to 147A.

Peduncles.—Length: First peduncle: About 13 cm. Fourth peduncle: About 19.51 cm. Seventh

5

peduncle: About 27.25 cm. Diameter: About 2 mm.
Angle: About 35 to 40° from vertical. Strength:
Strong. Texture: Pubescent. Color: Close to 146A.

Reproductive organs.—Gynoecium: Style length:
About 5 mm. Style color: Close to 144B to 144C.
Stigma color: Close to 9A.

Seed/fruit.—Seed and fruit production has not been
observed.

Disease/pest resistance: Resistance to pathogens and pests
common to *Chrysanthemums* has not been observed on
plants grown under commercial conditions.

6

Temperature tolerance: Plants of the new *Chrysanthemum*
have demonstrated good tolerance to low temperatures of
7° C. and high temperatures high temperatures of 38° C.

It is claimed:

1. A new and distinct cultivar of *Chrysanthemum* plant
named 'Yolace', as illustrated and described.

* * * * *



