



US00PP16859P3

(12) **United States Plant Patent**  
**Jamieson**

(10) **Patent No.:** **US PP16,859 P3**

(45) **Date of Patent:** **Jul. 25, 2006**

(54) **STRAWBERRY VARIETY NAMED**  
**'BRUNSWICK'**

(50) Latin Name: *Fragaria ananassa*  
Varietal Denomination: **Brunswick**

(75) Inventor: **Andrew R. Jamieson**, Nova Scotia  
(CA)

(73) Assignee: **Her Majesty the Queen in Right of**  
**Canada, as represented by the**  
**Minister of Agriculture and Agrifood,**  
**Atlantic Food and Horticulture**  
**Research Centre, Agriculture and**  
**Agri-Food Canada**, Nova Scotia (CA)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 65 days.

(21) Appl. No.: **10/770,360**

(22) Filed: **Feb. 2, 2004**

(65) **Prior Publication Data**

US 2004/0231019 P1 Nov. 18, 2004

**Related U.S. Application Data**

(63) Continuation of application No. 10/073,438, filed on Feb.  
11, 2002, now abandoned.

(51) **Int. Cl.**  
*A01H 5/00* (2006.01)

(52) **U.S. Cl.** ..... **Plt./208**

(58) **Field of Classification Search** ..... **Plt./208**  
See application file for complete search history.

(56) **References Cited**

**PUBLICATIONS**

UPOV-ROM GTITM Computer Database 2004/04, GTI  
Jouve Retrieval Software, Citation for 'Brunswick'.\*

\* cited by examiner

*Primary Examiner*—Anne Marie Grunberg

*Assistant Examiner*—Annette H Para

(74) *Attorney, Agent, or Firm*—The Webb Law Firm

(57) **ABSTRACT**

This invention relates to a new and distinct variety of  
strawberry (*Fragaria ananassa*) named 'Brunswick' char-  
acterized by its large petals, large calyx, ovoid fruit, and  
resistance to powdery mildew. The invention is distinct from  
other varieties, but most similar to 'Cavendish' and 'Hone-  
oye'.

**2 Drawing Sheets**

**1**

Botanical classification: *Fragaria ananassa*.  
Varietal denomination: 'Brunswick'.

**BACKGROUND OF THE INVENTION**

The present invention includes a new and distinct cultivar  
of *Fragaria ananassa* known by the varietal name  
'Brunswick', originally designated as "K90-12". The new  
variety resulted from a controlled cross in an ongoing  
breeding program between the strawberry plants 'Cavend-  
ish' (U.S. Plant Pat. No. 11,110) and 'Honeoye'  
(unpatented). 'Cavendish' was developed by the Atlantic  
Food and Horticulture Research Centre (a facility of Agri-  
culture and Agri-Food Canada) in Kentville and 'Honeoye'  
was developed by the New York State Agricultural Experi-  
ment Station in Geneva. 'Brunswick' was discovered in  
1990 as a seedling in a controlled breeding plot near  
Sheffield Mills, Nova Scotia at the Sheffield Farm, a field-  
station of the Atlantic Food and Horticulture Research  
Centre, where it was selected and propagated asexually by  
stolons at the Atlantic Food and Horticulture Research  
Centre in Kentville. Asexual propagules from this original  
source have been produced annually in a greenhouse at the  
Atlantic Food and Horticulture Research Centre, Kentville,  
Canada. 'Brunswick' has been tested at the Atlantic Food  
and Horticulture Research Centre (starting in 1991) and also  
at research centres in Charlottetown, Prince Edward Island,  
Buctouche, New Brunswick, Fredericton, New Brunswick,  
and Pynn's Brook, Newfoundland, all of Canada. This  
propagation and testing has demonstrated that the combina-  
tion of traits disclosed herein which characterize the new

**2**

variety are fixed and retained true to type through successive  
generations of asexual reproduction via stolons.

**DESCRIPTION OF THE DRAWINGS**

5 FIG. 1 shows plant parts of the new variety, typical in size,  
shape, and color; and

FIG. 2 shows the flowers of the new variety.

**DESCRIPTION OF THE PLANT**

10 The following detailed botanical description of the new  
variety is based upon measurements and observations taken  
of plants and fruit grown in Kentville, Nova Scotia, Canada.  
15 Observations were taken from each variety as grown in a  
side-by-side field trial. Color designations, color  
descriptions, and other phenotypical descriptions may devi-  
ate from the stated values and description depending upon  
variation in the environment, seasonal, climatic and cultural  
20 conditions, however, it is believed that this description will  
apply to the 'Brunswick' plants grown in similar conditions  
of soil and climate elsewhere. Color references generally are  
made to R.H.S. Colour Chart of The Royal Horticultural  
Society of London (1966 edition). Descriptive information  
25 on the new variety is presented in Tables 1 and 2. In the  
tables, the flowers described are secondary flowers. The fruit  
described is the secondary fruit of the maiden crop, thirteen  
or fourteen months after planting. The harvest data in Table  
3 is based on a complete harvest of the crop. Principal  
30 differences between 'Brunswick', the patented variety 'Cav-  
endish' and the unpatented variety 'Honeoye' are set forth

herein. Further, several differences between 'Brunswick' and 'Cavendish' and 'Honeoye' include:

1. 'Brunswick' has more dense petiole pubescence;
2. 'Brunswick' has a later beginning of the flowering period;
3. 'Brunswick' produces more flowers per truss and has a longer flower truss;
4. 'Brunswick' has broader fruit;
5. 'Brunswick's' petals are longer than they are wide, whereas the petals of the reference varieties are as long as they are wide; and
6. 'Brunswick's' inner calyx is smaller than the outer calyx, whereas the inner and outer calyces of the reference varieties have the same diameter.

Classification: The new variety is botanically identified as *Fragaria ananassa* and commercially classified as a short-day strawberry.

Plant and foliage characteristics: When propagated in the nursery, 'Brunswick' produces moderate numbers of runner plants, similar to 'Cavendish' but less than 'Honeoye'. Individual plants of all three varieties are medium in size. The plants of 'Brunswick' are of medium density with globose habit and strong vigor. 'Brunswick' rows are less dense and more open than rows of 'Honeoye', but more dense than 'Cavendish'. As shown in Table 1, leaf color of 'Brunswick', 'Cavendish' and 'Honeoye' are Green Group 137A on the upper surface. The color of the lower surface of 'Brunswick' and 'Honeoye' is Green Group 138B but Green Group 137C for 'Cavendish'. Leaflets of 'Brunswick' are larger than 'Cavendish' and slightly larger than 'Honeoye' and leaves of all three varieties have three leaflets. Leaflets of 'Brunswick' and 'Cavendish' are cupped whereas 'Honeoye' leaflets are flat. Leaflets of 'Brunswick' and 'Cavendish' have higher numbers of serrations than 'Honeoye'. Leaflet serrations of 'Brunswick' and 'Honeoye' are semi-round whereas serrations of 'Cavendish' are rounded. The venation of 'Brunswick' leaflets is pinnate. Petiole pubescence for 'Brunswick' tends to be higher in density than for 'Cavendish' and 'Honeoye' but all three varieties have a perpendicular orientation of the pubescence.

TABLE 1

Foliar Characteristics for 'Brunswick', 'Cavendish' and 'Honeoye'			
Foliar Character	Cultivar		
	'Brunswick'	'Cavendish'	'Honeoye'
Leaf color	Green Group	Green Group	Green Group
Upper surface	137A	137A	137A
Lower surface	138B	137C	138B
Central leaflet			
<u>Length (mm)</u>			
Mean	87.8	79.9	83.8
Range	80–100	64–89	65–97
<u>Width (mm)</u>			
Mean	77.0	74.6	70.3
Range	65–95	55–90	55–87
Length/width ratio	1.14	1.07	1.19
No. leaflets/leaf	3	3	3
Leaf convexity	cupped	cupped	flat

TABLE 1-continued

Foliar Characteristics for 'Brunswick', 'Cavendish' and 'Honeoye'			
Foliar Character	Cultivar		
	'Brunswick'	'Cavendish'	'Honeoye'
<u>Serrations</u>			
Number	many	many	moderate
Size	medium	medium	small
Shape	semi-round	rounded	semi-round
Tip serration size	small	small	small
Leaf pubescence	medium	medium	medium
<u>Petiole pubescence</u>			
Density	medium to dense	medium	medium
Direction	perpendicular	perpendicular	perpendicular

Flower and fruit characteristics: The length of bloom for 'Brunswick' is about three weeks when grown in Kentville, Nova Scotia, Canada in a matted row cultural system. 'Brunswick' typically has 2 to 4 crowns when counted at bloom time of the first harvest season. Flowering for 'Brunswick' typically begins on June 2 and ends on June 21, which is about three days later than 'Cavendish' and four days later than 'Honeoye'. As shown in Table 2, the flower truss of 'Brunswick' is longer than for 'Cavendish' and 'Honeoye'. 'Brunswick' flowers typically open slightly beneath the leaf canopy. 'Brunswick' flower trusses produce more flowers than trusses of 'Cavendish' and 'Honeoye'. Flowers of 'Brunswick' and the reference varieties are white. The anther color is Yellow-Orange Group 17A. Secondary flowers of 'Brunswick' are similar to 'Honeoye' in size but larger than 'Cavendish'. Secondary flowers of 'Brunswick' typically have 5 or 6 petals while 'Cavendish' usually has 6 petals and 'Honeoye' has 5. The petals of 'Brunswick' are slightly longer than wide whereas petals of 'Cavendish' and 'Honeoye' are as long as wide. The inner calyx of 'Brunswick' has a smaller diameter than the outer calyx while for 'Cavendish' and 'Honeoye' the inner and outer calyces have a similar diameter. Trusses of 'Brunswick' and 'Honeoye' are typically semi-erect at first picking in contrast to 'Cavendish' which is prostrate. The position of the calyx is even with the top of the berry for 'Brunswick'. The adherence of the calyx to the fruit is weak to medium for 'Brunswick' and 'Honeoye' but strong for 'Cavendish'. The fruit of 'Brunswick' are slightly larger than for 'Honeoye' but smaller than 'Cavendish'. The ratio of fruit length to width is 0.82 for 'Brunswick', 0.92 for 'Cavendish' and 0.97 for 'Honeoye'. 'Brunswick' fruit are ovoid to short-conic in shape. Fruit of all three varieties are medium in firmness. For 'Brunswick', the hollow center size ranges from zero in small tertiary fruit, to 1–3 mm (diameter) in medium-sized tertiary fruit, to 3–6 mm in large primary fruit. As shown in Table 2, the exterior fruit color of 'Brunswick' is Red Group 46A and on the inside the pith and cortex are Red Group 45B, indicating very even interior color. The reference varieties have a very similar color to 'Brunswick'. The achenes of 'Brunswick' are Green-Yellow Group 1A but darken to Orange-Red Group 34A on the shoulders of the fruit when exposed to sunlight.

TABLE 2

Character	Cultivar		
	'Brunswick'	'Cavendish'	'Honeoye'
Flower position (relative to leaf canopy)	slightly beneath	beneath	slightly beneath
Flower truss length (cm)	medium 24.2	short-medium 20.4	short-medium 21.8
Number of flowers/ truss	8.1	6.4	6.7
Number of petals	5.4	6.0	5.1
Flower size (mm diameter)	31.1	27.0	30.1
Flower color	White	White	White
Petal length (mm)	12.2	10.6	12.0
Petal width (mm)	11.6	10.5	12.0
Petal spacing	spaced to touching	touching to overlapping	spaced to touching
<u>Calyx size</u>			
Inner calyx (mm diam.)	24.4	22.8	23.1
Outer calyx (mm diam.)	26.7	22.4	23.8
Calyx position	even	even to in a basin	even to slightly above
Adherence of the calyx	weak to medium	strong	weak to medium
<u>Fruit size and shape</u>			
Length (mm)	28.1	31.8	29.9
Width (mm)	34.4	34.6	30.7
Length/width ratio	0.82	0.92	0.97
Subjective fruit shape	ovoid to short-conic	cordate	short-conic
Seed position	slight-indent	slight-indent	slight indent
Fruit firmness	medium	medium	medium
Color (R.H.S. Colour Chart)			
Calyx	Green Group 137C to 137D	Green Group 137C	Green Group 137C
Fruit exterior	Red Group 46A	Red Group 46A	Red Group 46A
Fruit interior	Red Group	Red Group	Red Group
Pith	45B	45B	45B
Cortex	45B	45B	45B

Disease resistance: 'Brunswick' and 'Cavendish' are resistant to red stele root rot (*Phytophthora fragariae*) while 'Honeoye' is susceptible. 'Brunswick' plants are slightly more resistant to powdery mildew (*Sphaerotheca macularis*) than 'Cavendish' and 'Honeoye', which are susceptible. 'Brunswick' and 'Cavendish' are susceptible to green petal phytoplasma while 'Honeoye' is resistant.

Production characteristics: 'Brunswick' has produced high yields; generally similar to 'Cavendish' and 'Honeoye', as shown in Table 3. The proportion of fruit considered marketable is similar between the three varieties. The fruit of 'Brunswick' are larger (by weight) than 'Honeoye' but smaller than 'Cavendish'. The mean harvest date for 'Brunswick' is similar to 'Cavendish' and 'Honeoye'. All three varieties have a harvest season which is early-midseason.

TABLE 3

Performance of 'Brunswick' and 'Cavendish' or 'Honeoye' in 1997 and 1998 Averaged over Several Sites*				
	Total Yield (t/ha)	% Yield marketable	Size (g/fruit)	Mean Harvest (day of year)
<u>1997 (five sites)</u>				
'Brunswick'	7.2	87.9	13.3	201.7
'Cavendish'	5.4	90.8	14.7	201.5
<u>1998 (four sites)</u>				
'Brunswick'	9.9	81.1	11.2	188.2
'Cavendish'	8.6	81.2	11.6	189.2
<u>1998 (one site)</u>				
'Brunswick'	17.1	94.2	12.8	181.0
'Honeoye'	15.8	96.1	10.8	182.8

\*Kentville NS, Charlottetown PEI, Fredericton NB, Buctouche NB, and Pynn's Brook Nfld. Plants were grown in matted rows and three blocks of 3 m long rows were harvested at each site.

I claim:

1. The new and distinct variety of strawberry plant named 'Brunswick' as described and illustrated.

\* \* \* \* \*




Fig. 1




Fig. 2

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : PP 16,859 P3  
APPLICATION NO. : 10/770360  
DATED : July 25, 2006  
INVENTOR(S) : Jamieson

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Title page, Item (73) Assignee, fourth line, "HortIculutre" should read -- Horticulture --

Signed and Sealed this

Twelfth Day of December, 2006

A handwritten signature in black ink on a light gray dotted background. The signature reads "Jon W. Dudas" in a cursive style.

JON W. DUDAS

*Director of the United States Patent and Trademark Office*