



(12) **United States Plant Patent**
Smith

(10) **Patent No.:** **US PP16,969 P2**
(45) **Date of Patent:** **Aug. 8, 2006**

(54) **CHRYSANTHEMUM PLANT NAMED**
‘YOBETHANY’

(50) Latin Name: *Chrysanthemum*×*morifolium*
Varietal Denomination: **Yobethany**

(75) Inventor: **Mark A. Smith**, Fort Myers, FL (US)

(73) Assignee: **Yoder Brothers, Inc.**, Barberton, OH
(US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/212,348**

(22) Filed: **Aug. 26, 2005**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./289**

(58) **Field of Classification Search** **Plt./289**
See application file for complete search history.

Primary Examiner—Anne Marie Grunberg
Assistant Examiner—Georgia Helmer
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Chrysanthemum* plant named
‘Yobethany’, characterized by its upright and somewhat
outwardly spreading plant habit; freely branching habit;
dense and full plant habit; uniform and freely flowering
habit; decorative-type inflorescences with elongated oblong-
shaped ray florets; yellow-colored ray florets; and natural
season flowering in early October in the Northern Hemi-
sphere.

2 Drawing Sheets

1

Botanical designation: *Chrysanthemum*×*morifolium*.
Cultivar denomination: ‘Yobethany’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Chrysanthemum* plant, botanically known as
Chrysanthemum×*morifolium*, commercially known as a
garden-type *Chrysanthemum* and hereinafter referred to by
the name ‘Yobethany’.

The new cultivar is a product of a planned breeding
program conducted by the Inventor in Salinas, Calif. and
Alva, Fla. The objective of the breeding program is to create
new garden-type *Chrysanthemum* cultivars having inflores-
cences with desirable inflorescence forms, attractive floret
colors and good garden performance.

The new *Chrysanthemum* originated from a cross-
pollination made in November, 1999 in Salinas, Calif., of a
proprietary selection of *Chrysanthemum*×*morifolium* iden-
tified as code number 96-L067, not patented, as the female,
or seed, parent with a proprietary selection of
Chrysanthemum×*morifolium* identified as code number
95-L443007, not patented, as the male, or pollen, parent.
The new *Chrysanthemum* was discovered and selected by
the Inventor as a single flowering plant within the progeny
of the stated cross-pollination grown in a controlled envi-
ronment in Alva, Fla. in October, 2001. The selection of this
plant was based on its desirable inflorescence form, attrac-
tive floret coloration and good garden performance.

Asexual reproduction of the new cultivar by terminal
vegetative cuttings in a controlled environment in Alva, Fla.
since January, 2002, has shown that the unique features of
this new *Chrysanthemum* are stable and reproduced true to
type in successive generations.

SUMMARY OF THE INVENTION

The cultivar Yobethany has not been observed under all
possible environmental conditions. The phenotype may vary
somewhat with variations in environment such as

2

temperature, daylength and light intensity, without,
however, any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Yobe-
thany’. These characteristics in combination distinguish
‘Yobethany’ as a new and distinct cultivar:

1. Upright and somewhat outwardly spreading plant habit.
2. Freely branching habit; dense and full plants.
3. Uniform and freely flowering habit.
4. Decorative-type inflorescences with elongated oblong-
shaped ray florets.
5. Yellow-colored ray florets.
6. Natural season flowering in early October in the
Northern Hemisphere.

In side-by-side comparisons conducted in Alva, Fla.,
plants of the new *Chrysanthemum* differed from plants of the
female parent selection in the following characteristics:

1. Plants of the new *Chrysanthemum* were slightly larger
than plants of the female parent selection.
2. Plants of the new *Chrysanthemum* and the female
parent selection differed in ray floret coloration as
plants of the female parent selection had white-colored
ray florets.
3. Plants of the new *Chrysanthemum* flowered about two
weeks later than plants of the female parent selection
when grown under natural season conditions.

In side-by-side comparisons conducted in Alva, Fla.,
plants of the new *Chrysanthemum* differed from plants of the
male parent selection in the following characteristics:

1. Plants of the new *Chrysanthemum* were stronger and
more uniform than plants of the male parent selection.
2. Plants of the new *Chrysanthemum* flowered more
uniformly than plants of the male parent selection.
3. Plants of the new *Chrysanthemum* had smaller inflo-
rescences than plants of the male parent selection.
4. Plants of the new *Chrysanthemum* had fewer disc
florets per inflorescence than plants of the male parent
selection.

- Plants of the new *Chrysanthemum* flowered about one week later than plants of the male parent selection when grown under natural season conditions.

Plants of the new *Chrysanthemum* can be compared to plants of the *Chrysanthemum* cultivar Sunny Denise, disclosed in U.S. Plant Pat. No. 8,980. In side-by-side comparisons conducted in Alva, Fla., plants of the new *Chrysanthemum* differed from plants of the cultivar Sunny Denise in the following characteristics:

- Plants of the new *Chrysanthemum* were smaller and more rounded than and not as upright as plants of the cultivar Sunny Denise.
- Plants of the new *Chrysanthemum* were more freely flowering than plants of the cultivar Sunny Denise.
- Plants of the new *Chrysanthemum* had smaller inflorescences than plants of the cultivar Sunny Denise.
- Ray florets of plants of the new *Chrysanthemum* were darker yellow in color than ray florets of plants of the cultivar Sunny Denise.

Plants of the new *Chrysanthemum* can also be compared to plants of the *Chrysanthemum* cultivar Cleagar gfl disclosed in U.S. Plant Pat. No. 10,463. In side-by-side comparisons conducted in Alva, Fla., plants of the new *Chrysanthemum* differed from plants of the cultivar Cleagar gfl in the following characteristics:

- Plants of the new *Chrysanthemum* were larger and more rounded than plants of the cultivar Cleagar gfl.
- Plants of the *Chrysanthemum* had better garden performance and were durable than plants of the cultivar Cleagar gfl.
- Plants of the new *Chrysanthemum* flowered more uniformly than plants of the cultivar Cleagar gfl.
- Plants of the new *Chrysanthemum* had smaller inflorescences than plants of the cultivar Cleagar gfl.
- Ray florets of plants of the new *Chrysanthemum* were brighter yellow in color than ray florets of plants of the cultivar Cleagar gfl.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall appearance of the new *Chrysanthemum*. These photographs show the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Chrysanthemum*.

The photograph on the first sheet comprises a side perspective view of a typical flowering plant of 'Yobethany' grown in a container.

The photograph on the second sheet comprises a close-up view of typical inflorescences of the cultivar 'Yobethany'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. The following observations and measurements describe plants grown in Leamington, Ontario, Canada during the summer in a glass-covered greenhouse and under conditions and practices which approximate those generally used in commercial garden-type *Chrysanthemum* production. Rooted cuttings were planted in 15.25-cm containers, grown under artificial long day conditions (four-

hour night interruption) and pinched about ten days later. About ten days after the pinch, plants were then exposed to artificial short day conditions (11.5 hours light) until flowering. During the production of the plants, temperatures ranged from 18° C. to 38° C. Measurements and numerical values represent averages for typical flowering plants.

Botanical classification: *Chrysanthemum*×*morifolium* cultivar Yobethany.

Commercial classification: Decorative-type garden *Chrysanthemum*.

Parentage:

Female, or seed, parent.—Proprietary selection of *Chrysanthemum*×*morifolium* identified as code number 96-L067, not patented.

Male, or pollen, parent.—Proprietary selection of *Chrysanthemum*×*morifolium* identified as code number 95-L443007, not patented.

Propagation:

Type.—Terminal vegetative cuttings.

Time to initiate roots.—About four days at 21° C.

Time to produce a rooted cutting.—About ten to twelve days at 21° C.

Root description.—Fine, fibrous; white in color.

Rooting habit.—Freely branching.

Plant description:

Plant form/growth habit.—Perennial herbaceous decorative-type garden *Chrysanthemum*. Inverted triangle with mounded crown. Stems initially upright, then somewhat outwardly spreading; rounded growth habit. Freely branching with about seven primary branches with secondary lateral branches potentially forming at every node. Moderately vigorous.

Plant height.—About 18 cm.

Plant diameter.—About 28 cm.

Lateral branches.—Length: About 16.5 cm. Diameter: About 4.5 mm. Internode length: About 1.3 cm. Aspect: Upright and somewhat outwardly spreading. Texture: Pubescent. Color: 146A.

Foliage description.—Leaf arrangement: Alternate. Length: About 4.6 cm. Width: About 2.9 cm. Apex: Cuspidate. Base: Attenuate. Margin: Palmately lobed, sinuses mostly divergent. Texture, upper surface: Slightly pubescent. Texture, lower surface: Pubescent; veins prominent. Color: Developing and fully expanded foliage, upper surface: Darker green than 147A. Developing and fully expanded foliage, lower surface: Close to 147B. Venation, upper surface: Close to 147A. Venation, lower surface: Close to 147B. Petiole length: About 1.3 cm. Petiole diameter: About 3 mm. Petiole color, upper surface: Close to 146A. Petiole color, lower surface: Close to 146B.

Inflorescence description:

Appearance.—Decorative-type inflorescence form with elongated oblong-shaped ray florets. Inflorescences borne on terminals above foliage, arising from leaf axils. Ray florets developing acropetally on a capitulum. About six inflorescences per secondary lateral branch.

Flowering response.—Under natural season conditions, plants flower in early October in the Northern Hemisphere.

Inflorescence bud (before showing color).—Height: About 4 mm. Diameter: About 5 mm. Shape: Oblate.

Color (lower surface of phyllaries): More green than 147A.

Inflorescence size.—Diameter: About 2.5 cm. Depth (height): About 1.1 cm. Disc diameter: About 2 mm; inconspicuous. Receptacle diameter: About 3.5 mm.

Ray florets.—Shape: Elongated oblong-shaped. Length: About 1.2 cm. Width: About 3.5 mm. Corolla tube length: About 2.5 mm. Corolla tube diameter: About 1 mm. Apex: Emarginate or rounded. Margin: Fused. Texture, upper and lower surfaces: Smooth, glabrous; satiny. Orientation: Initially upright, eventually perpendicular to the peduncle. Number of ray florets per inflorescence: About 155 in numerous whorls. Color: When opening and fully opened, upper surface: Close to 6A. When opening and fully opened, lower surface: Close to 6A to 6B.

Disc florets.—Shape: Tubular, elongated. Length: About 3 mm. Width, apex: About 1 mm. Width, base: About 1 mm. Number of disc florets per inflorescence: About five. Color: Immature: Close to 9A. Mature: Apex: Close to 9A. Mid-section: Close to 154D. Base: Close to 155D.

Phyllaries.—Quantity per inflorescence: About 18. Length: About 6 mm. Width: About 2 mm. Shape: Ligulate. Apex: Acute. Base: Truncate. Margin: Entire. Texture, upper surface: Smooth, waxy.

Texture, lower surface: Pubescent. Color, upper surface: Close to 146A. Color, lower surface: More green than 147A.

Peduncle.—Length: First peduncle: About 1.4 cm. Fourth peduncle: About 2 cm. Diameter: About 1.5 mm. Strength: Strong. Aspect: About 45° from vertical. Texture: Pubescent. Color: Close to 146A.

Reproductive organs.—Androecium: Present on disc florets only. Anther length: Less than 1 mm. Anther color: Close to 12A. Amount of pollen: None observed. Gynoecium: Present on both ray and disc florets. Style length: About 4 mm. Style color: Close to 154A. Stigma color: Close to 9A.

Seed/fruit.—Seed and fruit production has not been observed.

Disease/pest resistance: Plants of the new *Chrysanthemum* have not been shown to be resistant to pathogens and pests common to *Chrysanthemums*.

Garden performance: Plants of the new *Chrysanthemum* have been observed to be tolerant to rain, wind and temperatures ranging from 0° C. to more than 38° C.

It is claimed:

1. A new and distinct cultivar of *Chrysanthemum* plant named 'Yobethany', as illustrated and described.

* * * * *



