



US00PP17008P2

(12) **United States Plant Patent**
Smith(10) **Patent No.:** US PP17,008 P2
(45) **Date of Patent:** Aug. 15, 2006

- (54) **CHrysanthemum PLANT NAMED 'YOBARBIE'**
- (50) Latin Name: *Chrysanthemum×morifolium*
Varietal Denomination: Yobarbie
- (75) Inventor: **Mark A. Smith**, Fort Myers, FL (US)
- (73) Assignee: **Yoder Brothers, Inc.**, Barberton, OH (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **11/212,346**
- (22) Filed: **Aug. 26, 2005**
- (51) **Int. Cl.**
A01H 5/00 (2006.01)

- (52) **U.S. Cl.** **Plt./287**
- (58) **Field of Classification Search** Plt./287
See application file for complete search history.

Primary Examiner—Anne Marie Grunberg

Assistant Examiner—Georgia Helmer

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Chrysanthemum* plant named 'Yobarbie', characterized by its compact, upright and somewhat outwardly spreading plant habit; freely branching habit; dense and full plant habit; uniform and freely flowering habit; medium-sized decorative-type inflorescences with elongated oblong-shaped ray florets; lavender-colored ray florets; and natural season flowering in mid-October in the Northern Hemisphere.

2 Drawing Sheets

1

Botanical designation: *Chrysanthemum×morifolium*.
Cultivar denomination: 'Yobarbie'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Chrysanthemum* plant, botanically known as *Chrysanthemum×morifolium*, commercially known as a garden-type *Chrysanthemum* and hereinafter referred to by the name 'Yobarbie'.
The new cultivar is a product of a planned breeding program conducted by the Inventor in Salinas, Calif. and Alva, Fla. The objective of the breeding program is to create new garden-type *Chrysanthemum* cultivars having inflorescences with desirable inflorescence forms, attractive floret colors and good garden performance.

The new *Chrysanthemum* originated from a cross-pollination made in March, 2001 in Salinas, Calif., of a proprietary selection of *Chrysanthemum×morifolium* identified as code number 98-M306, not patented, as the female, or seed, parent with the *Chrysanthemum×morifolium* cultivar Empire Symphony, disclosed in U.S. Plant Pat. No. 8,782, as the male, or pollen, parent. The new *Chrysanthemum* was discovered and selected by the Inventor as a single flowering plant within the progeny of the stated cross-pollination grown in a controlled environment in Alva, Fla. in November, 2002. The selection of this plant was based on its desirable inflorescence form, attractive floret coloration and good garden performance.

Asexual reproduction of the new cultivar by terminal vegetative cuttings in a controlled environment in Alva, Fla. since January, 2003, has shown that the unique features of this new *Chrysanthemum* are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

The cultivar Yobarbie has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as

2

temperature, daylength and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Yobarbie'. These characteristics in combination distinguish 'Yobarbie' as a new and distinct cultivar:

1. Compact, upright and somewhat outwardly spreading plant habit.
2. Freely branching habit; dense and full plants.
3. Uniform and freely flowering habit.
4. Medium-sized decorative-type inflorescences with elongated oblong-shaped ray florets.
5. Lavender-colored ray florets.
6. Natural season flowering in mid-October in the Northern Hemisphere.

In side-by-side comparisons conducted in Alva, Fla., plants of the new *Chrysanthemum* differed from plants of the female parent selection in the following characteristics:

1. Plants of the new *Chrysanthemum* and the female parent selection differed in ray floret coloration as plants of the female parent selection had red bronze-colored ray florets.
2. Ray florets of plants of the new *Chrysanthemum* had better color retention than ray florets of plants of the female parent selection.
3. Plants of the new *Chrysanthemum* flowered about two weeks later than plants of the female parent selection when grown under natural season conditions.

In side-by-side comparisons conducted in Alva, Fla., plants of the new *Chrysanthemum* differed from plants of the male parent, the cultivar Empire Symphony, in the following characteristics:

1. Plants of the new *Chrysanthemum* were more rounded than plants of the cultivar Empire Symphony.
2. Plants of the new *Chrysanthemum* had smaller inflorescences than plants of the cultivar Empire Symphony.

3. Plants of the new *Chrysanthemum* flowered about one month later than plants of the cultivar Empire Symphony when grown under natural season conditions.

Plants of the new *Chrysanthemum* can be compared to plants of the *Chrysanthemum* cultivar Yodanielle, disclosed in U.S. Plant Pat. No. 13,802. In side-by-side comparisons conducted in Alva, Fla., plants of the new *Chrysanthemum* differed from plants of the cultivar Yodanielle in the following characteristics:

1. Plants of the new *Chrysanthemum* were larger than plants of the cultivar Yodanielle.
2. Plants of the new *Chrysanthemum* were less sensitive to root pathogens than plants of the cultivar Yodanielle.
3. Plants of the new *Chrysanthemum* flowered about five days later than plants of the cultivar Yodanielle when grown under natural season conditions.

Plants of the new *Chrysanthemum* can also be compared to plants of the *Chrysanthemum* cultivar Gedi Two Cop, disclosed in U.S. Plant Pat. No. 14,404. In side-by-side comparisons conducted in Alva, Fla., plants of the new *Chrysanthemum* differed from plants of the cultivar Gedi Two Cop in the following characteristics:

1. Plants of the new *Chrysanthemum* were larger than plants of the cultivar Gedi Two Cop.
2. Ray florets of plants of the new *Chrysanthemum* had better color retention than ray florets of plants of the cultivar Gedi Two Cop.
3. Plants of the new *Chrysanthemum* flowered about two to four days later than plants of the cultivar Gedi Two Cop when grown under natural season conditions.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall appearance of the new *Chrysanthemum*. These photographs show the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Chrysanthemum*.

The photograph on the first sheet comprises a side perspective view of a typical flowering plant of 'Yobarbie' grown in a container.

The photograph on the second sheet comprises a close-up view of typical inflorescences of the cultivar 'Yobarbie'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. The following observations and measurements describe plants grown in Leamington, Ontario, Canada during the summer in a glass-covered greenhouse and under conditions and practices which approximate those generally used in commercial garden-type *Chrysanthemum* production. Rooted cuttings were planted in 15.25-cm containers, grown under artificial long day conditions (four-hour night interruption) and pinched about ten days later. About ten days after the pinch, plants were then exposed to artificial short day conditions (11.5 hours light) until flowering. During the production of the plants, temperatures ranged from 18° C. to 38° C. Measurements and numerical values represent averages for typical flowering plants.

Botanical classification: *Chrysanthemum × morifolium* cultivar Yobarbie.

Commercial classification: Decorative-type garden *Chrysanthemum*.

Parentage:

Female, or seed, parent.—Proprietary selection of *Chrysanthemum × morifolium* identified as code number 98-M306, not patented.

Male, or pollen, parent.—*Chrysanthemum × morifolium* cultivar Empire Symphony, disclosed in U.S. Plant Pat. No. 8,782.

Propagation:

Type.—Terminal vegetative cuttings.

Time to initiate roots.—About four days at 21° C.

Time to produce a rooted cutting.—About ten to twelve days at 21° C.

Root description.—Fine, fibrous; white in color.

Rooting habit.—Freely branching.

Plant description:

Plant form/growth habit.—Perennial herbaceous decorative-type garden *Chrysanthemum*. Inverted triangle with rounded crown. Stems initially upright, then somewhat outwardly spreading; compact and rounded growth habit. Freely branching with about five primary branches with secondary lateral branches potentially forming at every node. Moderately vigorous.

Plant height.—About 16.5 cm.

Plant diameter.—About 21 cm.

Lateral branches.—Length: About 14.5 cm. Diameter: About 4 mm. Internode length: About 9 mm. Aspect: Upright to somewhat outwardly spreading. Texture: Pubescent. Color: 144A.

Foliage description.—Leaf arrangement: Alternate. Length: About 4.75 cm. Width: About 4.1 cm. Apex: Cuspidate. Base: Mostly truncate. Margin: Palmately lobed, sinuses parallel to divergent. Texture, upper surface: Pubescent. Texture, lower surface: Pubescent, veins prominent. Color: Developing and fully expanded foliage, upper surface: More green than 147A. Developing and fully expanded foliage, lower surface: Close to 147B. Venation, upper surface: Close to 146A. Venation, lower surface: Close to 147B. Petiole length: About 1.4 cm. Petiole diameter: About 3.5 mm. Petiole color, upper and lower surfaces: Close to 146B.

Inflorescence description:

Appearance.—Decorative-type inflorescence form with elongated oblong-shaped ray florets. Inflorescences borne on terminals above foliage, arising from leaf axils. Ray florets developing acropetally on a capitulum. About five inflorescences per secondary lateral branch.

Flowering response.—Under natural season conditions, plants flower in mid-October in the Northern Hemisphere.

Inflorescence bud (before showing color).—Height: About 5 mm. Diameter: About 7 mm. Shape: Oblate. Color (lower surface of phyllaries): Close to 146A to more green than 147A.

Inflorescence size.—Diameter: About 3.3 cm. Depth (height): About 1.4 cm. Disc diameter: About 2 mm; inconspicuous. Receptacle diameter: About 5 mm.

Ray florets.—Shape: Elongated oblong-shaped. Length: About 1.6 cm. Width: About 4.5 mm. Corolla tube length: About 3 mm. Corolla tube diameter: About 1

mm. Apex: Mostly emarginate. Margin: Fused. Texture, upper and lower surfaces: Smooth, glabrous; satiny. Orientation: Initially upright, then perpendicular to the peduncle. Number of ray florets per inflorescence: About 242 in numerous whorls. Color: When opening and fully opened, upper surface: Close to 155D overlain with close to 77D; color becoming lighter lavender with development. When opening and fully opened, lower surface: Close to 155D underlain with close to 79A; color becoming lighter lavender to white, close to 155D, with development.

Disc florets.—Shape: Tubular, elongated. Length: About 3 mm. Width, apex: About 1 mm. Width, base: About 1 mm. Number of disc florets per inflorescence: About five. Color: Immature: Close to 9A. Mature: Apex: Close to 9A. Mid-section: Close to 154D. Base: Close to 155D.

Phyllaries.—Quantity per inflorescence: About 16. Length: About 6 mm. Width: About 3 mm. Shape: Ligulate. Apex: Acute. Base: Truncate. Margin: Entire. Texture, upper surface: Smooth, waxy. Texture, lower surface: Pubescent. Color, upper sur-

face: Close to 146A. Color, lower surface: Close to 146A to more green than 147A.

Peduncle.—Length: First peduncle: About 2 cm. Fourth peduncle: About 2.4 cm. Diameter: About 2 mm. Strength: Strong. Aspect: About 40° from vertical. Texture: Pubescent. Color: Close to 144A to 146A.

Reproductive organs.—Androecium: Present on disc florets only. Anther length: Less than 1 mm. Anther color: Close to 12A. Amount of pollen: None observed. Gynoecium: Present on both ray and disc florets. Style length: About 4 mm. Style color: Close to 154A. Stigma color: Close to 9A.

Seed/fruit.—Seed and fruit production has not been observed.

Disease/pest resistance: Plants of the new *Chrysanthemum* have not been shown to be resistant to pathogens and pests common to *Chrysanthemums*.

Garden performance: Plants of the new *Chrysanthemum* have been observed to be tolerant to rain, wind and temperatures ranging from 0° C. to more than 38° C.

It is claimed:

1. A new and distinct cultivar of *Chrysanthemum* plant named 'Yobarbie', as illustrated and described.

* * * * *



