



US00PP17104P2

(12) **United States Plant Patent**
Danziger

(10) **Patent No.:** **US PP17,104 P2**
(45) **Date of Patent:** **Sep. 19, 2006**

(54) **CHRYSANTHEMUM PLANT NAMED**
'DANATDARPK'

(50) Latin Name: *Chrysanthemum morifolium*
Varietal Denomination: **Danatdarpink**

(75) Inventor: **Gabriel Danziger**, Moshav Nir-Zvi (IL)

(73) Assignee: **Danziger "Dan" Flower Farm**, Post
Beit Dagan (IL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/202,039**

(22) Filed: **Aug. 12, 2005**

(30) **Foreign Application Priority Data**

Feb. 25, 2005 (CO) A05889

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./293**

(58) **Field of Classification Search** **Plt./292,**
Plt./293

See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

Chan 1966 *Chrysanthemum* and rose mutations induced by
x-rays Am. Soc. Hort. Sci. Proc. pp. 613-620.*

Broertjes 1966 Mutation breeding of *chrysanthemums*
Euphytica 15: 156-162.*

Dowrick et al. 1966, The induction of mutation in *Chrysan-*
themum using x- and gamma radiation Euphytica 15:
204-210.*

* cited by examiner

Primary Examiner—Kent Bell

Assistant Examiner—Annette H Para

(74) *Attorney, Agent, or Firm*—Foley & Lardner LLP

(74) *Attorney, Agent, or Firm*—

(57) **ABSTRACT**

A new and distinct *Chrysanthemum* plant named 'Danat-
darpink' characterized by having inflorescence with oblong
shaped, red-purple ray florets, and bell shaped, yellow disc
florets; flowering time of 7.5 weeks; short day plant; and
vigorous growth habit.

2 Drawing Sheets

1

Botanical designation: *Chrysanthemum morifolium*.
Variety denomination: 'Danatdarpink'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Chrysanthemum* plant, botanically known as *Chrysanthe-*
mum morifolium, hereinafter referred to by the cultivar name
'Danatdarpink'.

The new cultivar 'Danatdarpink' is a product of a planned
mutant selection and monitoring program in Moshav Mish-
mar Hashiva, Israel, with the objective of finding color
mutants of the *Chrysanthemum morifolium* cultivar 'Dan-
chryatpink'. The new cultivar 'Danatdarpink' originated
from an induced, whole plant mutation on a single flowering
plant of the *Chrysanthemum morifolium* cultivar 'Dan-
atchryatpink' (unpatented, disclosed in U.S. patent applica-
tion Ser. No. 10/825,115) and grown in a controlled, culti-
vated environment in Moshav Mishmar Hashiva, Israel. The
parental cultivar 'Danchryatpink' is also known by the
commercial variety name Atlantis Pink.

The new cultivar 'Danatdarpink' was discovered and
selected by the inventor, Gabriel Danziger, as an induced,
whole mutation plant of 'Danchryatpink' in January of 2003.
The mutation arose by radiation.

Asexual reproduction of the new cultivar by stem cutting
was first performed in January of 2004 in Moshav Mishmar
Hashiva, Israel, and has demonstrated that the combination
of characteristics as herein disclosed for the new cultivar are
firmly fixed and retained through successive generations of

2

asexual reproduction. The new cultivar reproduces true to
type.

BRIEF SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and
are determined to be unique characteristics of 'Danat-
darpink' which in combination distinguish this *Chrysanthe-*
mum as a new and distinct cultivar:

1. Inflorescence with oblong shaped, red-purple ray
florets, and bell shaped, yellow disc florets;
2. Flowering time of 7.5 weeks;
3. Short day plant; and
4. Vigorous growth habit.

Plants of the new cultivar 'Danatdarpink' differ from
plants of the parental cultivar 'Danchryatpink' as described
in Table 1.

TABLE 1

Characteristic	'Danatdarpink'	'Danchryatpink' (unpatented)
Inflorescence Diameter	9.5 cm.	9.5 cm.
Flowering Time	7.5 weeks	7.5 weeks
Growth Potency	Medium-High	Medium-High
Ray Floret Color	Red purple, RHS N74-C	Violet, RHS 84-C
Disk Floret Color (when opening)	Yellow green, RHS N144-B	Yellow green, RHS 146-B

Of the many commercial cultivars known to the present
inventor, the most similar in comparison to 'Danatdarpink'
is the parental cultivar 'Danchryatpink' as described in Table
1 .

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying photographs illustrate the overall appearance of the new *Chrysanthemum* cultivar 'Danatdarpink' showing the colors as true as is reasonably possible with colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description, which accurately describe the colors of 'Danatdarpink'.

FIG. 1 shows a side view perspective of a corymb of 'Danatdarpink'.

FIG. 2 shows a close-up view of an inflorescence of 'Danatdarpink'.

DETAILED BOTANICAL DESCRIPTION

'Danatdarpink' has not been observed under all possible environmental conditions. The phenotype of the new cultivar may vary with variations in environment such as temperature, light intensity, and daylength without any change in the genotype of the plant.

The aforementioned photographs, together with the following observations, measurements and values describe the new cultivar 'Danatdarpink' as grown in a greenhouse in Moshav Mishmar Hashiva, Israel, under conditions which closely approximate those generally used in commercial practice. Plants of 'Danatdarpink' are mainly grown in greenhouses, but can also be grown outdoors during the summer. The recommended temperature range for growing plants of 'Danatdarpink' in a greenhouse are: day temperature range of 22° C. to 23° C., and evening temperature range of 17° C. to 18° C. For premium growth conditions, plants of 'Danatdarpink' are irrigated and fertilized on a regular basis using moderate regimen of 2.5 EC in summer and 3.0 EC in winter. Plants of 'Danatdarpink' are treated with the growth retardant, Alar (B-Nine), at a concentration of 3 gr/L.

Color references are made to The Royal Horticultural Society Colour Chart (R.H.S.), ed. 2001, except where general colors of ordinary significance are used. Color values were taken under daylight conditions at approximately 12:00 noon in Moshav Mishmar Hashiva, Israel. The age of the plant described is 11 weeks from planting.

Classification:

Botanical.—*Chrysanthemum morifolium*.

Commercial.—*Chrysanthemum* cv. Atlantis Dark Pink.

Parentage:

Parental cultivar.—Induced mutation of *Chrysanthemum morifolium* cultivar 'Danchryatpink' (unpatented, disclosed in U.S. Plant patent application Ser. No. 10/825,115).

Propagation: Stem cutting.

Plant:

General appearance and form.—Height: About 90 cm. Width: 20 cm.

Shape.—Compound corymbiform.

Flowering response.—Short day type.

Flowering season.—Autumn and beginning of winter.

Winter hardiness.—Hardy in moderate climate.

Lastingness of the individual bloom.—3 to 4 weeks (on the stem).

Rooting habit.—Adventitious roots from cutting base.

Time to initiate roots.—6 to 7 days.

Fragrance.—None.

Foliage:

Leaf length.—10 cm.

Leaf width.—5.5 cm.

Overall shape of leaf.—Pinnatipartite.

Apex shape.—Rounded.

Base shape.—Wedge-shaped.

Margin.—Serrate.

Texture.—Slightly warty.

Color of upper surface.—Mature leaf: Yellow-green, RHS 147-A. Immature leaf: Yellow-green, RHS 147-A.

Color of lower surface.—Mature leaf: Yellow-green, RHS 147-B. Immature leaf: Yellow-green, RHS 147-B.

Venation.—Pattern: Pinnate. Color: Upper surface: Yellow-green, RHS 147-B. Lower surface: Yellow-green, RHS 147-C.

Petiole.—Length: 2 cm. Diameter: 4 mm. Color: Yellow-green, RHS 146-A.

Stem.—Average length: 85 to 90 cm. Average diameter: 8 to 10 mm. Color: Yellow-green, RHS 146-B. Internode length: 15 to 20 mm.

Inflorescence:

Flowering response.—Short day.

Quantity of inflorescences on average.—7 to 8 on average per stem.

Inflorescence bud.—Height: 1 cm. Diameter: 16 mm. Color: Yellow-green, RHS 146-B.

Inflorescence size.—Diameter: 9.5 cm. Depth (height): 18 to 20 mm. Diameter of disc: 18 mm.

Ray florets.—Number of ray florets per inflorescence: 30 to 32. Orientation: Upward. Length: 35 to 40 mm. Width: 12 to 13 mm. Apex width: 6 to 8 mm. Overall shape: Oblong. Apex shape: Rounded (obtuse). Base shape: Fused. Margin: Entire. Texture: Smooth. Color when opening: Red purple, RHS N74-C. Color when fully opened: Upper surface: Red purple, RHS N74-C. Lower surface: Purple group, RHS 76-B.

Disc florets.—Number of disc florets per inflorescence: 200 to 220. Length: 4 to 5 mm. Width: 1 mm. Overall shape: Bell-shaped (campanulate). Apex shape: Toothed. Base shape: Fused. Margin: Entire. Texture: Smooth. Color when opening: Yellow-green, RHS N144-B. Color when fully opened: Upper surface: Yellow, RHS 9-A. Lower surface: Yellow, RHS 9-C.

Phyllaries.—Average number: 15 to 18. Average length: 6 to 9 mm. Average width: 1 to 2 mm. Overall shape: Oblong. Apex shape: Rounded (obtuse). Base shape: Truncate. Margin: Entire. Color: Upper surface: Yellow-green, RHS 148-B. Lower surface: Yellow-green, RHS 144-A. Peduncles: Aspect: Upward. Length: 15 cm. Diameter: 3.5 to 4 mm. Texture: Slightly warty. Color: Yellow-green, RHS 146-B.

Reproductive organs:

Androecium.—Present in disc florets only. Stamen: Number: 5, fused as a tube. Length: 5 mm. Color: Yellow, RHS 13-A. Anthers: Color: Yellow, RHS 13-A. Pollen: Color: Yellow, RHS 13-A.

Gynoecium.—Present in both ray and disc florets. Style: Length: 5 to 6 mm. Color: Yellow, RHS 2-B. Stigma: Width: 0.3 mm. Color: Yellow, RHS 12-B. Ovary: Color: Yellow-green, RHS 149-D. Seeds: Length: 1.5 mm. Width: 0.8 to 1 mm. Shape: Oblong. Color: Grayed-green, RHS 197-A. Fruit: Shape: Achene. Color: Grayed-green, RHS 197-A.

Disease resistance/susceptibility: None observed under regular growing condition.

I claim:

1. A new and distinct *Chrysanthemum* plant named 'Danatdarpink', substantially as illustrated and described herein.

FIGURE 1




FIGURE 2

