



US00PP17109P2

(12) **United States Plant Patent**
Sanders

(10) **Patent No.:** **US PP17,109 P2**
(45) **Date of Patent:** **Sep. 19, 2006**

(54) **IMPATIENS PLANT NAMED 'IMTRASALM'**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(50) Latin Name: *Impatiens walleriana*
Varietal Denomination: **Imtrasalm**

(52) **U.S. Cl.** **Plt./317**
(58) **Field of Classification Search** **Plt./317**
See application file for complete search history.

(75) Inventor: **Monica Maria Adelheid Sanders,**
Grootebroek (NL)

Primary Examiner—Kent Bell
(74) *Attorney, Agent, or Firm*—Bruce Vrana

(73) Assignee: **Syngenta Seeds B.V.,** Enkhuizen (NL)

(57) **ABSTRACT**

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

A new *Impatiens* plant, characterized particularly as to novelty by a flat and strong outward spreading growth habit, freely branching with branches that tend to grow horizontally, resulting in a big and very floriferous plant with large salmon flowers.

(21) Appl. No.: **11/205,887**

1 Drawing Sheet

(22) Filed: **Aug. 17, 2005**

1

2

Latin name of the genus and species of the plant claimed
Impatiens walleriana.

Varietal denomination: 'Imtrasalm'.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new distinct cultivar of *Impatiens*, botanically known as *Impatiens walleriana*.

The new *Impatiens* is a product of a planned breeding program conducted in Enkhuizen, Netherlands. The new *Impatiens* cultivar is characterized by a flat and spreading growing habit, freely branching with branches that tend to grow horizontally, resulting in a big and very floriferous plant with large salmon flowers.

The present cultivar originated from a cross-pollination made in 1999, of a proprietary *Impatiens walleriana* selection identified as 'P1991-1,' not patented, as the female, or seed parent, with a proprietary *Impatiens walleriana* selection identified as 'P1224-1,' not patented, as the male, or pollen parent. 'Imtrasalm' differs from its parents primarily in flower color.

The present cultivar has been repeatedly asexually reproduced by cuttings in Enkhuizen, Netherlands, in Angers, France and in Gilroy, Calif. over a four-year period. The distinctive characteristics of this new *Impatiens* are stable and reproduced true to type in successive generations of asexual reproduction. It takes 6 to 8 weeks to produce a finished plant, depending on the temperature.

This new *Impatiens* plant is an annual in most climatical zones in the USA, only in zones 9 and 10 it is a perennial plant.

DESCRIPTION OF THE DRAWING

This new *Impatiens* plant is illustrated by the accompanying photographic drawing which shows blooms, buds and foliage of the plant in full color, the color showing being as true as can be reasonably obtained by conventional photographic procedures.

DESCRIPTION OF THE NEW CULTIVAR

The following detailed descriptions set forth the distinctive characteristics of this new *Impatiens*. The data which

defines these characteristics were collected from asexual reproductions carried out in Enkhuizen, Netherlands. The plant history was taken on 25 week old plants, blossomed under natural light in the field.

5 Color readings were taken in the office under ambient light. Color references are primarily to the R.H.S. Color Chart of The Royal Horticultural Society of London.

TABLE 1

Differences between the new cultivar 'Imtrasalm' and a similar cultivar		
	'Imtrasalm'	'Imtrarose'
Flower color	Salmon	Rose
Habit	Medium long trailing branches	Long trailing branches

10
15
*Imtrarose' co-pending U.S. Plant Patent application no. 11/051,269

The Plant

20 Parentage:

Female parent.—'P1991-1,' not patented.

Male parent.—'P12247-1,' not patented.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—10–12 days at 20° C.

Time to produce a rooted cutting.—4 weeks.

Root description.—Fine, fibrous, and white in color.

Plant description:

General appearance.—Makes medium long branches that tend to grow horizontally.

Habit:

Growth and branching habit.—Freely branching habit, freely flowering, vigorous.

Plant height.—24–28 cm.

Plant spread.—80–92 cm.

Lateral branch description:

Length.—20–28 cm.

Diameter.—3–6 mm.

Texture.—Smooth, glabrous.

Internode length.—0.4–9 cm.

Color.—RHS 145A.

Foliage description:

Arrangement.—Alternate.

Length, mature leaves.—5–5.5 cm.
Width, mature leaves.—3.2–3.4 cm.
Shape.—Ovate.
Apex.—Acuminate.
Base.—Acute.
Margin.—Serrate.
Texture.—Smooth, glabrous.
Venation pattern.—Pinnate.
Color young foliage, upper surface.—RHS 146A.
Color young foliage, lower surface.—RHS 147C and 182D.
Fully expanded foliage, upper surface.—RHS 146A.
Fully expanded foliage, lower surface.—RHS 148C and 182B.
Venation, upper surface.—RHS 144A.
Venation, lower surface.—RHS 146A.
Petiole length.—About 1.5 cm.
Petiole diameter.—About 2.5 mm.
Texture.—Smooth, glabrous.
Color.—RHS 144D.

Flower description:

Flower type and flowering habit.—Single.
Number of flowers per leaf axil.—2–4.
Number of flowers and flower buds per lateral branch.—9–13.
Flower position.—Above and beyond the foliage and typically facing upwards and outward.
Flower shape.—Rounded, mostly flat.
Flowers lasting.—About 7–9 days on the plant depending on environmental conditions.
Flowers.—Non persistent and not fragrant.
Flowering.—Indeterminate and continuous.
Flowering season.—Year round under greenhouse conditions. In the garden, flowering from spring until fall.
Flower diameter.—About 4.5 cm.

Flower buds:

Rate of opening.—From flower bud to fully open flower, typically about 6–8 days, depending on temperature.
Bud length (at stage of showing color).—About 1.3 cm.
Bud diameter.—About 1 cm.
Bud shape.—Ovoid.
Texture.—Smooth, glabrous.
Color, just before opening.—RHS 52B.

Petals:

Quantity.—Single, five per flower, imbricate.
Length of banner petal.—About 1.6–1.9 cm.
Width of banner petal.—About 2.7–3 cm.
Length lateral and basal petals.—About 2.2–2.5 cm.
Width lateral and basal petals.—About 2.3 cm.
Lateral and basal petal shape.—Orbicular-obovate.
Banner petal shape.—Broadly obcordate.

Petal apex.—Retuse to emarginate.
Petal base.—Attenuate.
Petal margin.—Entire.
Petal texture.—Smooth, satiny.
Petal color.—Upper surface: When opening and fully opened: RHS 52A. Lower surface: When opening and fully opened: RHS 52C.

Sepals:

Lateral sepals.—2.
Shape.—Ovate-lanceolate.
Apex.—Acute.
Margin.—Entire.
Length.—3–7 mm.
Texture upper and lower surface.—Smooth, glabrous.

Spur:

Length.—About 3.4 cm.
Texture.—Smooth, glabrous.
Aspect.—Curved.
Color.—RHS 185D.

Peduncles:

Length.—About 2.1 cm.
Diameter.—1 cm.
Aspect.—About 45° from the stem.
Color.—RHS 149B.
Bracts.—At least 2.
Shape.—Linear-lanceolate.
Apex.—Acute.
Length.—3–6 mm.
Color.—RHS 145A.

Reproductive organs:

Androecium.—Stamen number: Five fused at anthers, filaments free, hooded.
Anther shape.—Obovate/oval.
Anther length.—About 4–5 mm.
Anther color.—RHS 54B to 54C.
Amount of pollen.—No pollen.

Gynoecium:

Pistil number.—One.
Pistil length.—About 4 mm.
Stigma shape.—Columar, five segmented.
Stigma color.—Red purple.
Style color.—RHS 54A.
Ovary arrangement.—Five celled.
Ovary color.—RHS 143A.

Seed development: Seed development has not been observed.

Disease/pest tolerance: No known susceptibility to diseases or insects noted to date.

What is claimed is:

1. A new and distinct cultivar of *Impatiens* plant named 'Imtrasalm', as substantially illustrated and described herein.

* * * * *

