



US00PP17189P2

(12) **United States Plant Patent**
Sanders

(10) **Patent No.:** **US PP17,189 P2**

(45) **Date of Patent:** **Nov. 7, 2006**

(54) **IMPATIENS PLANT NAMED ‘IMTRAORAN’**

(50) Latin Name: *Impatiens walleriana*
Varietal Denomination: **Imtraoran**

(75) Inventor: **Monica Maria Adelheid Sanders,**
Grootebroek (NL)

(73) Assignee: **Syngenta Seeds B.V.,** Enkhuizen (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 46 days.

(21) Appl. No.: **11/206,439**

(22) Filed: **Aug. 18, 2005**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./317**

(58) **Field of Classification Search** **Plt./317**
See application file for complete search history.

Primary Examiner—Kent Bell

(74) *Attorney, Agent, or Firm*—Bruce Vrana

(57) **ABSTRACT**

A new *Impatiens* plant, characterized particularly as to novelty by a flat and strong outward spreading growth habit, freely branching with branches that tend to grow horizontally, resulting in a big and very floriferous plant with very large orange flowers.

1 Drawing Sheet

1

Latin name of the genus and species of the plant claimed: *Impatiens walleriana*.

Varietal denomination: ‘Imtraoran’.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new distinct cultivar of *Impatiens*, botanically known as *Impatiens walleriana*.

The new *Impatiens* is a product of a planned breeding program conducted in Enkhuizen, Netherlands. The new *Impatiens* cultivar is characterized by a flat and strong outward spreading growing habit, freely branching with branches that tend to grow horizontally, resulting in a big and floriferous plant with very large orange flowers.

The present cultivar originated from a cross-pollination made in 2001, of a proprietary *Impatiens* selection identified as ‘R3627-1,’ not patented, as the female, or seed parent, with a proprietary *Impatiens* selection identified as ‘R2299-1,’ not patented, as the male, or pollen parent. ‘Imtraoran’ differs from its parents primarily in flower color.

The present cultivar has been repeatedly asexually reproduced by cuttings in Enkhuizen, Netherlands, in Angers, France and in Gilroy, Calif. over a three-year period. The distinctive characteristics of this new *Impatiens* are stable and reproduced true to type in successive generations of asexual reproduction. It takes 6 to 8 weeks to produce a finished plant, depending on the temperature.

This new *Impatiens* plant is an annual in most climatical zones in the USA, only in zones 9 and 10 it is a perennial plant.

DESCRIPTION OF THE DRAWING

This new *Impatiens* plant is illustrated by the accompanying photographic drawing which shows blooms, buds and foliage of the plant in full color, the color showing being as true as can be reasonably obtained by conventional photographic procedures.

DESCRIPTION OF THE NEW CULTIVAR

The following detailed descriptions set forth the distinctive characteristics of this new *Impatiens*. The data which defines these characteristics were collected from asexual

2

reproductions carried out in Enkhuizen, Netherlands. The plant history was taken on 25 week old plants, blossomed under natural light in the field.

Color readings were taken in the office under ambient light. Color references are primarily to The R.H.S. Color Chart of The Royal Horticultural Society of London.

TABLE 1

Difference between the new cultivar ‘Imtraoran’ and a similar cultivar

	‘Imtraoran’	‘Imtrarose’
Flower color	Orange	Rose
Habit	Spreading and slightly upright	Spreading with long trailing branches
Flower size	Very large	Large

‘Imtrarose’ co-pending U.S. plant patent application Ser. No. 11/051,269.

The Plant:

Parentage:

Female parent.—‘R3627-1,’ not patented.

Male parent.—‘R2299-1,’ not patented.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—10–12 days at 20° C.

Time to produce a rooted cutting.—4 weeks.

Root description.—Fine, fibrous, and white in color.

Plant description:

General appearance.—Makes long branches that tend to grow horizontally.

Habit:

Growth and branching habit.—Freely branching habit, freely flowering, vigorous.

Plant height.—30–32 cm.

Plant spread.—85–100 cm.

Lateral branch description:

Length.—37–47 cm.

Diameter.—4–5 mm.

Texture.—Smooth, glabrous.

Internode length.—0.5–7 cm.

Color.—RHS 145B and 183B.

Foliage description:

Arrangement.—Alternate.

Length, mature leaves.—6.3–7.4 cm.
Width, mature leaves.—2.9–3.3 cm.
Shape.—Ovate.
Apex.—Acuminate.
Base.—Attenuate.
Margin.—Serrate.
Texture.—Smooth, glabrous.
Venation pattern.—Pinnate.
Color young foliage, upper surface.—RHS 148A, beneath some 187A.
Color young foliage, lower surface.—RHS 183C and a bit 147C.
Fully expanded foliage, upper surface.—RHS 147A.
Fully expanded foliage, lower surface.—RHS 183D and a bit 147C.
Venation, upper surface.—RHS 144B.
Venation, lower surface.—RHS 183C.
Petiole length.—About 1.5–2 cm.
Petiole diameter.—About 3 mm.
Texture.—Smooth, glabrous.
Color.—RHS 184B.

Flower description:

Flower type and flowering habit.—Single.
Number of flowers per leaf axil.—2.
Number of flowers and flower buds per lateral branch.—15–21.
Flower position.—Above and beyond the foliage and typically facing upwards and outward.
Flower shape.—Rounded, mostly flat.
Flowers lasting.—About 7–9 days on the plant depending on environmental conditions.
Flowers.—Non persistent and not fragrant.
Flowering.—Indeterminate and continuous.
Flowering season.—Year round under greenhouse conditions. In the garden, flowering from spring until fall.
Flower diameter.—About 5 cm.

Flower buds:

Rate of opening.—From flower bud to fully open flower, typically about 6–8 days, depending on temperature.
Bud length (at stage of showing color).—About 1.4 cm.
Bud diameter.—About 1 cm.
Bud shape.—Ellipsoidal.
Texture.—Smooth, glabrous.
Color, just before opening.—RHS 40B.

Petals:

Quantity.—Single, five per flower, imbricate.
Length of banner petal.—About 1.9–2.3 cm.
Width of banner petal.—About 2.8–3 cm.
Length lateral and basal petals.—About 2.7–3 cm.
Width lateral and basal petals.—About 2.3–2.5 cm.
Lateral and basal petal shape.—Obovate.
Banner petal shape.—Broadly obcordate.

Petal apex.—Retuse to emarginate.
Petal base.—Banner: rounded; lateral and basal: attenuate.
Petal margin.—Entire.
Petal texture.—Smooth, satiny.
Petal color.—Upper surface: When opening and fully opened: RHS 44B. Lower surface: When opening and fully opened: RHS 40C.

Sepals:

Lateral sepals.—2.
Shape.—Ovate-lanceolate.
Apex.—Acute.
Margin.—Entire.
Length.—3–7 mm.
Texture upper and lower surface.—Smooth, glabrous.

Spur:

Length.—About 3.3 cm.
Texture.—Smooth, glabrous.
Aspect.—Curved.
Color.—RHS 185A.

Peduncles:

Length.—About 2.7 cm.
Diameter.—1 cm.
Aspect.—About 45° from the stem.
Color.—RHS 144B.
Bracts.—At least 2.
Shape.—Linear-lanceolate.
Apex.—Acute.
Length.—3–6 mm.
Color.—RHS 144A; top: RHS 183B.

Reproductive organs:

Androecium.—Stamen number: Five fused at anthers, filaments free, hooded.
Anther shape.—Obovate.
Anther length.—About 5 mm.
Anther color.—RHS 46A.
Amount of pollen.—No pollen.

Gynoecium:

Pistil number.—One.
Pistil length.—About 5 mm.
Stigma shape.—Columnar, five segmented.
Stigma color.—Red purple.
Style color.—RHS 187D.
Ovary arrangement.—Five celled.
Ovary color.—RHS 146A.

Seed development: Seed development has not been observed.

Disease/pest tolerance: No known susceptibility to diseases or insects noted to date.

What is claimed is:

1. A new and distinct cultivar of *Impatiens* plant named 'Imtraoran', as substantially illustrated and described herein.

* * * * *

