



US00PP17308P2

(12) **United States Plant Patent**
Sanders

(10) **Patent No.:** **US PP17,308 P2**
(45) **Date of Patent:** **Dec. 26, 2006**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED 'INGBWIT'**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(50) Latin Name: *Impatiens hawkeri*
Varietal Denomination: **Ingbwit**

(52) **U.S. Cl.** **Plt./318**

(58) **Field of Classification Search** **Plt./318**
See application file for complete search history.

(75) Inventor: **Monica Maria Adelheid Sanders,**
Grootebroek (NL)

Primary Examiner—Kent Bell
Assistant Examiner—Annette H Para
(74) *Attorney, Agent, or Firm*—Bruce Vrana

(73) Assignee: **Syngenta Seeds B.V.,** Enkhuizen (NL)

(57) **ABSTRACT**

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 32 days.

A new and distinct cultivar of New Guinea *Impatiens* plant named 'Ingbwit' characterized by its large, rounded, white flowers, early flowering, green foliage, ball shaped growth habit with good basal branching and good vigor.

(21) Appl. No.: **11/221,458**

1 Drawing Sheet

(22) Filed: **Sep. 8, 2005**

1

2

Latin name of the genus and species of the plant claimed:
Impatiens hawkeri.
Varietal denomination: 'Ingbwit'.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new and distinct New Guinea *Impatiens* plant botanically known as *Impatiens hawkeri* referred to by the cultivar name 'Ingbwit.'

The new cultivar was developed in a controlled breeding program conducted by the inventor in Enkhuizen, Netherlands. The new New Guinea *Impatiens* cultivar has an early flowering response, flowers with attractive coloration and a good flowering behavior under field conditions.

The new cultivar is propagated from cuttings resulting from the cross in September 1999 of the proprietary New Guinea *Impatiens* selection identified as 'F 62-1' as female parent and the proprietary New Guinea *Impatiens* selection identified as 'F 78-1' as male parent. 'F 62-1' is not commercially available and has not been patented. 'F 78-1' is not commercially available and has not been patented.

As a result of this cross the present cultivar was selected in September 2000 in Enkhuizen, Netherlands and has been repeatedly asexually reproduced by cuttings in Enkhuizen, Netherlands, in Gilroy, Calif., and in Angers, France over a period of several years. The distinctive characteristics of this new *Impatiens* are stable and reproduced true to type in successive generations of asexual reproduction. It takes 8 to 10 weeks to produce a finished plant, starting from a rooted plug and planted in a 12 cm pot, depending on the temperature.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Ingbwit.' These characteristics in combination distinguish 'Ingbwit' as a new and distinct New Guinea *Impatiens* cultivar:

1. Large, rounded, white flowers
2. Ball shaped growth habit with flowers on top of the foliage
3. Green foliage

4. Good basal branching character
 5. Good outdoor vigor
- Plants of the new New Guinea *Impatiens* differ primarily from the plants of the female parent selection in the following characteristic:

Plants of the New Guinea *Impatiens* have a good branching habit with shorter branches, whereas the plants of the female parent have a good branching habit with longer branches.

Plants of the new New Guinea *Impatiens* differ primarily from the plants of the male parent selection in the following characteristic:

Plants of the New Guinea *Impatiens* have a large flower, whereas the plants of the male have a medium sized flower.

Plants of the new New Guinea *Impatiens* can be compared to plants of 'Kiraya,' not patented in the United States. In side-by-side comparisons conducted in Enkhuizen, Netherlands, plants of the new New Guinea *Impatiens* differed from plants of 'Kiraya' in the following characteristics:

1. Plants of the new New Guinea *Impatiens* have light green leaves, whereas the plants of the cultivar 'Kiraya' have green leaves.
2. Plants of the new New Guinea *Impatiens* have flat, white flowers, whereas plants of the cultivar 'Kiraya' have slightly cup shaped, white flowers.

DESCRIPTION OF THE DRAWING

This new New Guinea *Impatiens* plant is illustrated by the accompanying photographic drawing which shows blooms, buds and foliage of the plant in full color, the color showing being as true as can be reasonably obtained by conventional photographic procedures.

DESCRIPTION OF THE NEW CULTIVAR

The following detailed descriptions set forth the distinctive characteristics of this new New Guinea *Impatiens*. The data which defines these characteristics were collected from asexual reproductions carried out in Enkhuizen, Nether-

lands. The plant history was taken on 30 week old plants, blossomed under natural light in the field.

Color readings were taken in the laboratory under ambient light. Color references are primarily to The R.H.S. Colour Chart of The Royal Horticultural Society of London.

The plant:

Classification.—Botanical: *Impatiens hawkeri*.

Parentage:

Female parent.—Proprietary *Impatiens hawkeri* selection identified as code number 'F 62-1,' not patented.

Male parent.—Proprietary *Impatiens hawkeri* selection identified as code number 'F 78-1,' not patented.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—Approximately 7–14 days.

Time to produce a rooted cutting.—Approximately 21 days.

Root description.—Fine, fibrous, and white in color.

Plant description:

General appearance.—Outwardly spreading, low rounded and uniform mounded plant growth.

Habit:

Growth and branching habit.—Freely branching habit, freely flowering, moderate vigorous.

Crop time:

Plant height.—18–20 cm.

Plant spread.—42–44 cm.

Lateral branch description:

Length.—18 cm.

Diameter.—4–8 mm.

Texture.—Smooth, glabrous.

Internode length.—0.5–7 cm.

Color.—RHS 146A.

Foliage description:

Arrangement.—Opposite or in whorls.

Length, mature leaves.—5.3 cm.

Width, mature leaves.—1.8 cm.

Shape.—Lanceolate.

Apex.—Acute to acuminate.

Base.—Acute.

Margin.—Serrulate with fine ciliation.

Texture.—Smooth, glabrous.

Venation pattern.—Pinnate.

Color young foliage, upper surface.—RHS 137A.

Color young foliage, lower surface.—RHS 146A.

Fully expanded foliage, upper surface.—RHS 143A.

Fully expanded foliage, lower surface.—RHS 144A.

Venation, upper surface.—RHS 145A.

Venation, lower surface.—RHS 144B.

Petiole length.—About 0.4 cm.

Petiole diameter.—About 2 mm.

Texture.—Smooth, glabrous.

Color.—RHS 145A.

Flower description:

Flower type and flowering habit.—Single.

Number of flowers per leaf axil.—3.

Number of flowers and flower buds per lateral branch.—8.

Flower position.—Above and beyond the foliage and typically facing upwards and outward.

Flower shape.—Rounded, mostly flat.

Flowers lasting.—About 10 days on the plant depending on environmental conditions.

Petals.—Self cleaning.

Gynoecium.—Persistent.

Flowers.—Not fragrant.

Flowering.—Indeterminate and continuous.

Flowering season.—Year round under greenhouse conditions. In the garden, flowering from spring until fall.

Flower length.—About 4–4.7 cm.

Flower width.—About 3.8–4.8 cm.

Flower depth.—About 0.2 cm.

Flower buds:

Bud length.—About 1.3 cm.

Bud diameter.—About 0.9 cm.

Bud shape.—Ovoid.

Texture.—Smooth, glabrous.

Color, just before opening.—RHS 157B.

Petals:

Quantity.—Single, five per flower, imbricate.

Length of banner petal.—About 2.5–3 cm.

Width of banner petal.—About 1.5–2 cm.

Length lateral and basal petals.—About 2.5 cm.

Width lateral and basal petals.—About 2.5 cm.

Lateral and basal petal shape.—Obovate.

Banner petal shape.—Obovate.

Petal apex.—Retuse to emarginate.

Petal base.—Attenuate.

Petal margin.—Entire.

Petal texture.—Smooth, satiny.

Petal color.—Upper surface: When opening and fully opened: RHS 155A. Lower surface: When opening and fully opened: RHS 155D.

Spur:

Length.—About 3.5 cm.

Texture.—Smooth, glabrous.

Aspect.—Curved.

Color.—RHS 151D.

Peduncles:

Length.—About 3–3.5 cm.

Strength.—Strong, flexible.

Diameter.—0.15 cm.

Aspect.—About 45° from the stem.

Color.—RHS 145B.

Reproductive organs:

Androecium.—Stamen number: Five fused at anthers, filaments free, hooded.

Anther shape.—Obovate.

Anther length.—About 0.5 mm.

Anther color.—RHS 155B.

Amount of pollen.—Moderate.

Pollen color.—RHS 158B.

Gynoecium:

Pistil number.—One.

Pistil length.—About 4 mm.

Stigma shape.—Columnar, five segmented.

Stigma color.—Colorless.

Style color.—RHS N144D.

Ovary arrangement.—Five celled.

Ovary color.—RHS 144A.

Seed development.—Seed development has not been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Impatiens* has not been observed.

What is claimed is:

1. A new and distinct cultivar of New Guinea *Impatiens* plant named 'Ingbwit', as substantially illustrated and described herein.

* * * * *

