

(12) **United States Plant Patent**
Sanders

(10) **Patent No.:** **US PP17,320 P2**
(45) **Date of Patent:** **Dec. 26, 2006**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED
'INGMAGE'**

(50) Latin Name: ***Impatiens hawkeri***
Varietal Denomination: **Ingmage**

(75) Inventor: **Monica Maria Adelheid Sanders,**
Grootebroek (NL)

(73) Assignee: **Syngenta Seeds B.V.,** Enkhuizen (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/222,998**

(22) Filed: **Sep. 9, 2005**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./318**
(58) **Field of Classification Search** **Plt./318**
See application file for complete search history.

Primary Examiner—Anne Marie Grunberg

Assistant Examiner—Annette H Para

(74) *Attorney, Agent, or Firm*—Bruce Vrana

(57) **ABSTRACT**

A new and distinct cultivar of New Guinea *Impatiens* plant
named 'Ingmage' characterized by its large, rounded, red
flowers, early flowering, dark green foliage, ball shaped
growth habit with excellent basal branching and good out-
door vigor.

1 Drawing Sheet

1

Latin name of the genus and species of the plant claimed:
Impatiens hawkeri.
Varietal denomination: 'Ingmage'.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new and distinct New
Guinea *Impatiens* plant botanically known as *Impatiens*
hawkeri and referred to by the cultivar name 'Ingmage'.

The new cultivar was developed in a controlled breeding
program conducted by the inventor in Enkhuizen, Nether-
lands. The new New Guinea *Impatiens* cultivar has an early
flowering response, large flowers with attractive coloration
and a good flowering behavior under field conditions.

The new cultivar is propagated from cuttings resulting
from the cross in September 1999 of the proprietary New
Guinea *Impatiens* selection identified as 'F 47-1' as female
parent and the proprietary New Guinea *Impatiens* selection
identified as 'E112-1' as male parent. 'F 47-1' is not com-
mercially available and has not been patented. 'E112-1' is
not commercially available and has not been patented.

As a result of this cross the present cultivar was selected
in September 2000 in Enkhuizen, Netherlands and has been
repeatedly asexually reproduced by cuttings in Enkhuizen,
Netherlands, in Gilroy, Calif., and in Angers, France over a
period of several years. The distinctive characteristics of this
new *Impatiens* are stable and reproduced true to type in
successive generations of asexual reproduction. It takes 7 to
9 weeks to produce a finished plant, starting from a rooted
plug and planted in a 12 cm pot, depending on the tempera-
ture.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and
are determined to be the unique characteristics of 'Ingmage.'
These characteristics in combination distinguish 'Ingmage'
as a new and distinct New Guinea *Impatiens* cultivar:

1. Large, rounded, red flowers
2. Ball shaped growth habit with flowers on top of the
foliage
3. Dark green foliage

2

4. Excellent basal branching character
5. Good outdoor vigor

Plants of the new New Guinea *Impatiens* differ primarily
from the plants of the female parent selection in the follow-
ing characteristic:

Plants of the New Guinea *Impatiens* have large red
flowers and dark green foliage, whereas the plants of the
female parent have light purple flowers and green foliage.

Plants of the new New Guinea *Impatiens* differ primarily
from the plants of the male parent selection in the following
characteristic:

Plants of the New Guinea *Impatiens* have red flowers with
dark green foliage, whereas the plants of the male parent
have purple flowers and dark foliage.

Plants of the new New Guinea *Impatiens* can be compared
to plants of 'Fisimp 551,' U.S. Plant Pat. No. 13,702. In
side-by-side comparisons conducted in Enkhuizen,
Netherlands, plants of the new New Guinea *Impatiens*
differed from plants of the cultivar 'Fisimp 551' in the
following characteristics:

1. Plants of the new New Guinea *Impatiens* have flat, red
flowers, whereas the plants of the cultivar 'Fisimp 551'
have slightly cup shaped, red purple colored flowers.
2. Plants of the new New Guinea *Impatiens* have a ball
shaped growth habit, whereas plants of the cultivar
'Fisimp 551' have a bushy shaped growth habit.

DESCRIPTION OF THE DRAWING

This new New Guinea *Impatiens* plant is illustrated by the
accompanying photographic drawing which shows blooms,
buds and foliage of the plant in full color, the color showing
being as true as can be reasonably obtained by conventional
photographic procedures.

DESCRIPTION OF THE NEW CULTIVAR

The following detailed descriptions set forth the distinc-
tive characteristics of this new New Guinea *Impatiens*. The
data which defines these characteristics were collected from
asexual reproductions carried out in Enkhuizen, Nether-

lands. The plant history was taken on 30 week old plants, blossomed under natural light in the field.

Color readings were taken in the laboratory under ambient light. Color references are primarily to The R.H.S. Colour Chart of The Royal Horticultural Society of London.

The plant:

Classification.—Botanical: *Impatiens hawkeri*.

Parentage:

Female parent.—Proprietary *impatiens hawkeri* selection identified as code number 'F 47-1,' not patented.

Male parent.—Proprietary *Impatiens hawkeri* selection identified as Code number 'E112-1,' not patented.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—Approximately 7–14 days.

Time to produce a rooted cutting.—Approximately 21 days.

Root description.—Fine, fibrous, and white in color.

Plant description:

General appearance.—Outwardly spreading, low rounded and uniform mounded plant growth.

Habit:

Growth and branching habit.—Freely branching habit, freely flowering, vigorous.

Crop time:

Plant height.—23–26 cm.

Plant spread.—43–46 cm.

Lateral branch description:

Length.—20 cm.

Diameter.—4–5 mm.

Texture.—Smooth, glabrous.

Internode length.—1.5–8 cm.

Color.—RHS 187A.

Foliage description:

Arrangement.—Primarily in whorls.

Length, mature leaves.—5.5–7 cm.

Width, mature leaves.—1.8–2 cm.

Shape.—Lanceolate.

Apex.—Acute to acuminate.

Base.—Acute.

Margin.—Serrulate with fine ciliation.

Texture.—Smooth, glabrous.

Venation pattern.—Pinnate.

Color young foliage, upper surface.—RHS 139A.

Color young foliage, lower surface.—RHS N77A.

Fully expanded foliage, upper surface.—RHS 139A.

Fully expanded foliage, lower surface.—RHS 59A.

Venation, upper surface.—RHS 53A.

Venation, lower surface.—RHS 59A.

Petiole length.—About 1 cm.

Petiole diameter.—About 3 mm.

Texture.—Smooth, glabrous.

Color.—RHS N79.

Flower description:

Flower type and flowering habit.—Single.

Number of flowers per leaf axil.—2–4.

Number of flowers and flower buds per lateral branch.—6.

Flower position.—Above and beyond the foliage and typically facing upwards and outward.

Flower shape.—Rounded, mostly flat.

Flowers lasting.—About 10 days on the plant depending on environmental conditions.

Petals.—Self cleaning.

Gynoecium.—Persistent.

Flowers.—Not fragrant.

Flowering.—Indeterminate and continuous.

Flowering season.—Year round under greenhouse conditions. In the garden, flowering from spring until fall.

Flower length.—About 4.5–6 cm.

Flower width.—About 4–6 cm.

Flower depth.—About 0.2 cm.

Flower buds:

Bud length.—About 1.3 cm.

Bud diameter.—About 1.2 cm.

Bud shape.—Ovoid.

Texture.—Smooth, glabrous.

Color, just before opening.—RHS 53B.

Petals:

Quantity.—Single, five per flower, imbricate.

Length of banner petal.—About 3–4 cm.

Width of banner petal.—About 2–2.5 cm.

Length lateral and basal petals.—About 2–3 cm.

Width lateral and basal petals.—About 2–3 cm.

Lateral and basal petal shape.—Obcordate.

Banner petal shape.—Obcordate.

Petal apex.—Retuse to emarginate.

Petal base.—Attenuate.

Petal margin.—Entire.

Petal texture.—Smooth, satiny.

Petal color.—Upper surface: When opening and fully opened: RHS 53B. Lower surface: When opening and fully opened: RHS 179A.

Spur:

Length.—About 3–4 cm.

Texture.—Smooth, glabrous.

Aspect.—Curved.

Color.—RHS 46A.

Peduncles:

Length.—About 3–4 cm.

Strength.—Strong, flexible.

Diameter.—0.2 cm.

Aspect.—About 45° from the stem.

Color.—RHS N34A.

Reproductive organs:

Androecium.—Stamen number: Five fused at anthers, filaments free, hooded.

Anther shape.—Obovate.

Anther length.—About 4.5 mm.

Anther color.—RHS 45A.

Amount of pollen.—Moderate.

Pollen color.—RHS 155D.

Gynoecium:

Pistil number.—One.

Pistil length.—About 5 mm.

Stigma shape.—Columnar, five segmented.

Stigma color.—Colorless.

Style color.—RHS 39C.

Ovary arrangement.—Five celled.

Ovary color.—RHS N144C.

Seed development.—Seed development has not been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Impatiens* has not been observed.

What is claimed is:

1. A new and distinct cultivar of New Guinea *Impatiens* plant named 'Ingmage', as substantially illustrated and described herein.

* * * * *

