



US00PP17363P2

(12) **United States Plant Patent**
Oudshoorn

(10) **Patent No.:** **US PP17,363 P2**
(45) **Date of Patent:** **Jan. 16, 2007**

(54) **PAPAYER PLANT NAMED ‘CASTANETTE’**

(50) Latin Name: *Papaver orientale*
Varietal Denomination: **Castanette**

(75) Inventor: **Hubertus Gerardus Oudshoorn,**
Rijpwetering (NL)

(73) Assignee: **Future Plants Licantie B.V.,**
Lisserbroek (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/141,585**

(22) Filed: **May 31, 2005**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./263**

(58) **Field of Classification Search** Plt./263
See application file for complete search history.

Primary Examiner—Kent Bell

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A distinct cultivar of *Papaver* plant named ‘Castanette’, characterized by its upright and relatively compact plant habit; moderately vigorous growth habit; freely flowering habit; and large flowers with light red-colored petals.

1 Drawing Sheet

1

Botanical designation: *Papaver orientale*.
Cultivar denomination: ‘Castanette’.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct cultivar of *Papaver* plant, botanically known as *Papaver orientale*, commercially referred to as Oriental Poppy, and hereinafter referred to by the cultivar name ‘Castanette’.

The new *Papaver* is a product of a planned breeding program conducted by the Inventor in Rijpwetering, The Netherlands. The objective of the breeding program is to develop new *Papaver* cultivars with compact plant habit and numerous flowers.

The new *Papaver* originated from a cross-pollination made by the Inventor in 2000 of two unnamed selections of *Papaver orientale*, not patented. The new *Papaver* was discovered and selected by the Inventor in 2001 in a controlled environment in Rijpwetering, The Netherlands, from the resultant progeny of the stated cross-pollination on the basis of its compact plant habit and flower color.

Asexual reproduction of the new cultivar by cuttings at Rijpwetering, The Netherlands since 2001, has shown that the unique features of this new *Papaver* are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the cultivar Castanette have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Castanette’. These characteristics in combination distinguish ‘Castanette’ as a new and distinct *Papaver* cultivar:

1. Upright and relatively compact plant habit.
2. Moderately vigorous growth habit.
3. Freely flowering habit.
4. Large flowers with light red-colored petals.

2

Plants of the cultivar Castanette differ from plants of the parent selections and other *Papaver* cultivars known to the Inventor, such as the cultivar Watermelon, not patented, in plant height as plants of the new *Papaver* are more compact than plants of the parent selections and other cultivars known to the Inventor.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the actual colors of the new *Papaver*.

The photograph at the top of the sheet comprises a close-up view of a typical flower of ‘Castanette’.

The photograph at the bottom of the sheet is a close-up view of a typical leaf of ‘Castanette’.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used. Plants used for the abovementioned photographs and following observations and measurements were grown in outdoor nursery in Roelofarendsveen, The Netherlands, and under conditions which closely approximate commercial production conditions during the spring and summer. The photographs, measurements and observations were taken during June when the plants were about one year old. During this time, day temperatures ranged from 14 to 27° C. and night temperatures ranged from 7 to 16° C.

Botanical classification: *Papaver orientale* cultivar Castanette.

Parentage:

Female, or seed, parent.—Unnamed selection of *Papaver orientale*, not patented.

Male, or pollen, parent.—Unnamed selection of *Papaver orientale*, not patented.

Propagation:

Type.—By cuttings.

Time to develop roots.—About 90 days at 18° C.

Root description.—Fleshy, thick; light brown to white in color.

Plant description:

Form.—Herbaceous perennial flowering plant; mostly upright; narrow inverted triangle. Basally branching, rosette; flowering stems arise from the base. Moderately vigorous.

Plant height.—About 56 cm.

Plant diameter.—About 34 cm.

Foliage description.—Arrangement: Alternate; simple. Shape: Lanceolate, pinnately and deeply dissected. Length: About 28.3 cm. Width: About 11.7 cm. Leaf segment shape: Lanceolate; mid-section and base, pinnatisect. Leaf segment apex: Acute. Leaf segment base: Cuneate. Leaf segment margin: Serrate; occasionally biserrate. Texture, upper and lower surfaces: Pubescent; very rough. Venation pattern: Pinnate. Color: Developing foliage, upper surface: 143A. Developing foliage, lower surface: 143A to 143B. Fully expanded foliage, upper surface: 137A. Fully expanded foliage, lower surface: 137C. Venation, upper surface: 138C to 138D. Venation, lower surface: 143B. Petioles: Length: Basal leaves, about 9.1 cm; stem leaves, about 7 mm. Diameter: About 7 mm. Color: 144B to 144C.

Flower description:

Flower type and habit.—Single rotate flowers; flowers terminal and face mostly upright. Flowers not persistent. One flower per flowering stem.

Natural flowering season.—Plants flower continuously from June to early July in The Netherlands.

Fragrance.—None detected.

Flower longevity on the plant.—About three days.

Flower size.—Diameter: About 9.7 cm. Depth (height): About 5.6 cm.

Flower buds.—Length: About 2.7 cm. Diameter: About 1.7 cm. Shape: Ovoid. Texture: Pubescent; rough. Color: 137A to 137B.

Petals.—Arrangement/appearance: Rotate; about four petals forming a cup-shaped corolla. Length: About 5 cm. Width: About 5.6 cm. Shape: Broadly obovate to reniform. Apex: Irregularly praemorse; slightly

ruffled. Margin: Entire; slightly ruffled. Texture, upper and lower surfaces: Smooth; glabrous. Color: When opening and fully opened, upper surface: 40A; towards the base, 187B to 187C; small basal spots, closest to 202A. When opening and fully opened, lower surface: 40A to 40B; towards the base, 187B; occasional green mark towards the apex, 143A to 143B.

Sepals.—Quantity: About three. Length: About 2.6 cm. Width: About 1.1 cm. Shape: Pinnatisect. Apex: Acute. Margin: Serrate. Texture, upper and lower surfaces: Densely pubescent. Color, upper and lower surfaces: Between 197A and N199A.

Flowering stems (peduncles).—Length: About 51.4 cm. Diameter: About 8 mm. Internode length: About 17.3 cm. Strength: Strong. Aspect: Erect. Texture: Pubescent; rough. Color: 144A.

Reproductive organs.—Stamens: Quantity per flower: Numerous, typically about 300. Stamen length: About 1.6 cm. Anther shape: Narrowly oblong; basifixed; occasionally curved. Anther length: About 2 mm. Anther color: Between N186C and 202A. Filament length: About 1.4 cm. Filament color: 187A; towards the base, 155D. Pollen amount: Moderate. Pollen color: Between 197A and 202A. Pistils: Quantity per flower: About 13. Pistil length: About 6 mm. Stigma shape: Narrowly oblong, flat. Stigma color: Slightly more purple than 200A. Ovary color: 145A to 145B.

Fruits.—Quantity per flower: One. Type: Capsule. Length: About 1.8 cm. Diameter: About 1.8 cm. Color: 137A.

Seeds.—Quantity per fruit: About 900. Length: About 1 mm. Diameter: About 1 mm. Color: Closest to 202A to 202B.

Disease/pest resistance: Plants of the new *Papaver* have been noted to be resistant to pathogens and pests common to *Papaver*.

Temperature tolerance: Plants of the new *Papaver* have been observed to tolerate temperatures from -25 to 40° C.

It is claimed:

1. A new and distinct cultivar of *Papaver* plant named 'Castanette', as illustrated and described.

* * * * *



‘Castanette’

