



US00PP17475P2

(12) **United States Plant Patent**
Bergman

(10) **Patent No.:** **US PP17,475 P2**
(45) **Date of Patent:** **Mar. 6, 2007**

(54) **CHRYSANTHEMUM PLANT NAMED**
‘YOROCKPORT’

(50) Latin Name: *Chrysanthemum*×*morifolium*
Varietal Denomination: **Yorockport**

(75) Inventor: **Wendy R. Bergman**, Lehigh Acres, FL
(US)

(73) Assignee: **Yoder Brothers, Inc.**, Barberton, OH
(US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 30 days.

(21) Appl. No.: **11/157,451**

(22) Filed: **Jun. 21, 2005**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./286**

(58) **Field of Classification Search** Plt./286
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

PP6,084 P * 1/1988 Hesse Plt./286

PP10,207 P * 1/1998 Pieters Plt./286
PP10,211 P * 1/1998 Pieters Plt./286
PP10,405 P * 5/1998 VandenBerg Plt./286
PP11,203 P * 2/2000 VandenBerg Plt./286
PP13,034 P2 * 10/2002 Bergman Plt./286

OTHER PUBLICATIONS

UPOV ROM GTITM Computer Database, GTI Jouve
Retrieval Software 2006/05 Citation for ‘Yorockport’.*

* cited by examiner

Primary Examiner—Wendy Haas

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Chrysanthemum* plant named
‘Yorockport’, characterized by its uniform and somewhat
outwardly spreading plant habit; strong and freely branching
growth habit; dark green-colored foliage; uniform flowering
response and habit; can be grown as a spray-type or without
bud removal; early flowering habit; daisy-type inflores-
cences with elongated oblong-shaped ray florets; intense red
purple-colored ray florets; and excellent postproduction lon-
gevity with plants maintaining good substance and color for
about four weeks in an interior environment.

2 Drawing Sheets

1

Botanical designation: *Chrysanthemum*×*morifolium*.
Cultivar denomination: ‘Yorockport’.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of *Chrysanthemum* plant, botanically known as
Chrysanthemum×*morifolium* and hereinafter referred to by
the name ‘Yorockport’.

The new *Chrysanthemum* is a product of a planned
breeding program conducted by the Inventor in Salinas,
Calif., and Alva, Fla. The objective of the program is to
create or discover new potted *Chrysanthemum* cultivars that
are suitable for year-round production with uniform plant
growth habit, good vigor and strong branching habit, numer-
ous inflorescences, desirable inflorescence form and floret
colors, fast and uniform flowering response, and good
postproduction longevity.

The new *Chrysanthemum* originated from a cross-
pollination made in October, 1997 in Salinas, Calif., of a
proprietary selection of *Chrysanthemum*×*morifolium* iden-
tified as code number YB-4891, not patented, as the female,
or seed, parent with a proprietary selection of
Chrysanthemum×*morifolium* identified as code number
YB-4554, not patented, as the male, or pollen, parent. The
new *Chrysanthemum* was discovered and selected by the
Inventor as a single flowering plant within the progeny of the
stated cross-pollination grown in a controlled environment
in Alva, Fla. in December, 1998. The selection of this plant

2

was based on its uniform plant growth habit, good vigor and
strong branching habit, desirable inflorescence form and
floret colors, fast and uniform flowering response, and good
postproduction longevity.

5 Asexual reproduction of the new *Chrysanthemum* by
vegetative tip cuttings was first conducted in Alva, Fla. in
March, 1999. Asexual reproduction by cuttings has shown
that the unique features of this new *Chrysanthemum* are
stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

The cultivar Yorockport has not been observed under all
possible environmental conditions. The phenotype may vary
somewhat with variations in environment such as
temperature, daylength, and/or light level, without, however,
any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Yorock-
port’. These characteristics in combination distinguish
‘Yorockport’ as a new and distinct *Chrysanthemum*:

1. Uniform and somewhat outwardly spreading plant
habit.
2. Strong and freely branching growth habit.
3. Dark green-colored foliage.
4. Uniform flowering response and habit.
5. Can be grown as a spray-type or without bud removal.
6. Early flowering, eight week response time.

7. Daisy-type inflorescences with elongated oblong-shaped ray florets.
8. Intense red purple-colored ray florets.
9. Excellent post-production longevity with plants maintaining good substance and color for about four weeks in an interior environment.

Plants of the new *Chrysanthemum* can be compared to plants of the female parent selection. Plants of the new *Chrysanthemum* differ from plants of the female parent selection primarily in plant form as plants of the female parent selection are more upright. In addition, plants of the new *Chrysanthemum* have darker colored ray florets than plants of the female parent selection.

Plants of the new *Chrysanthemum* can be compared to plants of the male parent selection. Plants of the new *Chrysanthemum* differ from plants of the male parent selection primarily in ray floret color as plants of the male parent selection have yellow-colored ray florets.

Plants of the new *Chrysanthemum* can be compared to plants of the cultivar Yolompoc, disclosed in U.S. Plant Pat. No. 11,203. In side-by-side comparisons conducted in Alva, Fla., plants of the new *Chrysanthemum* differed from plants of the cultivar Yolompoc in the following characteristics:

1. Plants of the new *Chrysanthemum* were more freely flowering than plants of the cultivar Yolompoc.
2. Ray florets of plants of the new *Chrysanthemum* were darker in color than ray florets of plants of the cultivar Yolompoc.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Chrysanthemum* showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ from the color values cited in the detailed botanical description which accurately describe the colors of the new *Chrysanthemum*.

The photograph on the first sheet comprises a side perspective view of typical flowering plants of 'Yorockport' grown as spray-types.

The photograph on the second sheet comprises a close-up view of typical inflorescences of 'Yorockport' grown as spray-types.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. The aforementioned photographs, following observations and measurements describe plants grown and flowered during the spring in Salinas, Calif., in a fiberglass-covered greenhouse and under conditions which approximate those generally used in commercial potted *Chrysanthemum* production. During the production of these plants, the following conditions were measured: day temperatures, 21° C. to 27° C.; night temperatures, 17° C. to 19° C.; and light levels, 5,000 to 6,000 foot-candles. Four unrooted cuttings were directly stuck in 15-cm containers, exposed to long day/short night conditions, and pinched once about two weeks later. At the time of the pinch, the photoinductive short day/long night treatments were initiated. Plants used for the description were grown as spray-types. Measurements and numerical values represent averages of typical flowering plants.

Botanical classification: *Chrysanthemum*×*morifolium* cultivar Yorockport.

Commercial classification: Daisy-type potted *Chrysanthemum*.

Parentage:

Female, or seed, parent.—Proprietary selection of *Chrysanthemum*×*morifolium* identified as code number YB-4891, not patented.

Male, or pollen, parent.—Proprietary selection of *Chrysanthemum*×*morifolium* identified as code number YB-4554, not patented.

Propagation:

Type.—Terminal tip cuttings.

Time to initiate roots.—About four days at 21° C.

Time to produce a rooted cutting.—About ten days at 21° C.

Root description.—Fibrous; white, close to 155D, in color.

Rooting habit.—Freely branching; moderately dense.

Plant description:

Appearance.—Herbaceous daisy-type potted *Chrysanthemum* that can be grown as a spray-type or without bud removal. Upright with lateral branches somewhat outwardly spreading; uniformly mounded crown. Strong and freely branching growth habit; about four lateral branches develop after removal of terminal apex (pinching); dense and full plants.

Plant height.—About 29 cm.

Plant width.—About 41 cm.

Lateral branches.—Length: About 21 cm. Diameter: About 4 mm. Internode length: About 1.5 cm. Strength: Strong. Texture: Pubescent. Color: 146A.

Foliage description.—Arrangement: Alternate; simple. Length: About 6.7 cm. Width: About 5.6 cm. Apex: Mucronate. Base: Truncate. Margin: Palmately lobed, sinuses between lateral lobes parallel to convergent. Texture, upper and lower surfaces: Pubescent. Color: Developing and fully expanded foliage, upper surface: Close to 147A. Developing and fully expanded foliage, lower surface: More green than 147B. Venation, upper surface: Close to 147A. Venation, lower surface: Close to 147B. Petiole length: About 2 cm. Petiole diameter: About 3 mm. Petiole texture, upper and lower surfaces: Pubescent. Petiole color, upper surface: More green than 147A. Petiole color, lower surface: Close to 147B.

Inflorescence description:

Appearance.—Daisy-type inflorescence form with elongated oblong-shaped ray florets. Inflorescences borne on terminals above foliage. Disk and ray florets develop acropetally on a capitulum. Inflorescences not fragrant.

Flowering response.—Under natural conditions, plants flower in the autumn/winter in the Northern Hemisphere. At other times of the year, inflorescence initiation and development can be induced under short day/long night conditions (at least 13.5 hours of darkness). Uniform and early flowering habit; plants exposed to two weeks of long day/short night conditions followed by photoinductive short day/long night conditions flower about eight weeks later.

Postproduction longevity.—Inflorescences maintain good color and substance for about four weeks in an interior environment.

Quantity of inflorescences.—Grown as a spray-type, about five per lateral branch.

5

Inflorescence bud.—Height: About 6 mm. Diameter: About 7 mm. Shape: Oblate. Color: 146A to 147A.

Inflorescence diameter.—About 7.4 cm.

Inflorescence height.—About 1.3 cm.

Diameter of disc.—About 2 cm.

Receptacle diameter.—About 7 mm.

Receptacle height.—About 7 mm.

Ray florets.—Length: About 3.7 mm. Width: About 9 mm. Corolla tube length: About 5 mm. Shape: Elongated oblong. Apex: Acute, rounded or emarginate. Base: Fused into a corolla tube. Margin: Entire. Orientation: Initially upright to eventually reflexed. Texture, upper and lower surfaces: Smooth, glabrous; satiny. Number of ray florets per inflorescence: About 22 arranged in a single whorl. Color: When opening and fully opened, upper surface: Closest to 72A. When opening and fully opened, lower surface: 155D underlain with close to 77A.

Disc florets.—Arrangement: Massed at center of receptacle. Length: About 8 mm. Diameter, apex: About 2.5 mm. Diameter, base: About 2 mm. Shape: Tubular; elongated. Apex: Five-pointed. Number of disc florets per inflorescence: About 150. Color: Immature: Close to 144A to 144B. Mature, apex: Close to 9A. Mature, mid-section: Close to 144B. Mature, base: Close to 155D.

6

Phyllaries.—Quantity per inflorescence: About 20 arranged in about two to three whorls. Length: About 9 mm. Width: About 3.5 mm. Shape: Lanceolate. Apex: Acute. Base: Truncate. Margin: Entire. Texture, upper surface: Waxy, smooth. Texture, lower surface: Pubescent. Color, upper surface: Close to 146A. Color, lower surface: 146A to 147A.

Peduncles.—Length, terminal peduncle: About 3.1 cm. Length, fourth peduncle: About 5.5 cm. Diameter: About 2.5 mm. Strength: Strong. Texture: Pubescent. Angle: About 45° from vertical. Color: Close to 146A.

Reproductive organs.—Androecium: Present on disc florets only. Anther length: Less than 1 mm. Anther color: Close to 15A. Pollen amount: Scarce. Pollen color: Close to 15A. Gynoecium: Present on both ray and disc florets. Style length: About 5 mm. Style color: Close to 154D. Stigma color: Close to 9A.

Seed/fruit.—Seed and fruit production has not been observed.

Disease/pest resistance: Resistance to pathogens and pests common to *Chrysanthemums* has not been observed on plants grown under commercial greenhouse conditions. It is claimed:

1. A new and distinct cultivar of *Chrysanthemum* plant named 'Yorockport', as illustrated and described.

* * * * *



