



(12) **United States Plant Patent**
Goemans

(10) **Patent No.:** **US PP17,499 P2**
(45) **Date of Patent:** **Mar. 13, 2007**

(54) **ALSTROEMERIA PLANT NAMED**
‘MAESTRO’

(50) Latin Name: *Alstroemeria hybrida*
Varietal Denomination: **Maestro**

(75) Inventor: **Frank Goemans**, Chichester (GB)

(73) Assignee: **Wulfinghoff Alstroemerias B.V.**,
Rijswijk (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 18 days.

(21) Appl. No.: **11/177,973**

(22) Filed: **Jul. 8, 2005**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./309**
(58) **Field of Classification Search** **Plt./309**
See application file for complete search history.

Primary Examiner—Anne Marie Grunberg
Assistant Examiner—S. B. McCormick-Ewoldt
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Alstroemeria* plant named
‘Maestro’, characterized by its upright and strong flowering
stems; durable and long-lasting leaves; freely flowering
habit; large umbels with large red purple-colored flowers;
and good postproduction longevity.

1 Drawing Sheet

1

Botanical designation: *Alstroemeria hybrida*.
Cultivar denomination: ‘Maestro’.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of *Alstroemeria* plant, botanically known as *Alstroeme-
ria hybrida*, commercially used as a cut flower *Alstroemeria*,
and hereinafter referred to by the name ‘Maestro’.

The new *Alstroemeria* is a product of a planned breeding
program conducted by the Inventor in Chichester, United
Kingdom. The objective of the breeding program was to
develop new cut flower *Alstroemeria* cultivars with freely
flowering habit, large flowers, attractive flower colors and
excellent postproduction longevity.

The new *Alstroemeria* originated from a cross-pollination
made by the Inventor in Chichester, United Kingdom, in
1999 of a selection of *Alstroemeria hybrida* identified as
code number 1302/19, not patented, as the female, or seed,
parent with a *Alstroemeria hybrida* selection identified as
code number 390/6, not patented, as the male, or pollen,
parent. The new *Alstroemeria* was discovered and selected
by the Inventor as a flowering plant within the progeny of
the stated cross-pollination in a controlled environment in
Chichester, United Kingdom in 1999.

Asexual reproduction of the new cultivar by tissue culture
in a controlled environment in Chichester, United Kingdom,
has shown that the unique features of this new *Alstroemeria*
are stable and reproduced true to type in successive genera-
tions of asexual propagation.

SUMMARY OF THE INVENTION

Plants of the cultivar Maestro have not been observed
under all possible environmental conditions. The phenotype
may vary somewhat with variations in environment such as
temperature and light intensity without, however, any vari-
ance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Maestro’.
These characteristics in combination distinguish ‘Maestro’
as a new and distinct cultivar:

2

1. Upright and strong flowering stems.
2. Durable and long-lasting leaves.
3. Freely flowering habit.
4. Large umbels with large red purple-colored flowers.
5. Good postproduction longevity.

Plants of the new *Alstroemeria* can be compared to plants
of the parent selections. Plants of the new *Alstroemeria* have
more durable foliage than plants of the parent selections. In
addition, plants of the new *Alstroemeria* have larger flowers
than plants of the parent selections.

Plants of the new *Alstroemeria* can also be compared to
plants of the cultivar Patricia, disclosed in a U.S. Plant
patent application Ser. No. 11/177,976 filed concurrently. In
side-by-side comparisons conducted in Geldern, Germany,
plants of the new *Alstroemeria* differed from plants of the
cultivar Patricia in the following characteristics:

1. Plants of the new *Alstroemeria* had shorter flowering
stems than plants of the cultivar Patricia.
2. Plants of the new *Alstroemeria* had larger inflores-
cences and larger flowers than plants of the cultivar
Patricia.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrative the
overall appearance of the new *Alstroemeria*, showing the
colors as true as it is reasonably possible to obtain in colored
reproductions of this type. Colors in the photographs may
differ slightly from the color values cited in the detailed
botanical description which accurately describe the colors of
the new *Alstroemeria*.

The photograph at the top of the sheet comprises a side
perspective view of a typical flowering stem of ‘Maestro’
grown in a container.

The photograph at the bottom of the sheet is a close-up
view of a typical flower and leaves of ‘Maestro’.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs, following observations and measurements describe plants of the new *Alstroemeria* grown during the spring in ground beds in Geldern, Germany in a glass-covered greenhouse for about five months. During the production of the plants, day temperatures were about 11° C. and night temperatures were about 7° C. Color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Alstroemeria hybrida* cultivar Maestro.

Parentage:

Female parent.—*Alstroemeria hybrida* selection identified as code number 1302/19, not patented.

Male parent.—*Alstroemeria hybrida* selection identified as code number 390/6, not patented.

Propagation:

Type.—By tissue culture.

Root description.—Fibrous, fleshy, thick; white, close to 155D, in color.

Rooting habit.—Freely branching.

Rhizomes.—Shape: Elongate; rounded. Length: About 5 cm. Diameter: About 9 mm. Texture: Smooth. Color: 155A.

Plant description:

Plant habit.—Upright and strong flowering stems; freely basal branching, bushy appearance. Vigorous growth habit.

Plant height.—About 160 cm.

Plant diameter (spread).—About 75 cm.

Number of harvestable flowering stems per year.—About 85 to 110 per plant.

Flowering stem description.—Aspect: Upright. Length: About 160 cm. Internode length: About 4 to 77 cm. Strength: Strong. Texture: Glabrous, smooth. Color: 137C.

Foliage description.—Length, largest: About 14 cm. Width, largest: About 3.5 cm. Shape: Lanceolate. Apex: Acute. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Venation pattern: Parallel. Durability: Durable, long-lasting. Petiole length: About 3 cm. Petiole diameter: About 6 mm. Petiole texture, upper and lower surfaces: Smooth, glabrous. Color: Developing foliage, upper and lower surfaces: 136B. Fully expanded foliage, upper surface: 136A. Fully expanded foliage, lower surface: 137B. Venation, upper surface: 139A. Venation, lower surface: 136B. Petiole, upper surface: Close to 136A. Petiole, lower surface: Close to 137B.

Flower description:

Flower type and habit.—Single cup-shaped flowers arranged in compound umbels. Flowers face mostly outward. Perianth segments separate. Freely and continuously flowering. Flowers not persistent.

Natural flowering season.—Flowering continuous year-round under greenhouse conditions.

Fragrance.—None detected.

Flower longevity on the plant.—About 7 to 18 days.

Flower longevity as a cut flower.—About seven to nine days.

Flower buds (showing color).—Length: About 7.5 cm. Diameter: About 2.5 cm. Shape: Ovoid. Color: 72B; towards the apex, 135B.

Umbel height.—About 22 cm.

Umbel diameter.—About 28 cm.

Number of flowers per umbel.—About three to six.

Flower diameter.—About 9 cm.

Flower height.—About 5 cm.

Perianth.—Arrangement: Six arranged in two whorls, each whorl with two lateral and one median segments. Size, inner perianth segments: Length, lateral segments: About 6 cm. Length, median segment: About 5.5 cm. Width, lateral segments: About 2 cm. Width, median segment: About 2 cm. Size, outer perianth segments: Length, lateral segments: About 6.3 cm. Length, median segment: About 6.3 cm. Width, lateral segments: About 3.4 cm. Width, median segment: About 3.4 cm. Shape, inner perianth, all segments: Lanceolate. Shape, outer perianth, all segments: Rounded. Apex, inner perianth, all segments: Acute; mucronate. Apex, outer perianth, all segments: Bracket-shaped; mucronate. Base, inner and outer perianths, all segments: Attenuate. Margin, inner and outer perianths, all segments: Entire. Texture, inner and outer perianths, all segments: Smooth, glabrous. Color, inner perianth: When opening and fully opened, lateral segments, upper surface: 72A; center section, 13B; spots, close to 59A. When opening and fully opened, median segment, upper surface: 72A; spots, close to 59A. When opening and fully opened, lateral and median segments, lower surface: 72A. Color, outer perianth: When opening and fully opened, lateral and median segments, upper surface: 72A; spots, close to 59A. When opening and fully opened, lateral and median segments, lower surface: 72A.

Pedicels.—Length: About 7 cm. Diameter: About 3 mm. Strength: Strong. Angle: About 45° from vertical. Texture: Smooth, glabrous. Color: 137A to 137B.

Reproductive organs.—Stamens: Quantity per flower: Six. Anther shape: Oval. Anther length: About 9 mm. Anther diameter: About 2 mm. Anther color: 183B. Pollen amount: Abundant. Pollen color: N186A. Pistils: Quantity per flower: One. Pistil length: About 5 cm. Style length: About 3 cm. Style color: 54A. Stigma color: 54A. Ovary color: 147A.

Fruits.—Shape: Globular. Length: About 7 mm. Diameter: About 6 mm. Color: 146B.

Seeds.—Seed development has not been observed.

Disease/pest resistance: Plants of the new *Alstroemeria* have not been observed to be resistant to pathogens and pests common to *Alstroemerias*.

Temperature tolerance: Plants of the new *Alstroemeria* have been observed to tolerate temperatures from 0 to 40° C. It is claimed:

1. A new and distinct cultivar of *Alstroemeria* plant named ‘Maestro’, as illustrated and described.

* * * * *

