



US00PP17784P2

(12) **United States Plant Patent**
Dekker

(10) **Patent No.:** **US PP17,784 P2**
(45) **Date of Patent:** **Jun. 5, 2007**

(54) **CHRYSANTHEMUM PLANT NAMED**
'DEKMALLORCA'

(50) Latin Name: *Chrysanthemum*×*morifolium*
Varietal Denomination: **Dekmallorca**

(75) Inventor: **Cornelis W. Dekker**, Hensbroek (NL)

(73) Assignee: **Dekker Breeding B.V.**, Hensbroek
(NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/267,901**

(22) Filed: **Nov. 4, 2005**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./291**

(58) **Field of Classification Search** **Plt./291,**
Plt./287

See application file for complete search history.

Primary Examiner—Kent Bell

Assistant Examiner—June Hwu

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Chrysanthemum* plant named
'Dekmallorca', characterized by its decorative-type inflo-
rescences with elliptic to oblong-shaped, dark pink-colored
ray florets; strong and upright flowering stems; freely flow-
ering habit; early and uniform flowering response; and good
postproduction longevity.

2 Drawing Sheets

1

Botanical designation: *Chrysanthemum*×*morifolium*.
Cultivar denomination: 'Dekmallorca'.

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application is co-pending with the following
related application (U.S. Plant application Ser. No. 11/267,
902): Title: *Chrysanthemum* Plant Named 'Dekalgarve'.
Applicant: Cornelis W. Dekker.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Chrysanthemum* plant, botanically known as
Chrysanthemum×*morifolium* and referred to by the name
'Dekmallorca'.

The new *Chrysanthemum* is the product of a planned
breeding program conducted by the Inventor in Hensbroek,
The Netherlands. The objective of the breeding program is
to create new cut *Chrysanthemum* cultivars with interesting
inflorescence forms and attractive floret coloration.

The new *Chrysanthemum* originated from a cross-
pollination made by the Inventor on Dec. 2, 2003, in
Hensbroek, The Netherlands, of a proprietary selection of
Chrysanthemum identified as code number 41222, not
patented, as the female, or seed, parent with a proprietary
Chrysanthemum selection identified as code number
01.5574.02, not patented, as the male, or pollen, parent. The
new *Chrysanthemum* was discovered and selected by the
Inventor as a single plant within the progeny of the stated
cross-pollination in a controlled environment in Hensbroek,
The Netherlands.

Asexual reproduction of the new *Chrysanthemum* by
terminal cuttings in Hensbroek, The Netherlands since April,
2003, has shown that the unique features of this new
Chrysanthemum are stable and reproduced true to type in
successive generations.

2

BRIEF SUMMARY OF THE INVENTION

The cultivar Dekmallorca has not been observed under all
possible environmental conditions. The phenotype may vary
somewhat with variations in environment such as
temperature, daylength and light intensity, without,
however, any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of 'Dekmal-
lorca'. These characteristics in combination distinguish
'Dekmallorca' as a new and distinct cultivar:

1. Decorative-type inflorescences with elliptic to oblong-
shaped, pink-colored ray florets.
2. Strong and upright flowering stems.
3. Early and uniform flowering response.
4. Good postproduction longevity.

Plants of the new *Chrysanthemum* can be compared to
plants of the female parent selection. In side-by-side com-
parisons conducted in Hensbroek, The Netherlands, plants
of the new *Chrysanthemum* differed from plants of the
female parent selection in the following characteristics:

1. Plants of the new *Chrysanthemum* had larger inflores-
cences than plants of the female parent selection.
2. Plants of the new *Chrysanthemum* flowered about three
to four days later than plants of the female parent
selection.
3. Plants of the new *Chrysanthemum* and the female
parent selection differed in ray floret color as plants of
the female parent selection had white-colored ray flo-
rets.

Plants of the new *Chrysanthemum* can be compared to
plants of the male parent selection. In side-by-side compari-
sons conducted in Hensbroek, The Netherlands, plants of the
new *Chrysanthemum* differed from plants of the male parent
selection primarily in growth habit as plants of the new
Chrysanthemum were less vigorous than plants of the male
parent selection.

Plants of the new *Chrysanthemum* can be compared to plants of the *Chrysanthemum* cultivar Dekalgarve. In side-by-side comparisons conducted in Hensbroek, The Netherlands, plants of the new *Chrysanthemum* differed primarily from plants of the cultivar Dekalgarve in the following characteristics:

1. Plants of the new *Chrysanthemum* had smaller inflorescences than plants of the cultivar Dekalgarve.
2. Ray florets of plants of the new *Chrysanthemum* were darker pink in color than ray florets of plants of the cultivar Dekalgarve.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the actual colors of the new *Chrysanthemum*.

The photograph on the first sheet comprises a side perspective view of a typical flowering stem of 'Dekmallorca'.

The photograph at the top of the second sheet is a close-up view of a typical inflorescence of 'Dekmallorca'.

The photograph at the bottom of the second sheet is a close-up view of upper and lower surfaces of typical leaves of 'Dekmallorca'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used. The aforementioned photographs and following observations and measurements describe plants grown during the summer in Hensbroek, The Netherlands, under commercial practice in a glass-covered greenhouse. Plants were initially given long day/short night treatments followed by short day/long night treatments to induce flower initiation and development. During the production of the plants, day temperatures ranged from 17.5° C. to 30° C., night temperatures ranged from 18.5° C. to 24° C. and light levels were about five kilolux. Plants were pinched once and were about ten weeks from planting when the photographs and the description were taken.

Botanical classification: *Chrysanthemum*×*morifolium* cultivar Dekmallorca.

Commercial classification: Decorative-type *Chrysanthemum* typically grown as a disbud-type cut flower.

Parentage:

Female or seed parent.—Proprietary selection of *Chrysanthemum*×*morifolium* identified as code number 41222, not patented.

Male or pollen parent.—Proprietary selection of *Chrysanthemum*×*morifolium* identified as code number 01.5574.02, not patented.

Propagation:

Type.—Terminal tip cuttings.

Time to initiate roots, summer.—About 6 days at 20° C.

Time to initiate roots, winter.—About 7 days at 20° C.

Time to produce a rooted cutting, summer.—About 14 days at 20° C.

Time to produce a rooted cutting, winter.—About 16 days at 20° C.

Root description.—Fine; light brown in color.

Rooting habit.—Freely branching.

Plant description:

Appearance.—Herbaceous decorative-type cut *Chrysanthemum*; typically grown as a disbud-type; erect and strong flowering stems. Moderately vigorous growth habit.

Flowering stem description.—Length: About 70 cm to 80 cm. Diameter: About 6 mm. Strength: Strong. Texture: Pubescent. Aspect: Erect. Branching habit: Plants are typically grown as single stems. Color: 146B.

Foliage description.—Arrangement: Alternate, simple. Length: About 7 cm to 13 cm. Width: About 5 cm to 8 cm. Apex: Cuspidate. Base: Attenuate. Margin: Pinnately lobed. Texture, upper and lower surface: Pubescent; rough; leathery. Petiole length: About 1 cm to 2.5 cm. Color: Developing foliage, upper surface: 137A to 137B. Developing foliage, lower surface: Close to 137D. Fully expanded foliage, upper surface: 137A; venation, 147B. Fully expanded foliage, lower surface: Close to 137C; venation, 146B. Petiole, upper and lower surfaces: Close to 147B.

Inflorescence description:

Appearance.—Decorative-type inflorescence form with elliptic to oblong-shaped ray florets. Inflorescences borne on terminals above foliage. Disk and ray florets develop acropetally on a capitulum. Inflorescences not fragrant. Typically grown as a disbud-type.

Flowering response.—Under natural conditions, plant typically flower in November in the Northern Hemisphere. At other times of the year, inflorescence initiation and development can be induced under short day/long night conditions (at least 13.5 hours of darkness). Plants exposed to long day/short night conditions after planting followed by photoinductive short day/long night conditions flower about 7.5 weeks later. Early and uniform flowering response.

Postproduction longevity.—Cut inflorescences will maintain good substance and form for about three weeks.

Quantity of inflorescences per flowering stem.—Grown as a disbud-type, only one inflorescence per stem is allowed to develop. Grown as a spray-type, about 12 to 15 inflorescences develop per flowering stem.

Inflorescence size.—Diameter: About 10 cm. Depth (height): About 3 cm to 4 cm. Diameter of disc: About 1.2 cm.

Inflorescence buds.—Height: About 5 mm to 6 mm. Diameter: About 7 mm to 9 mm. Shape: Oblate. Color: 137D.

Ray florets.—Length: About 2 cm to 4 cm. Width: About 1 cm to 2 cm. Shape: Elliptic to oblong. Apex: Emarginate. Base: Acute; fused. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Number of ray florets per inflorescence: About 300. Color: When opening and fully opened, upper surface: Close to 76C; color becoming closer to 76D with development. When opening and fully opened, lower surface: Close to 76D.

Disc florets.—Shape: Tubular; elongated. Length: About 5 mm. Width: About 1 mm. Number of disc florets per inflorescence: About 10. Color: Immature: 145D. Mature: Apex: 145A. Mid-section: 150B. Base: 145D.

5

Phyllaries.—Shape: Deltoid. Apex: Acute. Base: Truncate. Margin: Entire. Texture, upper surface: Smooth, glabrous. Texture, lower surface: Pubescent. Color, upper surface: Close to 146A. Color, lower surface: Close to 147A to 147B.

Peduncles.—Length: About 3 cm to 4 cm. Diameter: About 3 mm. Angle: Erect. Strength: Moderately strong. Texture: Pubescent. Color: 137C.

Reproductive organs.—Androecium: No anthers observed on disc florets. Gynoecium: Present on both ray and disc florets. Stigma length: About 6 mm to 8 mm. Stigma diameter: About 0.3 mm. Stigma

6

color: Towards the apex, 13A; towards the base, 150C.

Seed/fruit.—Seed and fruit production has not been observed.

Disease/pest resistance: Resistance to known *Chrysanthemum* pathogens and pests has not been observed on plants of the new *Chrysanthemum*.

It is claimed:

1. A new and distinct cultivar of *Chrysanthemum* plant named 'Dekmallorca', as illustrated and described.

* * * * *



