



US00PP17918P2

(12) **United States Plant Patent**  
**Li**

(10) **Patent No.:** **US PP17,918 P2**  
(45) **Date of Patent:** **Aug. 14, 2007**

(54) **VERBENA PLANT NAMED ‘KLEVE04343’**

(50) Latin Name: *Verbena hybrida*  
Varietal Denomination: **KLEVE04343**

(75) Inventor: **Ruijun Li**, North Parramata (AU)

(73) Assignee: **Klemm + Sohn GmbH + Co. KG.**,  
Stuttgart (DE)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/343,882**

(22) Filed: **Jan. 31, 2006**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(52) **U.S. Cl.** ..... **Plt./308**

(58) **Field of Classification Search** ..... **Plt./308**  
See application file for complete search history.

*Primary Examiner*—Anne Marie Grunberg  
*Assistant Examiner*—S. B. McCormick-Ewoldt  
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Verbena* plant named  
‘KLEVE04343’, characterized by its outwardly spreading to  
trailing and mounded plant habit; freely branching habit;  
dense and bushy growth habit; and large bright pink-colored  
flowers with flowers held above and beyond the foliage.

**1 Drawing Sheet**

**1**

Botanical designation: *Verbena hybrida*.  
Cultivar denomination: ‘KLEVE04343’.

**BACKGROUND OF THE INVENTION**

The present Invention relates to a new and distinct culti-  
var of *Verbena* plant, botanically known as *Verbena hybrida*,  
and hereinafter referred by the name ‘KLEVE04343’.

The new *Verbena* is a product of a planned breeding  
program conducted by the Inventor in Camden, New South  
Wales, Australia. The objective of the breeding program is to  
develop new compact *Verbena* cultivars with large and  
attractive flowers.

The new *Verbena* originated from a cross-pollination  
made by the Inventor in 2001 of a proprietary *Verbena*  
*hybrida* selection identified as 01.545.1, not patented, as the  
female, or seed, parent with a proprietary selection of  
*Verbena hybrida* identified as code number 01.603.1, not  
patented, as the male, or pollen, parent. The cultivar  
KLEVE04343 was discovered and selected by the Inventor  
as a flowering plant within the progeny from this cross-  
pollination in a controlled environment in Camden, New  
South Wales, Australia in 2002.

Asexual reproduction of the new cultivar by terminal  
cuttings in a controlled environment in Camden, New South  
Wales, Australia since 2002, has shown that the unique  
features of this new *Verbena* are stable and reproduced true  
to type in successive generations of asexual reproduction.

**SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of  
‘KLEVE04343’. These characteristics in combination dis-  
tinguish ‘KLEVE04343’ as a new and distinct cultivar:

1. Compact and trailing plant habit.
2. Freely branching habit; dense and bushy growth habit.
3. Large bright pink-colored flowers with flowers held  
above and beyond the foliage.

**2**

Plants of the new *Verbena* differ primarily from plants of  
the female parent selection in flower color as plants of the  
female parent selection have lavender-colored flowers. In  
addition, plants of the new *Verbena* and the female parent  
selection differ in leaf shape.

Plants of the new *Verbena* differ primarily from plants of  
the male parent selection in flower color as plants of the  
male parent selection have purple-colored flowers.

Plants of the new *Verbena* can be compared to plants of  
the cultivar Balazdapima, disclosed in U.S. Plant Pat. No.  
17,292. In side-by-side comparisons conducted in Camden,  
New South Wales, Australia, plants of the new *Verbena*  
differed from plants of the cultivar Balazdapima in the  
following characteristics:

1. Plants of the new *Verbena* had larger leaves than plants  
of the cultivar Balazdapima.
2. Plants of the new *Verbena* had larger flowers than plants  
of the cultivar Balazdapima.
3. Plants of the new *Verbena* and the cultivar Balazdapima  
differed in flower color.

**BRIEF DESCRIPTION OF THE PHOTOGRAPH**

The accompanying colored photograph illustrates the  
overall appearance of the new cultivar, showing the colors as  
true as it is reasonably possible to obtain in colored repro-  
ductions of this type. Colors in the photograph may differ  
slightly from the color values cited in the detailed botanical  
description which accurately describe the colors of the new  
*Verbena*. The photograph comprises a side perspective view  
of typical flowering plants of ‘KLEVE04343’.

**DETAILED BOTANICAL DESCRIPTION**

The cultivar KLEVE04343 has not been observed under  
all possible environmental conditions. The phenotype may  
vary somewhat with variations in environment such as  
temperature and light intensity without, however, any vari-  
ance in genotype. The aforementioned photograph and fol-  
lowing observations and measurements describe plants  
grown in Stuttgart, Germany, under commercial practice

during the spring in a glass-covered greenhouse with day temperatures ranging from 18° C. to 20° C., night temperatures ranging from 15° C. to 18° C. and light levels ranging from 20,000 to 55,000 lux. Plants were about five months old when the photograph and description were taken. Plants were pinched one time. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Verbena hybrida* cultivar KLEVE04343.

Parentage:

*Female, or seed, parent.*—Proprietary *Verbena hybrida* selection identified as 01.545.1, not patented.

*Male, or pollen, parent.*—Proprietary *Verbena hybrida* selection identified as code number 01.603.1, not patented.

Propagation:

*Type cutting.*—Terminal cuttings.

*Time to initiate roots, summer.*—About 5 to 8 days at 20° C.

*Time to initiate roots, winter.*—About 8 to 10 days at 20° C.

*Time to produce a rooted cutting or liner, summer.*—About 14 to 20 days at 20° C.

*Time to produce a rooted cutting or liner, winter.*—About 21 days at 20° C.

*Root description.*—Fine, fibrous; pale white in color.

*Rooting habit.*—Medium branching; moderately dense.

Plant description:

*Form.*—Compact and trailing plant habit.

*Growth and branching habit.*—Moderately vigorous and freely-branching with lateral branches potentially developing at every node; dense and bushy growth habit.

*Plant height.*—About 20 cm to 25 cm.

*Plant diameter or spread.*—About 40 cm.

*Lateral branches.*—Length: About 15 cm to 25 cm. Diameter: About 2.5 mm. Internode length: About 2.5 cm to 4 cm. Texture: Slightly pubescent. Strength: Moderately strong. Color: 138B.

*Foliage description.*—Arrangement: Opposite, simple. Length: About 1.5 cm to 3 cm. Width: About 1.2 cm to 2.5 cm. Shape: Deeply dissected. Apex: Acute. Base: Attenuate. Margin: Deeply dissected. Texture, upper and lower surfaces: Slightly pubescent. Venation pattern: Pinnate. Color: Developing and fully expanded foliage, upper surface: 138B; venation, 138C. Developing and fully expanded foliage, lower surface: 138C; venation, 138C. Petiole length: About 5 mm to 10 mm. Petiole diameter: About 1 mm. Petiole texture, upper and lower surfaces: Slightly pubescent. Petiole color, upper and lower surfaces: 138B.

Flower description:

*Flower type and habit.*—Single upright salverform flowers arranged on compact terminal racemes; flowers sessile. Freely flowering with about 12 to 15 flowers per raceme. Inflorescences positioned above and beyond the foliage. Flowers last about one week under greenhouse conditions. Flowers not persistent.

*Fragrance.*—Faint.

*Flowering season.*—In the garden, flowering is continuous from spring through the fall.

*Inflorescence height.*—About 3 cm to 5 cm.

*Inflorescence diameter.*—About 4 cm to 6 cm.

*Flower size.*—Diameter: About 1.5 cm. Depth: About 5 mm.

*Flower buds.*—Length: About 2 mm to 15 mm. Diameter: About 2 mm. Shape: Tubular, columnar. Color: 138C.

*Petals.*—Quantity/arrangement: Five per flower fused at base. Lobe length: About 5 mm to 15 mm. Lobe width: About 5 mm. Shape: Obovate. Apex: Rounded. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; velvety. Color: When opening, upper surface: 63B. When opening, lower surface: 63C. Fully opened, upper surface: 62B; color becoming closer to 70C with development. Fully opened, lower surface: 62C.

*Sepals.*—Quantity/arrangement: Five, fused into a tube. Length: About 1 cm. Diameter: About 1 mm. Shape: Lanceolate. Apex: Acute. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Coarse, pubescent. Color, upper surface: 138B. Color, lower surface: 138C.

*Peduncles.*—Length: About 5 cm. Diameter: About 1.5 mm. Strength: Strong. Color: Close to 138B.

*Reproductive organs.*—Stamens: Quantity per flower: Four, adnate to pistil. Anther shape: Elliptic. Anther length: About 1 mm. Anther color: 1C. Pollen amount: Scarce. Pollen color: 1C. Pistils: Quantity per flower: One. Pistil length: About 1.5 cm. Stigma shape: Spherical. Stigma color: 139C. Style length: About 1.5 cm. Style color: 139D. Ovary color: 139C.

*Fruit/seed.*—Fruit and seed production has not been observed.

Disease/pest resistance: Plants of the new *Verbena* have not been observed to be resistant to pathogens and pests common to *Verbena*.

Temperature tolerance: Plants of the new *Verbena* have been observed to be tolerant to temperatures ranges from about 1° C. to about 35° C.

It is claimed:

1. A new and distinct cultivar of *Verbena* plant named 'Kleve04343', as illustrated and described.

\* \* \* \* \*

