



(12) **United States Plant Patent**
Miedema-Jorna

(10) **Patent No.:** **US PP18,044 P2**
(45) **Date of Patent:** **Sep. 18, 2007**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED
'TAMAR DARK SALMON'**

(50) Latin Name: *Impatiens hawkeri*
Varietal Denomination: **Tamar Dark Salmon**

(75) Inventor: **Anita Miedema-Jorna**, De Lier (NL)

(73) Assignee: **Fides B.V.**, De Lier (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/358,857**

(22) Filed: **Feb. 17, 2006**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./318**

(58) **Field of Classification Search** **Plt./318**
See application file for complete search history.

Primary Examiner—Anne Marie Grunberg
Assistant Examiner—S. B. McCormick-Ewoldt
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of New Guinea *Impatiens* plant
named 'Tamar Dark Salmon', characterized by its upright,
outwardly spreading and uniformly mounded plant growth
habit; freely branching and freely flowering habit; dark
green-colored foliage; and large rounded salmon orange-
colored flowers.

2 Drawing Sheets

1

Botanical designation: *Impatiens hawkeri*.
Cultivar denomination: 'Tamar Dark Salmon'.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of New Guinea *Impatiens* plant, botanically known as
Impatiens hawkeri, and hereinafter referred to by the name
'Tamar Dark Salmon'.

The new *Impatiens* is a product of a planned breeding
program conducted by the Inventor in De Lier, The Neth-
erlands. The objective of the breeding program is to develop
new *Impatiens* cultivars with large rounded flowers and
attractive foliage and flower coloration.

The new *Impatiens* originated from a cross-pollination
made by the Inventor in 2002 with two unnamed proprietary
seedling selections of *Impatiens hawkeri*, not patented. The
cultivar Tamar Dark Salmon was discovered and selected by
the Inventor as a flowering plant within the progeny of the
stated cross-pollination in a controlled environment in De
Lier, The Netherlands.

Asexual reproduction of the new cultivar by terminal
cuttings in De Lier, The Netherlands since 2002, has shown
that the unique features of this new *Impatiens* are stable and
reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and
are determined to be the unique characteristics of 'Tamar
Dark Salmon'. These characteristics in combination distin-
guish 'Tamar Dark Salmon' as a new and distinct *Impatiens*
cultivar:

1. Upright, outwardly spreading and uniformly mounded
plant growth habit.
2. Freely branching and freely flowering habit.
3. Dark green-colored foliage.
4. Large rounded salmon orange-colored flowers.

Plants of the new *Impatiens* can be compared to plants of
the female parent selection. In side-by-side comparisons

2

conducted in De Lier, The Netherlands, plants of the new
Impatiens differed from plants of the female parent selection
in the following characteristics:

1. Plants of the new *Impatiens* flowered earlier than plants
of the female parent selection.
2. Plants of the new *Impatiens* had larger flowers than
plants of the female parent selection.

Plants of the new *Impatiens* can be compared to plants of
the male parent selection. In side-by-side comparisons con-
ducted in De Lier, The Netherlands, plants of the new
Impatiens differed from plants of the male parent selection
in the following characteristics:

1. Plants of the new *Impatiens* had a more uniform plant
habit than plants of the male parent selection.
2. Plants of the new *Impatiens* had larger flowers than
plants of the male parent selection.

Plants of the new *Impatiens* can also be compared to
plants of the cultivar Tamar Salmon, not patented. In side-
by-side comparisons conducted in De Lier, The Netherlands,
plants of the new *Impatiens* differed from plants of the
cultivar Tamar Salmon in the following characteristics:

1. Plants of the new *Impatiens* were more upright than
plants of the cultivar Tamar Salmon.
2. Plants of the new *Impatiens* and the cultivar Tamar
Salmon differed in flower coloration as plants of the
cultivar Tamar Salmon had lighter salmon orange-
colored flowers.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the
overall appearance of the new cultivar, showing the colors as
true as it is reasonably possible to obtain in colored repro-
ductions of this type. Colors in the photographs may differ
slightly from the color values cited in the detailed botanical
description which accurately describe the colors of the new
Impatiens.

The photograph on the first sheet comprises a side per-
spective view of a typical flowering plant of 'Tamar Dark
Salmon' grown in a container.

The photograph on the second sheet is a close-up view of typical flowers of 'Tamar Dark Salmon'.

DETAILED BOTANICAL DESCRIPTION

Plants of the cultivar Tamar Dark Salmon have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype.

The aforementioned photographs, following observations and measurements describe plants grown in De Lier, The Netherlands, under commercial production practice in a glass-covered greenhouse during the summer. Rooted young plants were planted in 12-cm containers and the aforementioned photographs and following observations and measurements were taken about ten weeks later. During the production of the plants, day and night temperatures averaged 20° C. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens hawkeri* cultivar Tamar Dark Salmon.

Parentage:

Female parent.—Unnamed proprietary seedling selection of *Impatiens hawkeri*, not patented.

Male parent.—Unnamed proprietary seedling selection of *Impatiens hawkeri*, not patented.

Propagation:

Type cutting.—Terminal tip cuttings.

Time to initiate roots, summer.—About 5 to 7 days at 19 to 20° C.

Time to initiate roots, winter.—About 6 to 8 days at 19 to 20° C.

Time to produce a rooted young plant, summer.—About 14 to 19 days at 19 to 20° C.

Time to produce a rooted young plant, winter.—About 14 to 21 days at 19 to 20° C.

Root description.—Fine; white to brown in color.

Plant description:

General appearance.—Upright, outwardly spreading and uniformly mounded plant growth habit; freely branching habit; bushy appearance; freely flowering. Moderately vigorous.

Crop time.—From a rooted cutting, about nine weeks are required to produce finished flowering plants in 12-cm containers.

Plant height.—About 23 cm.

Plant diameter or spread.—About 39 cm.

Lateral branches.—Quantity per plant: About seven. Length: About 15 cm. Diameter: About 7 mm. Internode length: About 4 cm. Strength: Moderate. Texture: Smooth, glabrous. Luster: Slightly glossy. Color: 146C tinted with 178A.

Foliage description.—Arrangement: Primarily in whorls with about five leaves per whorl; simple. Length: About 9.6 cm. Width: About 3.5 cm. Shape: Elliptic. Apex: Acute. Base: Attenuate. Margin: Serulate with ciliation. Texture, upper and lower surfaces: Smooth, glabrous. Luster, upper and lower surfaces: Moderately glossy. Venation pattern: Pinnate. Color: Developing foliage, upper surface: 146A to 146B. Developing foliage, lower surface: 139C. Fully expanded foliage, upper surface: Between 139A and 131A; venation, 194B. Fully expanded foliage, lower surface: 138B; towards the base, 194B; venation, similar to lamina. Petiole: Length:

About 3.5 cm. Diameter: About 3 mm. Texture, upper and lower surface: Smooth, glabrous. Color, upper and lower surfaces: 195B.

Flower description:

Flower type and flowering habit.—Single, rounded, salmon orange-colored flowers. Freely and continuously flowering; usually about 20 flowers per lateral branch. Flowers positioned just above the foliage; flowers typically face upright to outward, typically parallel to the leaf canopy. Petals not persistent; gynoecium persistent. Flowers not fragrant.

Flower longevity.—Flowers last about ten days on the plant.

Flowering season.—Year-round under greenhouse conditions. In the garden, plants flower from the spring throughout the summer.

Flower buds.—Length: About 1.4 cm. Diameter: About 6 mm. Shape: Ovoid. Color: Between 41D and 40C.

Flower diameter.—About 4.6 cm.

Flower depth (excluding spur).—About 1.5 cm.

Petals.—Quantity: Five per flower, imbricate. Length: Banner petals: About 2 cm. Lateral and base petals: About 2.4 cm. Width: Banner petal: About 3 cm. Lateral and base petals: About 2.2 cm. Shape: Banner petal: Reniform. Lateral and base petals: Broadly obcordate. Apex: Banner petal: Retuse. Lateral and base petals: Emarginate. Base, all petals: Attenuate. Margin, all petals: Entire. Texture, all petals, upper and lower surfaces: Smooth, glabrous; satiny. Color: When opening, banner petal, upper surface: 41C. When opening, lateral and base petals, upper surface: Mixture of 40C and 41C. When opening, banner petal, lower surface: 43D; towards the base, 52B. When opening, lateral and base petals, lower surface: 43D. Fully opened, banner petal, upper surface: Mixture of 43C and 43D; towards the margins, 41C. Fully opened, lateral and base petals, upper surface: Mixture of 43C to 43D and 41C; central stripe, 41B. Fully opened, banner petal, lower surface: 43D; towards the base, 52B. Fully opened, lateral and base petals, lower surface: 43D.

Spur.—Length: About 4.5 cm. Diameter: About 1.5 cm. Aspect: Curved. Color, immature and mature: Towards the flower, 185A; towards the apex, 155C.

Peduncles.—Length: About 5.5 cm. Diameter: About 1.5 mm. Texture: Smooth, glabrous. Strength: Moderately strong, flexible. Angle: About 45° from vertical. Color: 145B to 145C.

Reproductive organs.—Androecium: Stamen number: Five fused at anthers, hooded; filaments free. Anther length: About 2 mm. Anther shape: Obovate. Anther color: 182D. Pollen amount: Scarce. Pollen color: 158D. Gynoecium: Pistil quantity: One per flower. Pistil length: About 1 mm. Stigma color: 158D. Style: Not visible. Ovary: Five-celled. Ovary color: 146A.

Seed/fruit.—Seed and fruit development has not been observed.

Disease/pest resistance: Plants of the new *Impatiens* have not been observed to be resistant to pathogens and pests common to *Impatiens*.

Temperature tolerance: Plants of the new *Impatiens* have been observed to tolerate temperatures of about 10° C. to about 35° C.

It is claimed:

1. A new and distinct cultivar of New Guinea *Impatiens* plant named 'Tamar Dark Salmon', as illustrated and described.

* * * * *



