

(12) **United States Plant Patent**
Sanders

(10) **Patent No.:** **US PP18,064 P2**
(45) **Date of Patent:** **Sep. 18, 2007**

(54) **IMPATIENS PLANT NAMED ‘IMDOPITWO’**

(50) Latin Name: *Impatiens walleriana*
Varietal Denomination: **Imdopitwo**

(75) Inventor: **Monica Maria Adelheid Sanders,**
Grootebroek (NL)

(73) Assignee: **Syngenta Seeds B.V.,** Enkhuizen (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/440,274**

(22) Filed: **May 24, 2006**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./319**

(58) **Field of Classification Search** Plt./319
See application file for complete search history.

Primary Examiner—Kent Bell

Assistant Examiner—Annette H Para

(74) *Attorney, Agent, or Firm*—Bruce Vrana

(57) **ABSTRACT**

A new *Impatiens* plant, characterized particularly as to
novelty by a strong and upright growing habit, freely
branching, resulting in a big and floriferous plant with large,
double, light pink flowers, flowering above the foliage.

1 Drawing Sheet

1

Latin name of the genus and species of the plant claimed:
Impatiens walleriana.
Varietal denomination: ‘Imdopitwo’.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new distinct cultivar of
Impatiens, botanically known as *Impatiens walleriana*.

The new *Impatiens* is a product of a planned breeding
program conducted in Enkhuizen, Netherlands. The new
Impatiens cultivar has a strong, upright growing type, flo-
riferous with large, double flowers and with attractive foli-
age.

The present cultivar originated from a pollination made in
2002, of a proprietary *Impatiens* selection identified as
‘T2661-2,’ not patented, as the female, or seed parent, with
a proprietary *Impatiens* selection identified as ‘T2661-1,’ not
patented, as the male, or pollen parent.

The present cultivar has been repeatedly asexually repro-
duced by cuttings in Enkhuizen, Netherlands, in Angers,
France, in Gilroy, USA over a two year period. The distinc-
tive characters of this new *Impatiens* are stable and repro-
duced true to type in successive generations of asexual
reproduction. It takes 6 to 8 weeks to produce a finished
plant, depending on the temperature.

This new *Impatiens* plant is an annual in most climatical
zones in the US, only in zones 9 and 10 it is a perennial
plant.

DESCRIPTION OF THE DRAWING

This new *Impatiens* plant is illustrated by the accompa-
nying photographic drawing which shows blooms, buds and
foliage of the plant in full color, the color showing being as
true as can be reasonably obtained by conventional photo-
graphic procedures.

DESCRIPTION OF THE NEW CULTIVAR

The following detailed descriptions set forth the distinc-
tive characteristics of this new *Impatiens*. The data which
defines these characteristics were collected from asexual
reproductions carried out in Enkhuizen, Netherlands. The

2

plant history was taken on 25 week old plants, blossomed
under natural light in the field.

Color readings were taken in the office under ambient
light. Color references are primarily to The R.H.S. Colour
Chart of The Royal Horticultural Society of London.

Plants of the new *Impatiens* differ primarily from the
plants of the female parent selection in the following char-
acteristics:

Plants of the new *Impatiens* have pink flowers with only
petals, whereas the plants of the female have pink flowers
with pollen cap and stigma.

Plants of the new *Impatiens* differ primarily from the male
plant selection in the following characteristics:

Plants of the new *Impatiens* have pink flowers with only
petals, whereas the plants of the male have pink flowers with
pollen cap and stigma.

TABLE 1

**DIFFERENCES BETWEEN THE NEW CULTIVAR
‘IMDOPITWO’ AND A SIMILAR CULTIVAR**

	‘Imdopitwo’	‘Balfieplos’ (U.S. Pat. No. 13,637)
Flower color	Medium pink	Light pink
Leaf color	RHS 147A	RHS 139A
Color of branches	RHS 184A	RHS 146D

Parentage:

Female parent.—‘T2661-2,’ not patented.

Male parent.—‘T2661-1,’ not patented.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—10–12 days at 20 degrees C.

Time to produce a rooted cutting.—4 weeks.

Root descriptions.—Fine, fibrous and white in color.

Plant description:

General appearance.—Compact, mounded and out-
wardly spreading. Appropriate for 12 cm pots and
large containers.

Growth and branching habit.—Strong growing variety
with upright plant habit and large flowers.

Crop time.—6–8 weeks, starting from an unrooted cutting in a 10.5 cm pot.

Plant height.—14.3 cm.

Plant diameter or spread.—41 cm.

Lateral branches:

Length.—About 19.5 cm.

Diameter.—About 4.5 mm.

Internode length.—About 2 cm.

Texture.—Smooth.

Color.—184A.

Foliage description:

Arrangement.—Simple, alternate.

Length.—About 4.9 cm.

Width.—About 2.8 cm.

Shape.—Ovate.

Apex.—Acuminate.

Base.—Attenuate.

Margin.—Serrate, rounded.

Texture.—Glabrous, smooth.

Venation pattern.—Pinnate, arcuate.

Color, developing and fully expanded foliage, upper surface.—147A.

Color, developing and fully expanded foliage, lower surface.—138C.

Venation.—Upper surface: 147D. Lower surface: 147B.

Petiole, length.—About 0.9 cm.

Petiole, diameter.—About 2 mm.

Color.—35C to 35D with green.

Flower description:

Flower type and flowering habit.—Numerous and consistently large fully double flowers. Freely and continuously flowering. Flower buds open similar to a rose in fullness; flowers rounded. Flowers arise from leaf axils. Typically about 23 flowers and flower buds per lateral branch. Flowers positioned above and beyond the foliage and typically face mostly outward. Flowers last about five to seven days under greenhouse conditions. Flowers not persistent. Flowers not fragrant.

Flowering season.—Year round under greenhouse conditions. In the garden, flowering is continuous from Spring until Fall.

Flower size.—Diameter: About 4.6 cm. Depth: About 1.7 cm.

Flower buds:

Length, at stage for showing color.—About 1.3 cm.

Diameter, at stage for showing color.—About 10 mm.

Shape.—Ovoid.

Color.—N66D.

Petals:

Quantity/arrangement.—About 35 per flower; imbricate.

Length, outer petals.—About 2.2 cm.

Width, outer petals.—About 1.9 cm.

Shape.—Oval to obovate.

Apex.—Emarginate to retuse.

Base.—Shortly attenuate.

Margin.—Entire.

Texture, upper and lower surfaces.—Velvety, smooth.

Color.—When opening, upper surface: Center: N66C; Outer: N66D. When opening, lower surface: Center N66C and 65B. Fully opened, upper surface: N66D. Fully opened, lower surface: 65B.

Spurs:

Length.—About 2.1 cm.

Color.—181D.

Sepals:

Quantity/arrangement.—Five in a single whorl.

Calyx length.—About 3 mm.

Calyx diameter.—About 2 mm.

Shape.—Elliptic.

Apex.—Acute.

Margin.—Entire.

Texture, upper and lower surfaces.—Smooth.

Color.—Upper surface: 144B. Lower surface: 144B.

Peduncles:

Length.—About 2 cm.

Diameter.—About 2 mm.

Angle.—About 45 degrees to the lateral branch.

Strength.—Moderately strong.

Color.—166B.

Reproductive organs: None observed.

Seed/fruit development: Seed and fruit development has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Impatiens* plant named 'Imdopitwo,' as substantially illustrated and described herein.

* * * * *

