



US00PP18363P2

(12) **United States Plant Patent**  
**Kobayashi**(10) **Patent No.:** US PP18,363 P2  
(45) **Date of Patent:** Dec. 25, 2007

- (54) **POINSETTIA PLANT NAMED ‘PER11403’**
- (50) Latin Name: *Euphorbia pulcherrima* Willd.  
Varietal Denomination: **PER11403**
- (75) Inventor: **Ruth Kobayashi**, Carlsbad, CA (US)
- (73) Assignee: **Paul Ecke Ranch**, Encinitas, CA (US)
- (\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **11/358,922**
- (22) Filed: **Feb. 17, 2006**
- (51) **Int. Cl.**  
**A01H 5/00** (2006.01)
- (52) **U.S. Cl.** ..... **Plt./307**
- (58) **Field of Classification Search** ..... Plt./307,  
Plt./303

See application file for complete search history.

- (56) **References Cited**
- PUBLICATIONS
- UPOV-ROM GTITM, Plant Variety Database, 2006/04, GTI Jouve Retrieval Software, Citation for *Euphorbia ‘Per11403’* one page.\*  
Ecke Ranch retrieved on Mar. 13, 2007. Retrieved from the Internet <http://www.ecke.com/new1/contact.asp> 3 pages.\*
- \* cited by examiner
- Primary Examiner*—Kent Bell  
*Assistant Examiner*—June Hwu  
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of Poinsettia plant named ‘PER11403’, characterized by its uniform, compact, upright and mounded plant habit; strong stems; dark green-colored leaves; inflorescences with dark red-colored flower bracts; early-season flowering; natural season flower maturity date is late November for plants grown in Encinitas, Calif.; and excellent post-production longevity.

**1 Drawing Sheet****1**

Botanical designation: *Euphorbia pulcherrima* Willd.  
Cultivar denomination: ‘PER11403’.

**BACKGROUND OF THE INVENTION**

The present Invention relates to a new and distinct cultivar of Poinsettia plant, botanically known as *Euphorbia pulcherrima* Willd., and hereinafter referred to by the name ‘PER11403’.

The new Poinsettia is a naturally-occurring whole plant mutation of the *Euphorbia pulcherrima* Willd. cultivar 490, disclosed in U.S. Plant Pat. No. 7,825. The cultivar PER11403 was discovered and selected by the Inventor as a flowering plant within a population of plants of the parent cultivar in a controlled environment in Encinitas, Calif., on Dec. 1, 2003. The new Poinsettia was selected on the basis of its early flowering habit.

Asexual reproduction of the new Poinsettia by terminal cuttings propagated in a controlled environment in Encinitas, Calif., since January, 2003, has shown that the unique features of this new Poinsettia are stable and reproduced true to type in successive generations of asexual reproduction.

**BRIEF SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘PER11403’. These characteristics in combination distinguish ‘PER11403’ as a new and distinct cultivar:

1. Uniform, compact, upright and mounded plant habit.
2. Strong stems.
3. Dark green-colored leaves.
4. Inflorescences with dark red-colored flower bracts.

**2**

5. Early-season flowering; natural season flower maturity date is late November for plants grown in Encinitas, Calif.

6. Excellent post-production longevity.

Plants of the new Poinsettia differ from plants of the parent, the cultivar 490, primarily in time to flower as plants of the cultivar 490 flower about 10 days later than plants of the new Poinsettia.

Plants of the new Poinsettia can be compared to plants of the cultivar Bright Red Freedom, disclosed in U.S. Plant Pat. No. 10,572. In side-by-side comparisons conducted in Encinitas, Calif., plants of the new Poinsettia differed from plants of the cultivar Bright Red Freedom in the following characteristics:

1. Plants of the new Poinsettia flowered about two weeks earlier than plants of the cultivar Bright Red Freedom.
2. Plants of the new Poinsettia had darker red-colored flower bracts than plants of the cultivar Bright Red Freedom.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

25 The accompanying colored photographs illustrate the overall appearance of the new Poinsettia, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Poinsettia.

The photograph at the bottom of the sheet comprises a side perspective view of a typical flowering plant of ‘PER11403’ grown in a container.

35 The photograph at the top left of the sheet is a close-up view of typical inflorescences of ‘PER11403’.

The photograph at the top right of the sheet is a top perspective view of a typical plant of 'PER11403'.

#### DETAILED BOTANICAL DESCRIPTION

The new Poinsettia has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature, daylength and light intensity, without, however, any variance in genotype.

The aforementioned photographs, following observations and averaged measurements describe plants grown in Encinitas, Calif. during the winter under commercial practice in a polyethylene-covered greenhouse with day temperatures averaging about 24° C., night temperatures averaging about 19° C. and light levels about 4,000 foot-candles. Single plants were grown in 16.5-cm pots and pinched once. Plants were flowered under natural season short day/long night conditions. Plants were about 19 weeks from unrooted cuttings when the photographs and the detailed botanical description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

**Botanical classification:** *Euphorbia pulcherrima* Willd. cultivar PER11403.

**Parentage:** Naturally-occurring whole plant mutation of the *Euphorbia pulcherrima* Willd. cultivar 490, disclosed in U.S. Plant Pat. No. 7,825.

#### Propagation:

*Type cutting*.—Terminal cuttings.

*Time to initiate roots*.—About 10 days at 20° C. to 22° C.

*Time to develop roots*.—About four weeks at 20° C. to 22° C.

*Root description*.—Fibrous; white, close to 155D, in color.

*Rooting habit*.—Freely branching.

#### Plant description:

*Growth habit*.—Upright, compact, uniform and mounded plant habit; inverted triangle. Moderately vigorous growth habit.

*Plant height*.—About 23 cm.

*Plant diameter or spread*.—About 44 cm.

*Lateral branch description*.—Quantity: About six lateral branches develop after pinching. Length: About 20 cm. Diameter: About 6 mm. Internode length: About 2.3 cm. Strength: Strong. Texture: Smooth, glabrous. Color: 146B.

*Foliage description*.—Arrangement: Alternate, single. Length: About 12 cm. Width: About 10 cm. Shape: Elliptic. Apex: Acuminate. Base: Acute. Margin: Entire with irregular lobing. Venation pattern: Pinnate. Texture, upper surface: Glabrous, smooth. Texture, lower surface: Slightly pubescent. Color: Developing foliage, upper surface: 147A. Developing foliage, lower surface: 147B. Fully expanded

foliage, upper surface: Darker than 147A; venation, 147A. Fully expanded foliage, lower surface: More gray than 147A; venation, 148A. Petiole: Length: About 8.2 cm. Diameter: About 3 mm. Texture, upper and lower surfaces: Smooth; glabrous. Color, upper and lower surfaces: 187B.

#### Inflorescence description:

*Inflorescence type and habit*.—Inflorescences are compound corymbs of cyathia with colored flower bracts subtending the cyathia. One inflorescence per lateral branch. Flowers are not fragrant. Flowers persistent.

*Natural flowering season*.—Autumn/winter in Northern Hemisphere. Flower initiation and development is induced under long nyctoperiod conditions. Early-season flowering, response time, about seven weeks; natural season flower maturity date is early November for plants grown in Encinitas, Calif.

*Post-production longevity*.—Plants of the new Poinsettia maintain good substance and bract color for about four weeks under interior conditions.

*Inflorescence size*.—Diameter: About 32 cm. Height (depth): About 4 cm.

*flower bracts*.—Quantity per inflorescence: About 15. Length, largest bracts: About 15 cm. Width, largest bracts: About 12 cm. Shape: Elliptic. Apex: Acuminate. Base: Acute. Margin: Entire with occasional shallow lobing. Texture, upper and lower surfaces: Glabrous; velvety. Aspect: Mostly flat to recurving with development. Venation pattern: Pinnate. Color: Developing and transitional bracts, upper surface: 46A. Developing and transitional bracts, lower surface: 53B. Fully expanded bracts, upper surface: 53B. Fully expanded bracts, lower surface: Lighter than 53B. Venation, upper and lower surfaces: Similar to flower bract color. Bract petiole: Length: About 4.5 cm. Diameter: About 2 mm. Texture, upper and lower surfaces: Smooth; glabrous. Color, upper and lower surfaces: 185A.

*Cyathia*.—Quantity per corymb: About four. Diameter of cyathia cluster: About 2 cm. Length: About 1.2 cm. Width: About 6 mm. Shape: Ovoid. Color, immature: 144B. Color, mature: 144A. Nectaries: Quantity per cyathium: About one or two. Size: About 2 mm by 4 mm. Color: 13A. Peduncle: Length: About 2 mm. Diameter: About 1.5 mm. Strength: Strong. Aspect: Mostly upright. Texture: Smooth; glabrous. Color: 144A. Stamens: Quantity per cyathium: About six to ten. Anther shape: Bi-lobed. Anther length: Less than 1 mm. Anther color: 53A. Amount of pollen: Scarce. Pollen color: 13B. Pistils: None observed.

**Disease/pest resistance:** Resistance to pathogens and pests common to Poinsettias has not been observed on plants grown under commercial conditions.

**It is claimed:**

1. A new and distinct cultivar of Poinsettia plant named 'PER11403', as illustrated and described.

\* \* \* \* \*

