



(12) **United States Plant Patent**
Spencer

(10) **Patent No.:** **US PP18,366 P2**
(45) **Date of Patent:** **Dec. 25, 2007**

(54) **PHYGELIUS PLANT NAMED ‘CROSNOQUE’**

(50) Latin Name: *Phygelius aequalis*
Varietal Denomination: **CROSNOQUE**

(76) Inventor: **Malcolm Spencer**, Croftway Hall,
Yapton Road, Barnham, Bognor Regis,
West Sussex (GB), PO22 OBQ

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 22 days.

(21) Appl. No.: **11/494,374**

(22) Filed: **Jul. 26, 2006**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./263**

(58) **Field of Classification Search** **Plt./263**
See application file for complete search history.

Primary Examiner—Kent Bell
Assistant Examiner—S. B. McCormick-Ewoldt

(57) **ABSTRACT**

A new cultivar of *Phygelius* named ‘CROSNOQUE’ that is characterized by compact upright habit, medium-green foliage, and pure-white flowers that bloom late summer and fall. In combination these traits set ‘CROSNOQUE’ apart from all other existing varieties of *Phygelius* known to the inventor.

2 Drawing Sheets

1

Genus: *Phygelius*. Species: *aequalis*.
Denomination ‘CROSNOQUE’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Phygelius*, commonly known as Cape Fuchsia which was bred by the inventor at the inventor’s nursery in West Sussex, United Kingdom. The new cultivar is known botanically as *Phygelius aequalis* and is referred to hereinafter by the cultivar name ‘CROSNOQUE’. ‘CROSNOQUE’ is one of four co-pending applications by the inventor relating to new cultivars of *Phygelius*. The other three co-pending applications are titled ‘CROCOPRI’ (U.S. patent application Ser. No. 11/494,379), ‘CROPURPRI’ (U.S. patent application Ser. No. 11/493,957) and ‘CROYELSOV’ (U.S. patent application Ser. No. 11/494,350). Taken together, these four cultivars constitute the Croftway Series of *Phygelius* and individual plants are grown as annual container plants in most regions of the United States or as perennial plants in regions in U.S.D.A. hardiness zones of 8 or higher.

‘CROSNOQUE’ is a hybrid plant that resulted from a formal breeding program commenced by the inventor in 1997. The breeding program was developed with the objective of producing new varieties of *Phygelius* that exhibit new flower colors. In August 1999 the inventor chose the parents as breeding candidates. The inventor conducted controlled cross-pollination using an individual unnamed seedling of *Phygelius aequalis* as the female parent, and an individual *Phygelius aequalis* ‘Yellow Trumpet’ (unpatented) as the male parent. Cross-pollination was conducted at the inventor’s nursery in West Sussex, United Kingdom in 1999. The seed was collected and sown by the inventor the following February 2000. First flowering occurred, and initial assessment was made in July 2000. Back-crosses were carried out in September, from which new seed was collected and sown in February 2001. The first-flowering and final assessment, were made by the inventor in September 2001.

‘CROSNOQUE’ was selected by the inventor in West Sussex, United Kingdom in 2001. Selection was based on

2

the criteria of flower color and habit. ‘CROSNOQUE’ is distinguishable from the female parent by flower color and pronounced compact upright habit. The female parent exhibits deep-pink flowers, whereas ‘CROSNOQUE’ exhibits pure-white flowers. ‘CROSNOQUE’ is distinguishable from the male parent by flower color and pronounced compact upright habit. The male parent exhibits yellow flowers whereas ‘CROSNOQUE’ exhibits pure-white flowers. No other pure-white flowered comparison *Phygelius* varieties are known to the inventor.

The new *Phygelius* variety named ‘CROSNOQUE’ exhibits compact upright habit, medium-green foliage, and pure-white flowers. The plant dimensions at maturity are 75 cm. in height and 45 cm. in width. ‘CROSNOQUE’ blooms late summer and fall, and is hardy to USDA Zone 8. Cultural conditions include open well-draining soil with good humus content, full sun, and moderate to regular water.

‘CROSNOQUE’ was first asexually propagated by the inventor in 2001 in West Sussex, United Kingdom. The method of asexual propagation used was softwood cuttings. Since that time, under careful observation, the distinguishing characteristics have been determined stable, uniform, and reproduce true to type in successive generations of asexual propagation.

SUMMARY OF THE INVENTION

The following represent the distinguishing characteristics of the new *Phygelius* cultivar named ‘CROSNOQUE’. In combination these traits set ‘CROSNOQUE’ apart from all other existing varieties of *Phygelius* known to the inventor. ‘CROSNOQUE’ has not been tested under all possible conditions and phenotypic differences may be observed with variations in environmental, climatic, and cultural conditions, however, without any variance in genotype.

1. ‘CROSNOQUE’ exhibits compact upright habit.
2. ‘CROSNOQUE’ exhibits pure-white flowers late summer and fall.
3. ‘CROSNOQUE’ exhibits medium-green foliage.

4. 'CROSNOQUE' reaches 75 cm. in height and 45 cm. in width at maturity.
5. Cultural conditions of 'CROSNOQUE' include full sun, open well-draining soil with good humus content, and moderate to regular water.
6. 'CROSNOQUE' is hardy to USDA Zone 8.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying color drawings illustrate the overall appearance of the new *Phygelius* cultivar named 'CROSNOQUE' showing the color as true as it is reasonably possible to obtain in color reproductions of this type. Color in the drawings may differ from the color values cited in the detailed botanical description, which accurately describes the actual color of the new variety 'CROSNOQUE'.

The drawing labeled FIG. 1 depicts a side view of the plant illustrating habit.

The drawing labeled FIG. 2 depicts a close-up view of the flowers. All photographs were taken using conventional techniques and although colors may appear different from actual colors due to light reflectance, they are as accurate as possible by conventional photography.

BOTANICAL DESCRIPTION OF THE PLANT

The following is a detailed botanical description of the *Phygelius* cultivar named 'CROSNOQUE'. Data was collected in Arroyo Grande, Calif. from plants grown out-of-doors in 1-liter containers. Color determinations are in accordance with the 2001 Royal Horticultural Society Colour Chart of London, England except where general color terms of ordinary dictionary significance are used. The growing requirements are similar to the species. The age of the plant described is 4 months.

Botanical classification: *Phygelius aequalis* 'CROSNOQUE'.

Family: Schrophulariaceae.

Genus: *Phygelius*.

Species: *aequalis*.

Denomination: 'CROSNOQUE'.

Common name: Cape fuchsia.

Plant use: Ornamental for use in the landscape.

Parentage: *Phygelius aequalis* 'CROSNOQUE' is a hybrid plant seedling resulting from controlled cross-pollination of the following parents:

Female parent.—An individual unnamed *Phygelius aequalis*.

Male parent.—An individual *Phygelius aequalis* 'Yellow Trumpet'.

Type: Shrub.

Vigor: Vigorous.

Habit: Compact upright.

Dimensions (at one year): 45 cm. in height and 30 cm. in width.

Dimensions (at maturity): 75 cm. in height and 45 cm. in width.

Hardiness: USDA Zone 8.

Asexual propagation: Softwood cuttings.

Root system: Fine and fibrous.

Cultural conditions: Full sun, open well-draining soil with good humus content, and moderate to regular water.

Time to initiate roots: 4–6 weeks.

Crop time: 4–8 months to produce a finished 1-liter container.

Disease susceptibility: Powdery mildew when grown under cover.

Pest susceptibility: Whitefly and greenfly.

Seasonal interest: Flowers late summer and fall.

Stem:

Stem shape.—Angular.

Internode length (average).—4 cm.

Stem width.—0.40 cm. in width.

Stem length.—Ranges from 23–33 cm.

Stem surface.—Glabrous.

Stem color.—138A.

Foliage:

Foliage type.—Evergreen.

Leaf arrangement.—Opposite.

Leaf division.—Simple.

Leaf shape.—Broadly lanceolate.

Leaf base.—Rounded.

Leaf apex.—Acute.

Leaf venation.—Pinnate.

Vein color (adaxial and abaxial surfaces).—147D.

Leaf attachment.—Petiolate.

Petiole dimensions.—1.50 cm. in length and 2 mm. in width.

Petiole shape.—Sulcate.

Petiole color.—138A.

Petiole surface.—Glabrous.

Leaf margin.—Crenate.

Leaf dimensions (average).—4 cm. in length and 2.50 cm. in width.

Leaf color (adaxial surface).—N138A.

Leaf color (abaxial surface).—138A.

Leaf surfaces (adaxial and abaxial).—Glabrous.

Leaf appearance (adaxial surface).—Semi-glossy.

Leaf appearance (abaxial surface).—Matte.

Stipules.—Present.

Stipule color.—N138A.

Stipule apex.—Acute.

Stipule margin.—Crenate.

Stipule dimensions.—4 mm. in length and 2 mm. in width.

Foliar fragrance.—None observed.

Flower:

Inflorescence.—Raceme.

Inflorescence dimensions.—10 cm. in length and 5 cm. in width.

Inflorescence form.—One-sided.

Persistent or self-cleaning.—Self-cleaning.

Peduncle dimensions.—1.25 cm. in length and 2 mm. in diameter.

Peduncle shape.—Cylindroid.

Peduncle surface.—Glabrous.

Peduncle color.—138A.

Pedicel dimensions.—0.75 cm. in length and 1.50 mm. in width.

Pedicel shape.—Cylindroid.

Pedicel surface.—Glabrous.

Pedicel color.—138A.

Bud shape.—Claviform.

Bud surface.—Glabrous.

Bud dimensions.—2.50 cm. in length and 0.50 cm. in width.

Bud apex.—Truncate.

Bud color.—155D.

Flower shape.—Tubular.

Flower quantity.—30–40 flowers per inflorescence.

5

Flower diameter.—1.75 cm.
Flower depth.—4.50 cm.
Flower aspect.—Pendulous.
Corolla tube diameter.—1 cm.
Corolla tube surface.—Glabrous.
Corolla tube color.—155D.
Petals.—5 in number.
Petal dimensions.—5 mm. in length and 5 mm. in width.
Petal color (dorsal surface).—155D.
Petal color (ventral surface).—N144C.
Petals fused or unfused.—Fused.
Petal margin.—Entire.
Petal apex.—Subacute.
Petal surface (abaxial).—Glabrous.
Petal surface (adaxial).—Stipitate glandular.
Calyx dimensions.—0.50 cm. in diameter and 1 cm. in length.
Calyx surface.—Glabrous.
Calyx color.—138A.
Sepals.—Five in number.
Sepal margin.—Entire.
Sepal surface.—Glabrous.
Sepal dimensions.—6 mm. in length and 3 mm. in width.
Fused or unfused.—Sepals are overlapping.
Sepal apex.—Acute.

6

Blooming period.—Late summer and fall.
Fragrance.—None observed.
 Reproduction organs:
Stamens.—4 in number.
Stamen attachment.—Filament adnate to corolla tube.
Stamen color.—155D.
Stamen length.—5 cm.
Anther shape.—Rotund.
Anther dimensions.—2 mm. in width and 3 mm. in length.
Anther color.—160A.
Pollen color.—160A.
Pollen quantity.—Moderate.
Pistil.—One.
Pistil color.—N144C.
Pistil length.—5.50 cm.
Stigma.—Penicillate.
Stigma color.—N144C.
Ovary position.—Superior.
Ovary color.—N144C.
Ovary dimensions.—0.60 cm. in height and 0.40 cm. in width.
Ovary shape.—Ovoid.
 Seed: None observed to date.
 It is claimed:
 1. A new and distinct variety of *Phygelius* plant named ‘CROSNOQUE’ as described and illustrated.

* * * * *



FIG. 1




FIG. 2