



(12) **United States Plant Patent**
Sanders

(10) **Patent No.:** **US PP18,395 P3**
(45) **Date of Patent:** **Jan. 1, 2008**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED
'INGCARREB'**

(50) Latin Name: *Impatiens hawkeri*
Varietal Denomination: **Ingcarrreb**

(75) Inventor: **Monica Maria Adelheid Sanders,**
Grootebroek (NL)

(73) Assignee: **Syngenta Seeds B.V.,** Enkhuizen (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/438,957**

(22) Filed: **May 23, 2006**

(65) **Prior Publication Data**

US 2007/0277275 P1 Nov. 29, 2007

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./318**

(58) **Field of Classification Search** Plt./318
See application file for complete search history.

Primary Examiner—Kent Bell
Assistant Examiner—Annette H Para
(74) *Attorney, Agent, or Firm*—Bruce Vrana

(57) **ABSTRACT**

A new and distinct cultivar of New Guinea *Impatiens* plant
named 'Ingcarrreb' characterized by its good branching habit
with short internodes, medium sized dark green foliage,
large magenta flowers, and good vigor.

1 Drawing Sheet

1

Latin name of the genus and species of the plant claimed:
Impatiens hawkeri.

Varietal denomination: 'Ingcarrreb'.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new and distinct New
Guinea *Impatiens* plant botanically known as *Impatiens*
hawkeri and referred to by the cultivar name 'Ingcarrreb.'

The new cultivar was developed in a controlled breeding
program conducted by the inventor in Enkhuizen, Nether-
lands. The new New Guinea *Impatiens* cultivar is medium
sized with an early flowering response, large flowers with
attractive coloration and a good flowering behavior under
field conditions.

The new cultivar is propagated from cuttings resulting
from the cross in September 2002 of the proprietary New
Guinea *Impatiens* selection identified as 'MM 232-1' as
female parent and the proprietary New Guinea *Impatiens*
selection identified as 'I 150-1' as male parent. 'MM 232-1'
is not commercially available and has not been patented. 'I
150-1' is not commercially available and has not been
patented.

As a result of this cross the present cultivar was selected
in September 2003 in Enkhuizen, Netherlands and has been
repeatedly asexually reproduced by cuttings in Enkhuizen,
Netherlands, in Gilroy, Calif., and in Angers, France over a
period of two years. The distinctive characteristics of this
new *Impatiens* are stable and reproduced true to type in
successive generations of asexual reproduction. It takes 8 to
10 weeks to produce a finished plant, starting from a rooted
plug and planted in a 12 cm pot, depending on the tempera-
ture.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and
are determined to be the unique characteristics of 'Ingcarr-
reb.' These characteristics in combination distinguish 'Ingcarr-
reb' as a new and distinct New Guinea *Impatiens* cultivar:

1. Large magenta flowers with a stable color
2. Ball shaped plant habit with flowers on top of the
foliage

2

3. Dark green foliage
4. Good basal branching character with short internodes
5. Good outdoor vigor

Plants of the new New Guinea *Impatiens* differ primarily
from the plants of the female parent selection in the follow-
ing characteristic:

Plants of the New Guinea *Impatiens* have large magenta
flowers, whereas the plants of the female parent selection
have medium sized, violet flowers.

Plants of the new New Guinea *Impatiens* differ primarily
from the plants of the male parent selection in the following
characteristic:

Plants of the New Guinea *Impatiens* have large magenta
flowers, whereas the plants of the male parent selection have
pink flowers.

Plants of the new New Guinea *Impatiens* can be compared
to plants of 'Moala,' U.S. Pat. No. 10,301. In side-by-side
comparisons conducted in Enkhuizen, Netherlands, plants of
the new New Guinea *Impatiens* differed from plants of the
cultivar 'Moala' in the following characteristics:

1. Plants of the new New Guinea *Impatiens* have medium
sized, dark green leaves, whereas the cultivar 'Moala'
has large, green colored leaves.
2. Plants of the new New Guinea *Impatiens* have a good
branching plant habit with short internodes, whereas
the plant habit of the cultivar 'Moala' is upright grow-
ing.
3. Plants of the new New Guinea *Impatiens* have magenta
flowers with a light red purple spot in the center,
whereas the cultivar 'Moala' has red flowers.

DESCRIPTION OF THE DRAWING

This new New Guinea *Impatiens* plant is illustrated by the
accompanying photographic drawing which shows blooms,
buds and foliage of the plant in full color, the color showing
being as true as can be reasonably obtained by conventional
photographic procedures.

DESCRIPTION OF THE NEW CULTIVAR

The following detailed descriptions set forth the distinctive characteristics of this new New Guinea *Impatiens*. The data which defines these characteristics were collected from asexual reproductions carried out in Enkhuizen, Netherlands. The plant history was taken on 30 week old plants, blossomed under natural light in the field.

Color readings were taken in the laboratory under ambient light. Color references are primarily to The R.H.S. Colour Chart of The Royal Horticultural Society of London.

The plant:

Classification — Botanical.—*Impatiens hawkeri*.

Parentage:

Female parent.—Proprietary *Impatiens hawkeri* selection identified as code number 'MM 232-1,' not patented.

Male parent.—Proprietary *Impatiens hawkeri* selection identified as code number 'I 150-1,' not patented.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—Approximately 7–14 days.

Time to produce a rooted cutting.—Approximately 21 days.

Root description.—Fine, fibrous, and white in color.

Plant description:

General appearance.—Outwardly spreading, low rounded and uniform mounded plant growth.

Habit:

Growth and branching habit.—Freely branching habit, freely flowering, moderate vigorous.

Crop time:

Plant height.—26 cm.

Plant spread.—39 cm.

Lateral branch description:

Length.—14–18 cm.

Diameter.—1–10 mm.

Texture.—Smooth, glabrous.

Internode length.—1–7.5 cm.

Color.—RHS 187A.

Foliage description:

Arrangement.—Primarily in whorls.

Length, mature leaves.—9.0 cm.

Width, mature leaves.—2.2 cm.

Shape.—Lanceolate.

Apex.—Acute to acuminate.

Base.—Acute.

Margin.—Serrulate with fine ciliation.

Texture.—Smooth, glabrous, leathery.

Venation pattern.—Pinnate.

Color young foliage, upper surface.—RHS 147A.

Color young foliage, lower surface.—RHS 187A.

Fully expanded foliage, upper surface.—RHS 147A.

Fully expanded foliage, lower surface.—RHS 197A.

Venation, upper surface.—RHS 187A.

Venation, lower surface.—RHS 187B.

Petiole length.—About 1.5 cm.

Petiole diameter.—About 3.0 mm.

Texture.—Smooth, glabrous.

Color.—RHS 185A.

Flower description:

Flower type and flowering habit.—Single.

Number of flowers per leaf axil.—1.

Number of flowers and flower buds per lateral branch.—6.

Flower position.—Above and beyond the foliage and typically facing upwards and outward.

Flower shape.—Rounded, mostly flat.

Flowers lasting.—About 10 days on the plant depending on environmental conditions.

Petals.—Self cleaning.

Gynoecium.—Persistent.

Flowers.—Not fragrant.

Flowering.—Indeterminate and continuous.

Flowering season.—Year round under greenhouse conditions. In the garden, flowering from spring until fall.

Flower length.—About 6.3 cm.

Flower width.—About 6.3 cm.

Flower depth.—About 0.9 cm.

Flower buds:

Bud length.—About 2.1 cm.

Bud diameter.—About 1.4 cm.

Bud shape.—Ellipsoidal.

Texture.—Smooth, glabrous.

Color, just before opening.—RHS 53C.

Petals:

Quantity.—Single, five per flower, imbricate.

Length of banner petal.—About 3.0 cm.

Width of banner petal.—About 4.2 cm.

Length lateral and basal petals.—About 3.2 cm.

Width lateral and basal petals.—About 3.5 cm.

Lateral and basal petal shape.—Cordate.

Banner petal shape.—Roughly cordate.

Petal apex.—Emarginate.

Petal base.—Attenuate.

Petal margin.—Entire.

Petal texture.—Smooth, satiny.

Petal color.—Upper surface: When opening and fully opened: RHS 53B; spot in center RHS 67C. Lower surface: When opening and fully opened: RHS 53D.

Spur:

Length.—About 4.5 cm.

Texture.—Smooth, glabrous.

Aspect.—Curved.

Color.—RHS 46A.

Peduncles:

Length.—About 4.5 cm.

Strength.—Strong, flexible.

Diameter.—0.2 cm.

Aspect.—About 45° from the stem.

Color.—RHS 46A.

Reproductive organs:

Androecium.—Stamen number: Five fused at anthers, filaments free, hooded.

Anther shape.—Obovate.

Anther length.—About 6 mm.

Anther color.—RHS 56B to 11C.

Amount of pollen.—Moderate.

Pollen color.—RHS 4D.

Gynoecium:

Pistil number.—One.

Pistil length.—About 5 mm.

Stigma shape.—Columnar, five segmented.

Stigma color.—Red purple.

Style color.—RHS 187A.

Ovary arrangement.—Five celled.

Ovary color.—RHS 144A.

Seed development: Seed development has not been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Impatiens* has not been observed.

What is claimed is:

1. A new and distinct cultivar of New Guinea *Impatiens* plant named 'Ingcarreb', as substantially illustrated and described herein.

* * * * *

