



US00PP18463P2

(12) **United States Plant Patent**  
**Misato et al.**(10) **Patent No.:** US PP18,463 P2  
(45) **Date of Patent:** Jan. 29, 2008(54) **ANTIRRHINUM PLANT NAMED  
'SUNKISUHOWA'**(50) Latin Name: *Antirrhinum majus*×*Antirrhinum hispanicum*  
Varietal Denomination: Sunkisuhowa

(75) Inventors: Tomoya Misato, Yamanashi (JP); Kazunari Iwaki, Shiga (JP); Yasunori Yomo, Kanagawa (JP)

(73) Assignee: Suntory Flowers Ltd., Tokyo (JP)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: 11/509,540

(22) Filed: Aug. 24, 2006

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)(52) **U.S. Cl.** ..... **Plt./322**  
(58) **Field of Classification Search** ..... Plt./322  
See application file for complete search history.

*Primary Examiner*—Kent Bell  
*Assistant Examiner*—S. B. McCormick-Ewoldt  
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Antirrhinum* plant named 'Sunkisuhowa', characterized by its upright and mounded plant habit; freely branching habit and short internodes; dense and bushy plant form; numerous white-colored flowers; and long flowering period.

**1 Drawing Sheet****1**

Botanical designation: *Antirrhinum majus*×*Antirrhinum hispanicum*.

Cultivar denomination: 'Sunkisuhowa'.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of *Antirrhinum*, botanically known as *Antirrhinum majus*×*Antirrhinum hispanicum* and hereinafter referred to by the name 'Sunkisuhowa'.

The new *Antirrhinum* is a product of a planned breeding program conducted by the Inventor in Shiga, Japan. The objective of the breeding program is to create new freely flowering potted *Antirrhinums*.

The new *Antirrhinum* originated from a cross-pollination in Shiga, Japan in May, 2001, of a proprietary selection of *Antirrhinum majus*×*Antirrhinum hispanicum* identified as code number His15-6, not patented, as the female, or seed, parent with a proprietary selection of *Antirrhinum majus*×*Antirrhinum hispanicum* identified as code number His15-1, not patented. The cultivar Sunkisuhowa was discovered and selected by the Inventors as a flowering plant from within the progeny of the stated cross-pollination in a controlled environment in Shiga, Japan.

Asexual reproduction of the new *Antirrhinum* by cuttings in a controlled environment in Shiga, Japan since April, 2002, has shown that the unique features of this new *Antirrhinum* are stable and reproduced true to type in successive generations.

**SUMMARY OF THE INVENTION**

The cultivar Sunkisuhowa has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment and cultural practices such as temperature and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Sunkisuhowa'. These characteristics in combination distinguish

**2**

'Sunkisuhowa' as a new and distinct cultivar of *Antirrhinum*:

1. Upright and mounded plant habit.
2. Freely branching habit and short internodes; dense and bushy plant form.
3. Numerous white-colored flowers.
4. Long flowering period.

Plants of the new *Antirrhinum* differ from plants of the female parent selection primarily in flower color as plants of the female parent selection have white-colored flowers with violet-colored venation. In addition, plants of the new *Antirrhinum* have larger flowers than plants of the female parent selection.

Plants of the new *Antirrhinum* differ from plants of the male parent selection primarily in flower size as plants of the new *Antirrhinum* have larger flowers than plants of the male parent selection.

Plants of the new *Antirrhinum* can be compared to plants of the cultivar Festa White, not patented. Plants of the new *Antirrhinum* and the cultivar Festa White differ in the following characteristics:

1. Plants of the new *Antirrhinum* are taller and broader than plants of the cultivar Festa White.
2. Plants of the new *Antirrhinum* have longer and thicker later branches than plants of the cultivar Festa White.
3. Plants of the new *Antirrhinum* have much longer inflorescences than plants of the cultivar Festa White.
4. Plants of the new *Antirrhinum* are more freely flowering than plants of the cultivar Festa White.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying colored photographs illustrate the overall appearance of the new *Antirrhinum*, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Antirrhinum*.

The photograph at the top of the sheet comprises a side perspective view of a typical flowering plant of 'Sunkisuhowa' grown in a container.

The photograph at the bottom of the sheet is a close-up view of a typical inflorescence of 'Sunkisuhowa'.

#### DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown in Shiga, Japan, under commercial practice in a polyethylene-covered greenhouse with day temperatures ranging from 10° C. to 20° C. and night temperatures averaging 0° C. Plants had been growing for about six months in 15-cm containers when the photographs and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used.

**Botanical classification:** *Antirrhinum majus*×*Antirrhinum hispanicum* cultivar Sunkisuhowa.

**Parentage:**

*Female, or seed, parent.*—Proprietary selection of *Antirrhinum majus*×*Antirrhinum hispanicum* identified as code number His 15-6, not patented.

*Male, or pollen, parent.*—Proprietary selection of *Antirrhinum majus*×*Antirrhinum hispanicum* identified as code number His 15-1, not patented.

**Propagation:**

*Type.*—By cuttings.

*Time to initiate roots.*—About 10 days at 20° C. to 25° C.

*Time to produce a rooted young plant.*—About 20 days at 20° C. to 25° C.

*Root description.*—Fine, fibrous and fleshy; white in color.

*Rooting habit.*—Freely branching.

**Plant description:**

*Plant form/habit.*—Upright and mounded plant habit; inverted triangle. Freely branching habit with short internodes; dense and bushy plant form; about nine lateral branches develop per plant; pinching will enhance branching.

*Plant height.*—About 35.2 cm.

*Plant width (spread).*—About 26 cm.

*Lateral branches.*—Length: About 30.8 cm. Diameter: About 3.3 mm. Internode length: About 1 cm. Strength: Moderately strong. Texture: Smooth, glabrous. Color: 144B.

**Foliage description:**

*Arrangement.*—Before flowering, opposite; after flowering, alternate; simple.

*Length.*—About 2.9 cm.

*Width.*—About 1 cm.

*Shape.*—Elliptic.

*Apex.*—Acute to obtuse.

*Base.*—Cuneate.

*Margin.*—Entire.

*Texture, upper and lower surfaces.*—Smooth, glabrous.

*Venation pattern.*—Pinnate; reticulate.

*Color.*—Developing and fully expanded leaves, upper surface: 144A; venation, 144C. Developing and fully expanded leaves, lower surface: 144B; venation, 144C.

*Petiole.*—Length: About 3.8 mm. Diameter: About 1.1 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: 144B.

**Flower description:**

*Flower type/habit.*—Single bi-labiate flowers arranged in terminal and lateral racemes; flowers face obliquely upright to outward. Freely flowering habit with about 31 flowers per inflorescence and about 11 inflorescences per plant.

*Fragrance.*—None detected.

*Natural flowering season.*—Continuously flowering from early autumn until early summer in Japan. Flowers not persistent.

*Postproduction longevity.*—Flowers last about five days on the plant.

*Flower buds.*—Height: About 2.6 cm. Diameter: About 1.2 cm. Shape: Obovate. Color: 145C.

*Inflorescence height.*—About 20.2 cm.

*Inflorescence diameter.*—About 6.2 cm.

*Flower diameter.*—About 3.8 cm by 3.4 cm.

*Flower depth.*—About 3.3 cm.

*Petals.*—Quantity per flower: Upper lip, two-lobed; lower lip, three-lobed; petals fused at the base. Upper lip: Length: About 2.2 cm. Width: About 3.1 cm. Shape: Broadly obovate. Apex: Rounded. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; velvety. Color: When opening, upper and lower surfaces: 155C. Fully opened, upper and lower surfaces: 155C. Lower lip: Length: About 1.6 cm. Width: About 3.3 cm. Shape: Obovate. Apex: Rounded. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; velvety. Color: When opening, upper and lower surfaces: 155C. Fully opened, upper and lower surfaces: 155C. Tube/throat: Length: About 1.6 cm. Diameter: About 1.1 cm. Texture, upper and lower surfaces: Sparsely pubescent. Color: Tube, 155C; throat, 155C.

*Sepals.*—Quantity per flower: Typically five in a single whorl. Length: About 7.8 mm. Width: About 3.4 mm. Shape: Ovate. Apex: Acute. Margin: Entire. Texture, upper surface: Smooth, glabrous. Texture, lower surface: Pubescent. Color, upper and lower surfaces: 144A.

*Pedicels.*—Length: About 8.2 mm. Diameter: About 1.2 mm. Strength: Strong. Texture: Smooth, glabrous. Color: 144A.

*Reproductive organs.*—Stamens: Quantity per flower: Typically four. Anther size: About 1.7 mm by 1.7 mm. Anther shape: Oval, bi-lobed. Anther color: 19A. Filament color: 155D. Pollen amount: Moderate. Pollen color: 9A. Pistils: Quantity per flower: One. Pistil length: About 1.7 cm. Stigma shape: Club-shaped. Stigma color: 145B. Style length: About 1.6 cm. Style color: 145C. Ovary color: 145B.

*Seed/fruit.*—Seed and fruit development have not been observed.

**Disease/pest resistance:** Plants of the new *Antirrhinum* have not been noted to be resistant to pathogens and pests common to *Antirrhinum*.

**Garden performance:** Plants of the new *Antirrhinum* have been observed to have good garden performance and tolerate rain, wind and temperatures ranging from about -10° C. to about 30° C.

**It is claimed:**

1. A new and distinct *Antirrhinum* plant named 'Sunkisuhowa' as illustrated and described.

\* \* \* \* \*

