



US00PP18510P2

(12) **United States Plant Patent**
Kientzler

(10) **Patent No.:** **US PP18,510 P2**
(45) **Date of Patent:** **Feb. 26, 2008**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED
'KIE283'**

(50) Latin Name: *Impatiens hawkeri*
Varietal Denomination: **KIE283**

(75) Inventor: **Ludwig Kientzler**, Gensingen (DE)

(73) Assignee: **The Paul Ecke Ranch**, Encinitas, CA
(US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/518,584**

(22) Filed: **Sep. 8, 2006**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./318**

(58) **Field of Classification Search** **Plt./318**
See application file for complete search history.

Primary Examiner—Kent Bell

Assistant Examiner—June Hwu

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of New Guinea *Impatiens* plant
named 'KIE283', characterized by its upright, outwardly
spreading and mounded plant habit; freely branching habit;
dense and bushy growth habit; dark green-colored leaves;
freely flowering habit; and large red and red purple-colored
flowers that are positioned above and beyond the foliage.

1 Drawing Sheet

1

Botanical designation: *Impatiens hawkeri*.
Cultivar denomination: 'KIE283'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of New Guinea *Impatiens* plant, botanically known as *Impa-
tiens hawkeri*, and hereinafter referred to by the name
'KIE283'.

The new *Impatiens* is a product of a planned breeding
program conducted by the Inventor in Gensingen, Germany.
The objective of the breeding program was to develop new
Impatiens cultivars with numerous large flowers and attrac-
tive foliage and flower coloration.

The new *Impatiens* originated from a cross-pollination
made by the Inventor in October, 2000, of the *Impatiens*
hawkeri cultivar Antigua, disclosed in U.S. Plant Pat. No.
8,283, as the female, or seed, parent with a proprietary
selection of *Impatiens hawkeri* identified as code number
00-421, not patented, as the male, or pollen, parent. The
cultivar KIE283 was discovered and selected by the Inventor
as a flowering plant within the progeny of the stated cross-
pollination in a controlled environment in Gensingen, Ger-
many in April, 2001.

Asexual reproduction of the new cultivar by terminal
cuttings propagated in a controlled environment in
Gensingen, Germany since May, 2001 has shown that the
unique features of this new *Impatiens* are stable and repro-
duced true to type in successive generations of asexual
reproduction.

SUMMARY OF THE INVENTION

The cultivar KIE283 has not been observed under all
possible environmental conditions. The phenotype may vary
somewhat with variations in environment such as tempera-
ture and light intensity, without, however, any variance in
genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of 'KIE283'.

2

These characteristics in combination distinguish 'KIE283'
as a new and distinct cultivar of New Guinea *Impatiens*:

1. Compact, upright, outwardly spreading and mounded
plant habit.
2. Freely branching habit; dense and bushy growth habit.
3. Dark green-colored leaves.
4. Freely flowering habit.
5. Large red and red purple-colored flowers that are
positioned above and beyond the foliage.

In side-by-side comparisons conducted in Gensingen,
Germany, plants of the new *Impatiens* differed from plants
of the female parent, the cultivar Antigua, in the following
characteristics:

1. Plants of the new *Impatiens* were more vigorous than
plants of the cultivar Antigua.
2. Plants of the new *Impatiens* were more freely flowering
than plants of the cultivar Antigua.
3. Plants of the new *Impatiens* had larger flowers than
plants of the cultivar Antigua.

In side-by-side comparisons conducted in Gensingen,
Germany, plants of the new *Impatiens* differed from plants
of the male parent selection in the following characteristics:

1. Plants of the new *Impatiens* had darker green-colored
leaves than plants of the male parent selection.
2. Plants of the new *Impatiens* flowered earlier than plants
of the male parent selection.

Plants of the new *Impatiens* can be compared to plants of
the New Guinea *Impatiens* cultivar BFP-368 Rose, disclosed
in U.S. Plant Pat. No. 9,532. In side-by-side comparisons
conducted in Gensingen, Germany, plants of the new *Impa-
tiens* differed from plants of the cultivar BFP-368 Rose in
the following characteristics:

1. Plants of the new *Impatiens* were more vigorous than
plants of the cultivar BFP-368 Rose.
2. Plants of the new *Impatiens* had darker green-colored
leaves than plants of the cultivar BFP-368 Rose.
3. Plants of the new *Impatiens* and the cultivar BFP-368
Rose differed in flower color.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall appearance of the new *Impatiens*. The photographs show the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Impatiens*.

The photograph at the bottom of the sheet comprises a side perspective view of a typical flowering plant of 'KIE283' grown in a container.

The photograph at the top of the sheet is a close-up view of a typical flower of 'KIE283'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. The following observations and measurements describe plants grown in Encinitas, Calif. during the winter and early spring in a polyethylene-covered greenhouse and under conditions and practices which approximate those generally used in commercial New Guinea *Impatiens* production. During the production of the plants, day temperatures averaged 24° C., night temperatures averaged 19° C. and light levels averaged 4,000 foot candles. Measurements and numerical values represent averages for typical flowering plants. Single plants were grown in one-gallon containers and were about 18 weeks old when the photographs and the detailed description were taken.

Botanical classification: *Impatiens hawkeri* cultivar KIE283.
Parentage:

Female, or seed, parent.—*Impatiens hawkeri* cultivar Antigua, disclosed in U.S. Plant Pat. No. 8,283.

Male, or pollen, parent.—Proprietary selection of *Impatiens hawkeri* identified as code number 00-421, not patented.

Propagation:

Type.—Vegetative cuttings.

Time to initiate roots, summer.—About 14 days at 23° C.

Time to initiate roots, winter.—About 17 days at 20° C.

Time to produce a rooted young plant, summer.—About 21 days at 23° C.

Time to produce a rooted young plant, winter.—About 24 days at 20° C.

Root description.—Fine, fibrous; white to brown in color.

Rooting habit.—Freely branching.

Plant description:

Plant form.—Compact, upright, outwardly spreading and mounded plant habit.

Growth and branching habit.—Vigorous and freely branching habit; about 16 lateral branches developing at the base; dense and bushy growth habit. Pinching, that is, removal of the terminal apices, is typically not required, but will enhance branching.

Plant height.—About 16 cm.

Plant diameter or spread.—About 30 cm.

Lateral branches.—Length: About 11 cm. Diameter: About 8 mm. Internode length: About 2.7 cm. Texture: Smooth, glabrous. Color: 183A.

Foliage description:

Arrangement.—Opposite or in whorls, simple.

Length.—About 8.8 cm.

Width.—About 3.2 cm.

Shape.—Elliptical.

Apex.—Acuminate.

Base.—Attenuate.

Margin.—Serrate with ciliation.

Texture, upper and lower surfaces.—Smooth, glabrous.

Venation pattern.—Pinnate.

Color.—Developing foliage, upper surface: 147A.

Developing foliage, lower surface: 183B. Fully expanded foliage, upper surface: 147A; venation, 184A. Fully expanded foliage, lower surface: Slightly more grey than 183A; venation, 183B.

Petiole.—Length: About 1.7 cm. Diameter: About 3 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: 184A.

Flower description:

Flower type and flowering habit.—Single axillary flowers. Freely flowering habit; usually about six to seven open flowers and flower buds per lateral branch. Flowers positioned above the foliage and typically face upright or outward. Flowers last about one week under greenhouse conditions. Petals self-cleaning, gynoecium persistent. Flowers not fragrant.

Natural flowering season.—Year-round under greenhouse conditions. In the garden, flowering from spring until fall in Germany. Plants begin flowering about eight to twelve weeks after planting.

Flower size.—Length: About 6.4 cm. Width: About 6.4 cm. Depth: About 2.4 cm.

Flower buds.—Length: About 2.5 cm. Diameter: About 1.2 cm. Shape: Ovoid; pointed. Color: 53C.

Petals.—Quantity/arrangement: Five per flower in a single whorl. Length, banner petal: About 2.6 cm. Length, lateral petals: About 3.1 cm. Length, lower petals: About 3.4 cm. Width, banner petal: About 4.8 cm. Width, lateral petals: About 3.5 cm. Width, lower petals: About 3.8 cm. Shape: Cordate. Apex: Emarginate. Base: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening, upper surface: More red than 67A; center, 71C. When opening, lower surface: 53C. Fully opened, upper surface: Brighter than 53C; towards the base, 61B; at base, 53A. Fully opened, lower surface: 53C.

Sepals.—Quantity/arrangement: Three; one modified into an elongated spur. Length: About 1.5 cm. Width: About 6 mm. Shape: Elliptical. Apex: Acuminate. Base: Obtuse. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper surface: 72B to 72C. Color, lower surface: 70B to 70C. Spur length: About 5.3 cm. Spur diameter: At flower, about 5 mm; at apex, less than 1 mm. Spur texture: Smooth, glabrous. Spur color: 59B.

Peduncles.—Length: About 4.6 cm. Diameter: About 2 mm. Angle: About 30° from vertical. Strength: Strong. Texture: Smooth, glabrous. Color: 182A.

Reproductive organs.—Stamens: Quantity: Five fused at anthers; filaments free. Anther size: About 3 mm by 6 mm. Anther color: 10C. Pollen amount: Moderate. Pollen color: 10D. Pistils: Quantity per flower: One. Pistil length: About 5 mm. Stigma shape: Rounded. Stigma color: 61A. Style color: 145A. Ovary color: 145A.

Seed/fruit.—Seed and fruit production have not been observed.

5

Disease/pest resistance: Plants of the new *Impatiens* have not been observed to be resistant to pathogens and pests common to New Guinea *Impatiens*.

Temperature tolerance: Plants of the new *Impatiens* have been observed to tolerate temperatures from about 16° C. to about 30° C.

6

It is claimed:

1. A new and distinct New Guinea *Impatiens* plant named ‘KIE283’ as illustrated and described.

* * * * *

