



US00PP18555P2

(12) **United States Plant Patent**  
**Trees**(10) **Patent No.:** US PP18,555 P2  
(45) **Date of Patent:** Mar. 4, 2008(54) **LANTANA PLANT NAMED  
'BALANDROGLIM'**(50) Latin Name: ***Lantana camara***  
Varietal Denomination: **Balandroglim**(75) Inventor: **Scott C. Trees**, Shell Beach, CA (US)(73) Assignee: **Ball Horticultural Company**, West  
Chicago, IL (US)(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 16 days.(21) Appl. No.: **11/594,392**(22) Filed: **Nov. 8, 2006**(51) **Int. Cl.**  
**A01H 5/00** (2006.01)(52) **U.S. Cl.** ..... **Plt./227**(58) **Field of Classification Search** ..... Plt./227  
See application file for complete search history.

(56)

**References Cited**

## U.S. PATENT DOCUMENTS

PP10,156 P \* 12/1997 Roberson ..... Plt./227  
PP15,246 P2 \* 10/2004 Trees ..... Plt./227

## OTHER PUBLICATIONS

UPOV ROM GTITM Computer Database, GTI Jouve  
Retrieval Software 2007/02 Citation for 'BALANDRO-  
GLIM'.\*

\* cited by examiner

*Primary Examiner*—Wendy Haas(74) *Attorney, Agent, or Firm*—Audrey Charles(57) **ABSTRACT**A new and distinct cultivar of *Lantana* plant named  
'Balandroglim', characterized by its rose-colored flowers,  
dark green-colored foliage, and moderately vigorous, semi-  
upright growth habit.**1 Drawing Sheet****1**Latin name of genus and species of plant claimed: *Lan-  
tana camara*.

Variety denomination: 'Balandroglim'.

## BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar  
of *Lantana* plant botanically known as *Lantana camara* and  
hereinafter referred to by the cultivar name 'Balandroglim'.The new cultivar originated in a controlled breeding  
program in Arroyo Grande, Calif. during August 2002. The  
objective of the breeding program was the development of  
*Lantana* cultivars with continuous flowering, dark green-  
colored foliage, and as well-branched, vigorous growth  
habit.The new *Lantana* cultivar is the result of cross-  
pollination. The female (seed) parent of the new cultivar is  
the proprietary *Lantana camara* breeding selection designated  
484-1, not patented, characterized by its medium  
yellow-colored flowers, dark green-colored foliage, and  
compact, well-branched growth habit. The male (pollen)  
parent of the new cultivar is PATRIOT Classic Desert Sunset  
'Robpatdes', U.S. Plant Pat. No. 10,156, characterized by its  
red yellow-colored flowers, dark green-colored foliage, and  
open growth habit. The new cultivar was discovered and  
selected as a single flowering plant within the progeny of the  
above stated cross-pollination during May 2003 in a con-  
trolled environment at Arroyo Grande, Calif.Asexual reproduction of the new cultivar by terminal stem  
cuttings since May 2003 at Arroyo Grande, Calif. and West  
Chicago, Ill. has demonstrated that the new cultivar repro-  
duces true to type with all of the characteristics, as herein  
described, firmly fixed and retained through successive  
generations of such asexual propagation.**2**

## SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have  
been repeatedly observed and can be used to distinguish  
'Balandroglim' as a new and distinct cultivar of *Lantana*  
plant:

1. Rose-colored flowers;
2. Dark green-colored foliage; and
3. Moderately vigorous, semi-upright growth habit.

Plants of the new cultivar differ from plants of the female  
parent primarily in flower color and growth habit and from  
plants of the male parent primarily in flower color and  
growth habit.Of the many commercially available *Lantana* cultivars  
known to the inventor, the most similar in comparison to the  
new cultivar is LANDMARK Rose Glow 'Balandroglo',  
U.S. Plant Pat. No. 15,246. However, in side by side  
comparisons, plants of the new cultivar differ from plants of  
'Balandroglo' in the following characteristics:

1. Plants of the new cultivar are taller than plants of  
'Balandroglo';
2. Plants of the new cultivar are narrower than plants of  
'Balandroglo'; and
3. Plants of the new cultivar have a flower color different  
from plants of 'Balandroglo'.

## BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it  
is reasonably possible to make the same in color illustrations  
of this type, typical flower and foliage characteristics of the  
new cultivar. Colors in the photographs differ slightly from  
the color values cited in the detailed description, which  
accurately describes the colors of 'Balandroglim'. The

plants were grown in 4.5 inch pots for 17 weeks in a greenhouse at West Chicago, Ill.

FIG. 1 illustrates a side view of the overall growth and flowering habit of 'Balandroglim'.

FIG. 2 illustrates a close-up view of an individual inflorescence of 'Balandroglim' with approximately one-third of the flowers open.

FIG. 3 illustrates a close-up view of an individual inflorescence of 'Balandroglim' with approximately two-thirds of the flowers open.

FIG. 4 illustrates a close-up view of an individual inflorescence of 'Balandroglim' with all of the flowers open.

#### DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible environmental conditions to date. Accordingly, it is possible that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2001 edition, except where general color terms of ordinary significance are used. The color values were determined on Apr. 3, 2006 between 9:00 a.m. and 11:00 a.m. under natural light conditions in West Chicago, Ill.

The following descriptions and measurements describe plants produced from cuttings taken from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown at West Chicago, Ill. in 4.5 inch pots for 17 weeks utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 70° F. to 77° F. (21° C. to 25° C.) during the day and approximately 65° F. to 68° F. (18° C. to 20° C.) during the night. Greenhouse light levels of 2,500 footcandles to 6,000 footcandles were maintained during the day.

Botanical classification: *Lantana camara* cultivar Balandroglim.

Parentage:

*Female parent*.—Proprietary *Lantana camara* breeding selection designated 484-1, not patented.

*Male parent*.—PATRIOT Classic Desert Sunset 'Robpatdes', U.S. Plant Pat. No. 10,156.

Propagation:

*Type cutting*.—Terminal stem.

*Time to initiate roots*.—Approximately 7 to 11 days.

*Time to produce a rooted cutting*.—Approximately 24 to 28 days.

*Root description*.—Fibrous.

*Rooting habit*.—Freely branching.

Plant Description:

*Commercial crop time*.—Approximately 7 to 8 weeks from a rooted cutting.

*Growth habit and generated appearance*.—Moderately vigorous, semi-upright.

*Size*.—Height from soil level to top of plant plane: Approximately 24.8 cm. Width: Approximately 43.8 cm.

*Branching habit*.—Freely branching. Pinching enhances basal branching. Quantity of main branches per plant: Approximately 3.

*Branch*.—Shape: Square in cross section: Strength: Strong, becomes woody with age. Length: Approximate-

mately 29.1 cm. Diameter: Approximately 3.6 mm. Length of central internode: Approximately 4.1 cm. Texture: Densely pubescent with a mixture of glandular and nonglandular hairs. Gland color: Colorless. Color of young stem: 144A. Color of mature stem: 199B.

Foliage description:

*General description*.—Quantity of leaves per main branch: Approximately 15. Fragrance: Strong, spicy. Form: Simple. Arrangement: Opposite.

*Leaves*.—Aspect: Perpendicular to obtuse angle to stem. Shape: Ovate. Margin: Serrate. Apex: Acute. Base: Obtuse. Venation pattern: Pinnate. Length of mature leaf: Approximately 9.0 cm. Width of mature leaf: Approximately 5.2 cm. Texture of upper surface: Scabrous. Texture of lower surface: Scabrous with a mixture of glandular pubescence. Gland color: Colorless. Color of upper surface of young foliage: Closest to 137C with venation of 143C. Color of lower surface of young and mature foliage: 138B with venation of 138C. Color of upper surface of mature foliage: 137A with venation of 143C.

*Petiole*.—Length: Approximately 1.1 cm. Diameter: Approximately 2.0 mm. Texture: Scabrous and glandular pubescent. Gland color: Colorless. Color: 143C.

Flowering description:

*Flowering habit*.—'Balandroglim' is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

*Lastingness of individual inflorescence on the plant*.—Approximately 7 to 8 days from first color of outer buds to dropping of last flower.

Inflorescence description:

*General description*.—Type: Corymb. Quantity per plant: Approximately 7. Fragrance: Light, sweet. Aspect: Facing upward or outward. Height: Approximately 1.9 cm. Width: Approximately 3.6 cm. Quantity of fully open flowers per inflorescence: Approximately 28.

*Peduncle*.—Strength: Strong. Shape: Square in cross section. Aspect: Acute angle to stem. Length: Approximately 1.9 cm. Diameter: Approximately 1.0 mm. Texture: Scabrous and glandular pubescent. Gland color: Colorless. Color: 144B.

Flower description:

*General description*.—Type: Salverform.

*Bud*.—Rate of opening: Generally takes 1 to 2 days for bud to progress from first color to fully open flower. Buds open in progression from the margin to the center of the inflorescence.

*Bud just before opening*.—Shape: Elongated, rectangular at apex. Length: Approximately 9.3 mm. Diameter: Approximately 2.9 mm. Color: 43D.

*Corolla*.—Depth: Approximately 1.4 cm. Diameter: Approximately 1.0 cm.

*Petals*.—Quantity: 4, non-imbricate, non-symmetrical petals. Petals are fused at base forming a corolla tube. Shape: Obovate. Appearance: Dull. Aspect: Flat. Margin: Entire, ruffled. Apex: Obtuse. Length of upper petal from throat: Approximately 4.0 mm. Width of upper petal: Approximately 5.0 mm. Length of lateral petals from throat: Approximately 4.0 mm. Width of lateral petals: Approximately 3.0 mm. Length of lower petal from throat: Approximate-

mately 4.0 mm. Width of lower petal: Approximately 6.0 mm. Texture of upper surface: Glabrous. Texture of lower surface: Sparsely pubescent with a mixture of glandular and nonglandular hairs. Gland color: Mixture of colorless and N74C. Color of upper surface when first open: 4B with 9A at throat opening. Color of lower surface when first open: 4D. Color of upper surface when fully open; N74C with 9A and an overlay of 33A at throat opening. Petals darken with age to N74B with 33A at throat opening. Color of lower surface when fully open: 75C darkening to N74C with age.

*Corolla tube*.—Length: Approximately 1.1 cm. Diameter of tube opening: Approximately 2.0 mm. Diameter at base: Approximately 1.0 mm. Texture of inner surface: Sparsely pubescent. Texture of outer surface: Glabrous at base, gradually becoming densely pubescent at tube opening with a mixture of glandular and nonglandular hairs. Pubescence is a mixture of colorless and 187D. Color of inner surface: 155D. Color of outer surface: 155D with an overlay of 187D.

*Calyx*.—Shape: Tubular with two broadly acute tips. Length: Approximately 3.0 mm. Diameter at tip: Approximately 2.0 mm. Diameter at base: Approximately 1.0 mm. Texture of inner surface: Glabrous. Texture of outer surface: Densely pubescent with a mixture of glandular and nonglandular hairs. Gland color: Colorless. Color of inner and outer surfaces: 142D.

*Bracts*.—Quantity per flower. Shape: Lanceolate. Length: Approximately 5.0 mm. Width: Approximately 1.0 mm. Texture of upper surface: Sparsely pubescent with a mixture of glandular and nonglandular hairs. Gland color: Colorless. Texture of lower surface: Densely pubescent with a mixture of glandular and nonglandular hairs. Gland color: Colorless. Color of upper and lower surfaces: 144B.

*Reproductive organs*.—Androecium: Stamen quantity: 4 per flower, adnate to corolla tube. Stamen length: Approximately 1.0 mm. Anther shape: Ovoid, bilobed. Anther length: Less than 1 mm. Anther color: 9A Pollen amount: Sparse. Pollen color: 9C. Gynoecium: Pistil quantity: 1 per flower. Pistil length: Approximately 5.0 mm. Stigma shape: Funnel. Stigma length: Less than 1 mm. Stigma color: N144D, translucent. Style length: Approximately 4.0 mm. Style color: 155A. Ovary diameter: Approximately 1.0 mm. Ovary color: N144D.

Seed and fruit production: Neither seed nor fruit production has been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Lantana* has not been observed.


What is claimed is:

1. A new and distinct cultivar of *Lantana* plant named ‘Balandroglim’, substantially as herein shown and described.

\* \* \* \* \*



**FIG. 1**



**FIG. 2**

**FIG. 3**

**FIG. 4**