



US00PP18641P2

(12) **United States Plant Patent**
Klemm

(10) **Patent No.:** **US PP18,641 P2**
(45) **Date of Patent:** **Mar. 25, 2008**

(54) **OSTEOSPERMUM PLANT NAMED**
'KLEOE05119'

(50) Latin Name: ***Osteospermum ecklonis***
Varietal Denomination: **KLEOE05119**

(75) Inventor: **Nils Klemm**, Stuttgart (DE)

(73) Assignee: **Klemm + Sohn GmbH&Co. KG**,
Stuttgart (DE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/583,664**

(22) Filed: **Oct. 19, 2006**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./360**
(58) **Field of Classification Search** **Plt./360**
See application file for complete search history.

Primary Examiner—Kent Bell

Assistant Examiner—S. B. McCormick-Ewoldt

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Osteospermum* plant named
'KLEOE05119', characterized by its upright and compact
plant habit; freely branching growth habit; early and freely
flowering habit; and large daisy-type inflorescences with
yellow-colored ray florets.

1 Drawing Sheet

1

Botanical designation: *Osteospermum ecklonis*.
Cultivar denomination: 'KLEOE05119'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Osteospermum* plant, botanically known as *Osteosper-*
mum ecklonis, and hereinafter referred to by the name
'KLEOE05119'.

The new *Osteospermum* is a product of a planned breed-
ing program conducted by the Inventor in Stuttgart, Ger-
many. The objective of the program is to create and develop
new compact *Osteospermum* cultivars with freely branching
and uniformly mounded plant habit, early and freely flow-
ering habit and attractive inflorescence coloration.

The new *Osteospermum* originated from an open-
pollination by the Inventor in 2002 of the *Osteospermum*
ecklonis cultivar Seikimora, disclosed in U.S. Plant Pat. No.
13,409, as the female, or seed, parent with an unknown
selection of *Osteospermum ecklonis* as the male, or pollen,
parent. The new *Osteospermum* was discovered and selected
by the Inventor as a single flowering plant within the
progeny of the stated open-pollination in a controlled envi-
ronment in Stuttgart, Germany in April, 2003.

Asexual reproduction of the new *Osteospermum* by ter-
minal cuttings in a controlled environment in Stuttgart,
Germany since May, 2003, has shown that the unique
features of this new *Osteospermum* are stable and repro-
duced true to type in successive generations.

SUMMARY OF THE INVENTION

The cultivar KLEOE05119 has not been observed under
all possible environmental conditions. The phenotype may
vary somewhat with variations in environment such as
temperature, daylength and light intensity, without,
however, any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of
'KLEOE05119'. These characteristics in combination dis-
tinguish 'KLEOE05119' as a new and distinct cultivar of
Osteospermum.

2

1. Upright and compact plant habit.
2. Freely branching growth habit.
3. Early and freely flowering habit.
4. Large daisy-type inflorescences with yellow-colored
ray florets.

In side-by-side comparisons conducted in Stuttgart,
Germany, plants of the new *Osteospermum* differ primarily
from plants of the female parent, the cultivar Seikimora, in
the following characteristics:

1. Plants of the new *Osteospermum* are larger than plants
of the cultivar Seikimora.
2. Plants of the new *Osteospermum* have broader leaves
than plants of the cultivar Seikimora.
3. Plants of the new *Osteospermum* and the cultivar
Seikimora differ in ray floret color as plants of the
cultivar Seikimora have orange-colored florets.

Plants of the new *Osteospermum* can be compared to
plants of the *Osteospermum* cultivar KLEOE05118, dis-
closed in a U.S. Plant patent application Ser. No. 11/583,
663, filed concurrently. In side-by-side comparisons con-
ducted in Stuttgart, Germany, plants of the new
Osteospermum differed primarily from plants of the cultivar
KLEOE05118 in ray floret color as plants of the cultivar
KLEOE05118 had light orange-colored ray florets. In
addition, plants of the new *Osteospermum* had longer
peduncles and were more freely flowering as plants of the
cultivar KLEOE05118.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying photograph illustrates the overall
appearance of the new *Osteospermum*. This photograph
shows the colors as true as it is reasonably possible to obtain
in colored reproductions of this type. Colors in the photo-
graph may differ slightly from the color values cited in the
detailed botanical description which accurately describe the
colors of the new *Osteospermum*. The photograph comprises
a close-up view of typical flowering plants of
'KLEOE05119'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to
The Royal Horticultural Society Colour Chart, 1995 Edition,

except where general terms of ordinary dictionary significance are used. The aforementioned photograph, following observations and measurements describe plants grown in Stuttgart, Germany during the summer in an outdoor nursery and under conditions and practices which approximate those generally used in commercial *Osteospermum* production. During the production of the plants, day temperatures ranged from 15° C. to 20° C. and night ranged from 10° C. to 15° C. Measurements and numerical values represent averages for typical flowering plants. Plants were about three months old when the photograph and description were taken.

Botanical classification: *Osteospermum ecklonis* cultivar KLEOE05119.

Parentage:

Female, or seed, parent.—*Osteospermum ecklonis* cultivar Seikimora, disclosed in U.S. Plant Pat. No. 13,409.

Male, or pollen, parent.—Unknown selection of *Osteospermum ecklonis*, not patented.

Propagation:

Type.—Terminal vegetative cuttings.

Time to initiate roots, summer.—About ten days at 25° C.

Time to initiate roots, winter.—About two weeks at 18° C. to 20° C.

Time to produce a rooted cutting, summer.—About three to four weeks at 25° C.

Time to produce a rooted cutting, winter.—About four to five weeks at 18° C. to 20° C.

Root description.—Fibrous.

Rooting habit.—Freely branching; dense.

Plant description:

Plant form/growth habit.—Upright and compact plant habit. Inflorescences positioned well above the foliar plane. Moderately vigorous growth habit.

Plant height.—About 14 cm to 16 cm.

Plant diameter.—About 24 cm to 28 cm.

Lateral branches.—Quantity per plant: Freely branching, about three to five lateral branches per plant. Length: About 13 cm to 14 cm. Diameter: About 1 cm. Internode length: About 0.5 cm to 2 cm. Strength: Strong. Texture: Pubescent. Color: 143B.

Foliage description.—Arrangement: Alternate, simple. Length: About 2 cm. Width: About 1.3 cm. Shape: Obovate. Apex: Acute. Base: Acute. Margin: Entire. Texture, upper and lower surfaces: Sparsely pubescent; leathery. Venation pattern: Pinnate. Color: Developing foliage, upper and lower surfaces: 146B. Fully expanded foliage, upper surface: 147A; venation, 147B. Fully expanded foliage, lower surface: 147B; venation, 146B.

Inflorescence description:

Appearance.—Daisy-type inflorescence form with elongated oblong-shaped ray florets. Inflorescences positioned above the foliage, arising from leaf axils. Disc and ray florets developing acropetally on a capitulum. Inflorescences face mostly upright. Freely flowering habit; about 50 inflorescences

develop over time per plant. Inflorescences persistent. Inflorescences not fragrant.

Flowering response.—In Germany, plants of the new *Osteospermum* flower continuously from spring to autumn. Early flowering habit, plants begin flowering about eight to twelve weeks after planting. Inflorescences last about five to ten days on the plant.

Inflorescence bud.—Height: About 1.5 cm. Diameter: About 1 cm. Shape: Ovate. Color: 147B.

Inflorescence size.—Diameter: About 6 cm. Depth (height): About 2 cm. Disc diameter: About 8 mm to 11 mm. Receptacle diameter: About 9 mm. Receptacle height: About 7 mm.

Ray florets.—Shape: Elongate oblong. Length: About 3 cm. Width: About 6 mm. Apex: Acute to emarginate. Base: Acute. Margin: Entire. Texture: Smooth, glabrous; velvety. Orientation: Upright to outwardly. Number of ray florets per inflorescence: About 14 to 17 in a single whorl. Color: When opening, upper surface: 13A; towards the base, 93B. When opening, lower surface: 13A. Fully opened, upper surface: 13B; towards the base, 93B; color becoming closer to 13C with development. Fully opened, lower surface: 13A.

Disc florets.—Shape: Tubular; apex dentate, five-pointed. Length: About 8 mm. Diameter, apex: About 2 mm. Diameter, base: About 1 mm. Number of disc florets per inflorescence: About 20 to 30. Color: Immature: 93A. Mature: Apex: 93A. Mid-section and base: 1C.

Phyllaries.—Quantity per inflorescence: About 14. Length: About 1.3 cm. Width: About 3 mm. Shape: Lanceolate. Apex: Acuminate. Base: Truncate. Margin: Entire. Texture, upper surface: Smooth, glabrous. Texture, lower surface: Pubescent. Color, upper and lower surfaces: 147B.

Peduncles.—Length: About 11 cm to 13 cm. Diameter: About 2 mm to 4 mm. Strength: Strong. Aspect: Mostly upright. Texture: Pubescent; rough. Color: 146C.

Reproductive organs.—Androecium: Present on disc florets only. Anther shape: Lanceolate. Anther length: About 2 mm. Anther color: 79B. Pollen amount: Abundant. Pollen color: 23A. Gynoecium: Present on both ray and disc florets. Pistil length: About 4 mm. Stigma shape: Lanceolate. Stigma color: Close to 143D. Style length: About 2 mm. Style color: Close to 83A.

Seeds.—Length: About 6 mm to 8 mm. Diameter: About 4 mm to 6 mm. Color: 200A.

Disease/pest resistance: Plants of the new *Osteospermum* have not been shown to be resistant to pathogens and pests common to *Osteospermums*.

Temperature tolerance: Plants of the new *Osteospermum* have been observed to tolerate temperatures ranging from about 0° C. to about 35° C.

It is claimed:

1. A new and distinct *Osteospermum* plant named 'KLEOE05119' as illustrated and described.

* * * * *

